

## On Weak Stability Of Iteration Procedures For Some Multi-valued Contractive Maps In Metric space

Amal M. Hashim

[amalmhashim@yhoo.com](mailto:amalmhashim@yhoo.com)

Shereen J. Abbas

Department of Mathematics, College of Science,  
University of Basrah, Basrah- Iraq

### Abstract :

In this paper we prove  $w^2$ -stability for certain class of multi-valued maps that satisfy some general contractive condition in generalized Hausdorff metric space. Some well known results are also derived as a special cases.

**Keywords:** Coincidence and Fixed points ,Weak Stable iteration,  $w^2$ -stability.

حول الاستقرار الضعيف للعمليات المكررة لبعض دوال الانكماش  
ذات القيم المتعددة في الفضاء المترى  
أمل محمد هاشم البطاط  
شيرين عباس  
قسم الرياضيات / كلية العلوم / جامعة البصرة

المخلص

في هذا البحث برهنا  $w^2$ -stability لفئة معينة من الدوال ذات القيم المتعددة التي تحقق شرط التقليل العام في فضاء هاوزدورف المعمم المترى اشتقت بعض النتائج المعروفة كنتائج خاصة

### 1- Introduction

Brined [1] introduced a weaker concept of stability, called weak stability. Timis [10] established weaker notion, named  $w^2$ -stability because of the restriction of an approximation sequence. Some fixed point iteration are not weakly stable so it is used a weaker type sequence named equivalent sequence and gave weak stability results of Picard iteration for various contractive maps.

However, a formal definition of stability of general iterative procedures has been studied by Harder in her Ph. D. Thesis [5] and published in the papers [3] and [4], but the concept of stability is not very precise because of the sequence  $\{y_n\}$  which is arbitrary taken.

In this paper, we give  $w^2$ -stability results of picard iteration for multi-valued maps with coincidence points satisfying some contractive type mappings. Since the stability results for multi-valued contractions have been found useful in the area of generalized differential equations and other contexts (see, for instance [2], [6], [11] and [12]).

## 2- Preliminaries

Throughout the paper, let  $(X, d)$  be a metric space. We shall follow the following notions and definitions.

$CL(X) = \{A : A \text{ is non empty closed subset of } X\}$ . We consider

$$H(A, B) = \max\{\sup d(a, B) ; a \in A, \\ \sup d(A, b) ; b \in B\},$$

For all  $A, B \in CL(X)$  and  $d(a, B) = \inf\{d(a, b) , b \in B\}$ .

$H$  is called the generalized Hausdorff metric for  $CL(X)$  induced by  $d$ .

In [9] Timis presented some mappings  $T : X \rightarrow X$  satisfying various contractive conditions for which the associated Picard iteration is  $w^2$ -stability.

Their corresponding condition in case of a pair  $(S, T)$  mapping, where  $S : Y \rightarrow X$  single-valued map and  $T : Y \rightarrow CL(X)$  multi-valued map with  $x, y \in Y$  and  $x \neq y$  are in the following form.

$$(1.1) \quad H(Tx, Ty) < m(x, y)$$

Where

$$m(x, y) = \max\{d(Sx, Sy), \\ [d(Sx, Tx) + d(Sy, Ty)] / 2, \\ [d(Sx, Ty) + d(Sy, Tx)] / 2\}$$

$$(1.2) \quad H(Tx, Ty) < M(x, y)$$

Where:

$$M(x, y) = \max\{d(Sx, Sy), \\ d(Sx, Tx), d(Sy, Ty), \\ [d(Sx, Ty) + d(Sy, Tx)] / 2\},$$

$$(1.3) \quad H(Tx, Ty) < N(x, y)$$

$$N(x, y) = \max\{d(Sx, Sy),$$

$$\text{Where: } d(Sx, Tx), d(Sy, Ty) \\ , d(Sx, Ty), d(Sy, Tx)\},$$

$$H(Tx, Ty) < M(x, y) + Ld(Sx, Ty) \quad \text{Where } L \geq 0.$$

We remarked that in case of single-valued maps with  $S = \text{identity map}$  in the metric space  $(X, d)$ .

- i. Condition (1.1) implies (1.2) that is any mapping which satisfies condition (1.1) also satisfies condition (1.2).
- ii. (1.2) implies (1.3) and (1.3), (1.4) are independent, and (1.2) implies (1.4) for more details ( see [7] for instance ).

### Definition 2.1 [1]

Let  $(X, d)$  be a metric space and  $\{x_n\}_{n=1}^{\infty} \subset X$  be given sequence. We shall say that  $\{y_n\}_{n=1}^{\infty} \subset X$  is an approximate sequence of  $\{x_n\}$  if, for any  $k \in \mathbb{N}$  (Natural numbers), there exists  $\eta = \eta(k)$  such that  $d(x_n, y_n) \leq \eta$ , for all  $n \geq k$ .

**Lemma 2.2 [1]**

The sequence  $\{y_n\}$  is an approximate sequence of  $\{x_n\}$  if and only if, there exists a decreasing sequence of positive numbers  $\{\eta_n\}$  converging to  $\eta \geq 0$  such that

$$d(x_n, y_n) \leq \eta_n, \text{ for all } n \geq k.$$

**Definition 2.3 [9]**

Let  $(X, d)$  be a metric space and let  $S : X \rightarrow X$ . two sequence  $\{x_n\}_{n=0}^\infty$  and  $\{y_n\}_{n=0}^\infty$  are called S-equivalent sequences if  $d(Sx_n, Sy_n) \rightarrow 0$ , as  $n \rightarrow \infty$ .

**Lemma 2.4 [8]**

Let  $B \in CL(X)$  and  $a \in X$ , then for any  $b \in B$ ,  $d(a, b) \leq H(a, B)$ .

We remarked that any equivalent sequence is an approximate sequence but the reverse is not true, as it shown in the next example.

**Example 2.5 [10]**

Let  $\{x_n\}_{n=0}^\infty$  to be the sequence with  $x_n = n$ . first we take an equivalent sequence of  $\{x_n\}_{n=0}^\infty$  to be  $\{y_n\}_{n=0}^\infty$ ,  $y_n = n + \frac{1}{n}$ . in this case, we have that

$$d(y_n, x_n) = d(n, n + \frac{1}{n}) = \frac{1}{n} \rightarrow 0, \text{ as } n \rightarrow \infty.$$

Now we take an approximate sequence of  $\{x_n\}_{n=0}^\infty$  to be

$\{y_n\}_{n=0}^\infty$ ,  $y_n = n + \frac{n}{2n+1}$ , Then,

$$\begin{aligned} d(y_n, x_n) &= d\left(n, \frac{n}{2n+1}\right) \\ &= \frac{n}{2n+1} \rightarrow \frac{1}{2} > 0 \end{aligned}$$

as  $n \rightarrow \infty$ .

**Definition 2.6 [13]**

Let  $(X, d)$  be a metric space and  $Y \subseteq X$  let  $S : Y \rightarrow X$ ,  $T : Y \rightarrow CL(X)$  be such that  $TY \subseteq TX$  and  $z$  is a coincidence point of  $S$  and  $T$ , that is  $u = Sz \in Tz$ . for any  $x_0 \in Y$ , let the sequence  $\{Sx_n\}$  be generated by the general procedure,

$$Sx_{n+1} \in f(T, x_n), \quad n = 0, 1, \dots \quad (1.5)$$

Converges to an element  $u \in X$ . let  $\{Sy_n\}$  be an approximate sequence of  $\{Sx_n\}$ , we have that  $H(Sy_{n+1}, f(T, y_n)) = 0$  implies  $\lim_{n \rightarrow \infty} Sy_n = u$ . Then (1.5) is called weakly  $(S, T)$ -stable or weakly stable with respect to  $(S, T)$ .

### 3- Main Results

We shall introduce the following definition.

#### Definition 3.1

Let  $(X, d)$  be a metric space. Let  $S : Y \rightarrow X$  and  $T : Y \rightarrow CL(X)$  be such as  $TY \subseteq SY$  and  $z$  is a coincidence point of  $S$  and  $T$ , that is,  $u = Sz \in Tz$ .

For any  $x_0 \in Y$ , let the sequence  $\{Sx_n\}$  be generated by the general iteration procedure, (1.5) converges to an element  $u \in X$ . let  $\{Sy_n\}_{n=0}^\infty \subset X$  be an equivalent sequence of  $\{Sx_n\}$ , and we have that  $H(Sy_{n+1}, f(T, y_n)) = 0$  implies  $\lim_{n \rightarrow \infty} Sy_n = u$ . then (1.5) is called  $w^2$ -stable with respect to  $(S, T)$ .

#### Theorem 3.2

Let  $(X, d)$  be a metric space and  $Y \subseteq X$ . Let  $T : Y \rightarrow CL(X)$ ,  $S : Y \rightarrow X$  such as  $TY \subseteq SY$  and one of  $SY$  or  $TY$  is a complete subspace of  $X$ . let  $z$  be a coincidence point of  $T$  and  $S$ , that is,  $u = Sz \in Tz$ .

For any  $x_0 \in Y$ , let the sequence  $\{Sx_n\}$  generated by Picard iteration  $Sx_{n+1} \in Tx_n$  converges to  $u$ .

Let  $\{Sy_n\} \subseteq X$  be an equivalent sequence of  $\{Sx_n\}$  and define  $\varepsilon_n = H(Sy_{n+1}, Ty_n)$ ,  $n = 0, 1, \dots$ .

If the pair  $(S, T)$  satisfy condition (1.2), and if  $Tz$  is singleton, then the Picard iteration is  $w^2$ -stable with respect to  $(S, T)$ .

#### Proof :

Consider  $\{Sy_n\}$  to be equivalent sequence of  $\{Sx_n\}$ . Then according to definition 3.1, if  $\lim_{n \rightarrow \infty} H(Sy_{n+1}, Ty_n) = 0$  implies that  $\lim_{n \rightarrow \infty} Sy_n = u$ , then the Picard iteration is  $w^2$ - $(S, T)$  stable.

In order to prove this, we suppose that  $\lim_{n \rightarrow \infty} H(Sy_{n+1}, Ty_n) = 0$ ,

therefore,  $\forall \varepsilon > 0$ ,  $\exists n_0 = n(\varepsilon)$  such that  $H(Sy_{n+1}, Ty_n) < \varepsilon \quad \forall n \geq n_0$

$$\begin{aligned} d(Sy_{n+1}, u) &\leq d(Sy_{n+1}, Sx_{n+1}) + d(Sx_{n+1}, u) \\ &\leq d(Sx_{n+1}, Ty_n) + H(Ty_n, Sy_{n+1}) \\ &\quad + d(Sx_{n+1}, u). \\ &\leq H(Tx_n, Ty_n) + H(Ty_n, Sy_{n+1}) \\ &\quad + d(Sx_{n+1}, u) \\ &\leq \max\{d(Sx_n, Sy_n), d(Sx_n, Tx_n), \\ &\quad d(Sy_n, Ty_n), \frac{1}{2}[d(Sx_n, Ty_n) \\ &\quad + d(Tx_n, Sy_n)]\} + H(Ty_n, Sy_{n+1}) \\ &\quad + d(Sx_{n+1}, u). \end{aligned}$$

From the hypothesis,  $Sx_n \rightarrow u$  and

$Sx_{n+1} \in Tx_n$  we have that

$$d(Sx_n, Tx_n) \leq d(Sx_n, Sx_{n+1}) + d(Sx_{n+1}, Tx_n)$$

$$\leq d(Sx_n, u) + d(u, Sx_{n+1}) + d(Sx_{n+1}, Tx_n) \rightarrow 0$$

Now if  $M(x, y) = d(Sx_n, Tx_n)$  by taking limit we obtain  $d(Sy_{n+1}, u) \rightarrow 0$ .

If  $M(x, y) = d(Sy_n, Ty_n)$ , we have

$$d(Sy_n, Ty_n) \leq d(Sy_n, Sx_n) + d(Sx_n, Sx_{n+1}) \\ + d(Sx_{n+1}, Sy_{n+1}) + d(Sy_{n+1}, Ty_n).$$

From definition 2.3, we have that  $d(Sx_n, Sy_n) \rightarrow 0$  and by taking limit, we obtain

$$d(Sy_{n+1}, u) \rightarrow 0$$

If  $M(x, y) = d(Sx_n, Sy_n)$ , from definition 2.3, we have  $d(Sx_n, Sy_n) \rightarrow 0$  and by taking limit, we obtain  $d(Sy_{n+1}, u) \rightarrow 0$ . If

$$M(x, y) = \frac{1}{2} [d(Sx_n, Ty_n) + d(Sy_n, Tx_n)] \\ \leq \frac{1}{2} [d(Sx_n, Sy_n) + d(Sy_n, Ty_n) \\ + d(Sy_n, Tx_n)]$$

Taking limit, we obtain  $d(Sy_{n+1}, u) \rightarrow 0$

Hence  $\lim_{n \rightarrow \infty} Sy_n = 0$

This complete the proof of the theorem.

**Theorem 3.3**

Let  $(X, d)$  be a metric space and  $Y \subseteq X$ . Let  $T : Y \rightarrow CL(X)$ ,  $S : Y \rightarrow X$  such as  $TY \subseteq SY$  and one of  $SY$  or  $TY$  is a complete subspace of  $X$ . Let  $z$  be a coincidence point of  $T$  and  $S$ , that is,  $u = Sz \in Tz$ .

For any  $x_0 \in Y$ , let the sequence  $\{Sx_n\}$  generated by Picard iteration  $Sx_{n+1} \in Tx_n$  converges to  $u$ .

Let  $\{Sy_n\} \subseteq X$  be an equivalent sequence of  $\{Sx_n\}$  and define  $\varepsilon_n = H(Sy_{n+1}, Ty_n)$ ,  $n = 0, 1, \dots$ .

If the pair  $(S, T)$  satisfy condition (1.4), and if  $Tz$  is singleton, then the Picard iteration is  $w^2$ -stable with respect to  $(S, T)$ .

The following example show that  $(S, T)$  is not stable but weakly stable and hence  $w^2$ -stable with respect to  $(S, T)$ .

**Example 3.4**

Let  $X = [0, 1]$  and  $T : X \rightarrow X$ ,  $S : X \rightarrow X$  such as  $TX \subseteq SX$  and

$$TX = \begin{cases} \{0\}, & \text{if } x \in \left[0, \frac{1}{2}\right] \\ \left\{\frac{1}{2}\right\}, & \text{if } x \in \left(\frac{1}{2}, 1\right] \end{cases}$$

$$TX = \{0\} \cup \left\{\frac{1}{2}\right\} = \left\{0, \frac{1}{2}\right\} \subseteq SX = X = [0, 1]$$

$$Sx = x$$

Where  $[0,1]$  endowed with the usual metric  $T$  is continuous at each point of  $[0,1]$  except at  $\frac{1}{2}$ .  $T$  has a unique fixed point at 0, i. e,  $0 \in T(0) = \{0\}$ .

$T$  satisfies condition (1.3).

If  $0 \leq x \leq \frac{1}{2}$ ,  $0 \leq y \leq \frac{1}{2}$  and  $x \neq y$ . Then

$$\begin{aligned} H(Tx, Ty) &= 0 < |x - y| \\ &= \max\{|x - y|, |x - Tx|, |y - Ty|\}, \\ &\quad \frac{1}{2}[|x - Ty| + |y - Tx|] \end{aligned}$$

If  $\frac{1}{2} < x \leq 1$  and  $\frac{1}{2} < y \leq 1$  and  $x \neq y$ . then

$$\begin{aligned} H(Tx, Ty) &= 0 < |x - y| = \max\{|x - y|, |x - Tx|, \\ &\quad |y - Ty|, \frac{1}{2}[|x - Ty| + |y - Tx|]\} \end{aligned}$$

If  $0 \leq x \leq \frac{1}{2}$  and  $\frac{1}{2} < y \leq 1$  and  $x \neq y$ . then

$$\begin{aligned} H(Tx, Ty) &= \frac{1}{2} < y = \max\left\{\left|\frac{1}{2} - x\right|, y\right\} \\ &= \max\{|x - Tx|, |y - Ty|\}, \end{aligned}$$

$$H(Tx, Ty) < \max\{|x - y|, |x - Tx|,$$

Thus

$$|y - Ty|, \frac{1}{2}[|x - Ty| + |y - Tx|]\}$$

In order to study the  $(S, T)$ -stability, let  $x_0 \in [0,1]$ ,  $Sx_{n+1} = x_{n+1} \in Tx_n$ , for  $n = 0, 1, \dots$

$$\text{Then, } x_1 \in Tx_0 = \begin{cases} \{0\} & \text{if } x_0 \in \left[0, \frac{1}{2}\right] \\ \left\{\frac{1}{2}\right\} & \text{if } x_0 \in \left(\frac{1}{2}, 1\right] \end{cases}$$

In each case,  $x_2 \in Tx_1 = \{0\}$  and  $x_n = 0$ ,  $\forall n \geq 2$ .

so,  $\lim_{n \rightarrow \infty} Sx_n = \lim_{n \rightarrow \infty} x_n = 0 \in T\{0\}$

To show that the picard iteration is not  $(S, T)$ -stable,

Let  $Sy_n = y_n = \frac{n^2 + 1}{2n^2}$ ,  $n \geq 1$ .

$$\varepsilon_n = H(Sy_{n+1}, Ty_n) = |y_{n+1} - Ty_n|$$

$$\text{Then } \varepsilon_n = \left| \frac{(n+1)^2 + 1}{2(n+1)^2} - \frac{1}{2} \right|,$$

because of  $y_n \geq \frac{1}{2}$ , for  $n \geq 1$ .

Therefore,  $\lim_{n \rightarrow \infty} \varepsilon_n = 0$  but  $\lim_{n \rightarrow \infty} Sy_n = \lim_{n \rightarrow \infty} y_n = \frac{1}{2}$ , so the Picard iteration is not  $(S, T)$ -stable.

In order to show that  $(S, T)$  weak stability. We take an approximate sequence  $\{Sy_n\}$  of  $\{Sx_n\}$ , from lemma (2.3)

$$|Sx_n - Sy_n| = |x_n - y_n| \leq \eta_n, \quad n \geq k.$$

$$-\eta_n \leq x_n - y_n \leq \eta_n$$

$$0 \leq y_n \leq x_n + \eta_n, \quad n \geq k$$

Since  $x_n = 0$  for  $n \geq 2$ ,

$$0 \leq y_n \leq \eta_n, \quad n \geq k_1 = \max\{2, k\}$$

We can choose  $\{\eta_n\}$  such that  $\eta_n \leq \frac{1}{2}$ ,  $n \geq k_1$  and therefore,  $0 \leq y_n \leq \frac{1}{2}$ ,  $\forall n \geq k_1$

So  $Ty_n = \{0\}$  and the results that  $\varepsilon_n = H(Sy_{n+1}, Ty_n) = Sy_{n+1} = y_{n+1}$

Now  $\lim_{n \rightarrow \infty} \varepsilon_n = \lim_{n \rightarrow \infty} y_{n+1} = 0$ ,

So the iteration procedure is weakly stable with respect to  $(S, T)$ .

Hence, it is  $w^2$ -stable with respect to  $(S, T)$ .

### Corollary 3.5 [9]

Let  $(X, d)$  be a complete metric space and  $S, T : X \rightarrow X$  such that  $TX \subseteq SX$  satisfying the following condition :

$$d(Tx, Ty) < \max\{d(Sx, Tx), d(Sy, Ty)\},$$

$$d(Sx, Sy), \frac{1}{2}[d(Sx, Ty) + d(Sy, Tx)]\}, \quad \text{For all } x, y \in X \text{ and } x \neq y.$$

Let  $\{Sy_n\}_{n=0}^{\infty}$  an iteration procedure defined by  $x_0 \in X$  and  $Sx_{n+1} = Tx_n$ , for all  $n \geq 0$  and the sequence  $\{Sx_n\}$  converge to  $u$ , where  $u$  is a coincidence point of  $S$  and  $T$ , then the Picard iteration is  $w^2$ -stable with respect to  $(S, T)$ .

The following example explains the stability, weakly stability and  $w^2$ -stability for some contraction condition in case of single-valued map.

### Example 3.5

Let  $X = [0, 1]$  and let  $T : X \rightarrow X$  be such that  $Tx = \frac{x}{2}$ , where  $X$  has the usual metric, Evidently, every point of  $X$  is a fixed point of  $T$ . let  $x_0 = \frac{1}{2}$ .

Then  $x_{n+1} = Tx_n = T^{n+1}x_0 = \frac{1}{2^{n+1}}$ ,  $n = 0, 1, 2, \dots$ ,

thus  $\lim_{n \rightarrow \infty} x_n = 0$ . let  $\{y_n\}_{n=0}^{\infty}$  be an arbitrary sequence in  $X$  such that  $y_0 = x_0$  and

$$y_0 = \frac{1}{2}, \quad n = 1, 2, 3, \dots$$

Thus  $\lim_{n \rightarrow \infty} |y_{n+1} - Ty_n| = \left| \frac{1}{n+1} - \frac{1}{n} \right| = \frac{1}{n(n+1)} \rightarrow 0$  as  $n \rightarrow \infty$ , however,

$$\lim_{n \rightarrow \infty} y_n = 0 \neq \lim_{n \rightarrow \infty} x_n = \frac{1}{2}.$$

Therefore, the iterative procedure  $x_{n+1} = Tx_n$  is not stable.

Now, if we choose  $\{y_n\}$  an approximate sequence of  $\{x_n\}$  such that  $y_0 = x_0$  and

$$y_n = \frac{2n+1}{2n}. \text{ In this case, we have } d(y_n, x_n) = \frac{n+1}{2n} \rightarrow \frac{1}{2} > 0$$

as  $n \rightarrow \infty$ , however,  $\lim_{n \rightarrow \infty} y_n = \frac{1}{2} = \lim_{n \rightarrow \infty} x_n$ . Therefore, the iterative procedure  $x_{n+1} = Tx_n$  is weakly stable.

Finally, let  $\{y_n\}$  be an equivalent sequence of  $\{x_n\}$  such that  $y_0 = x_0$  and  $y_n = \frac{n+1}{2}$ .

in this case, we have  $d(y_n, x_n) = \frac{1}{2n} \rightarrow 0$  as  $n \rightarrow \infty$ .

$$\text{However, } \lim_{n \rightarrow \infty} y_n = \frac{1}{2} = \lim_{n \rightarrow \infty} x_n = T\left(\frac{1}{2}\right).$$

Therefore, the iterative procedures  $x_{n+1} = Tx_n$ ,  $n = 0, 1, 2, \dots$  is  $w^2$ -stable.

Note that  $T$  is non expansive, that is,  $d(Tx, Ty) \leq d(x, y)$ , for all  $x, y \in X$ .

### Remark 3.7

- i. If  $S$  is the identity mapping in corollary 3.4 we have theorem 2.4 [10].
- ii. Every stable iteration is weakly stable but the reverse may not true (see [11]).
- iii. Every weakly stable iteration is  $w^2$ -stable but the reverse may not true (see [9]).
- iv. There is some mappings that satisfy contraction condition and for which the Picard iteration is not  $(S, T)$ -stable, it is not  $(S, T)$ -weakly stable but it is  $(S, T)$ - $w^2$ stable (see [9]).

### References:

- [1] Berinde, V. Iterative approximation of fixed points. Springer Verlag, lectures notes in Mathematics, 2007.
- [2] Cardinali. T. and Rubbioni. P, A generalization of the Caristi fixed point theorem in metric space, Fixed point theory. 11(2010), 3-10.
- [3] Harder, A. M. and Hicks, T. L. A stable iteration procedure for non expansive mapping. Math. Japon. 33(1998), 687-692.
- [4] Harder, A. M. and Hicks, T. L. Stability results for fixed point iteration procedures. Math. Japon. 33(1988), 693-706.
- [5] Harder, A. M. Fixed point theory and stability results for fixed point iteration procedures. Ph. D. thesis. University of Missouri-Rolla. Missouri. (1987).
- [6] Lim, T. C. , Fixed point stability for set-valued contractive mappings with applications to generalized differential equations. J. Math. Anal. Appl. 110 (1985), 436-441.
- [7] Rhoades, B. E. , A comparison of various definition of contractive mappings, Tran. Amer. Math. Soc. 226 (1977), 257-290.
- [8] Singh S. L. , Chary Bhatnagar and Hashim A. M. Round of stability of picard iterative procedure for multi-valued operators. Non linear Anal. Forum 10(1) (2005), 13-19.
- [9] Timis, I. and Berinde, V. Weak stability of iterative procedures for some coincidence theorems. Creative Math. & Inf. 19 (2010), 85-95.
- [10] Timis, I. On the weak stability of picard iteration for some contractive type mappings and coincidence theorems. Int. J. Com. Appl. 37 (2012), 9-13.



- [11] Timis, I. On the weak stability of picard iteration for some contractive type mappings. Annals of university of Craiova – Mathematics, and computer science series.37 (2) (2010), 106-114.
- [12] Tolostonogo V. A. Differential inclusions in Banach space kluwer Academic publishers, 2000 (2008), 2942-2949.
- [13] Zaineb Sami Almousawy; Fixed point theorems and Multi-valued operators. M. SC. Thesis. University of Basrah (supervised by Dr. Amal M. Hashim) 2012.