

<http://dx.doi.org/10.52113/1/2/2023-227-239>

Overview of urinary tract infection caused by bacteria

Ahmed Talib Abdulkarem¹, Shaimaa M.S. Zainulabdeen², Saif Mazeel Abed³



Abstract

Urinary tract infections [UTIs], a significant public health concern, can be brought on by a wide variety of bacteria, including *Escherichia coli*, *Proteus mirabilis*, *Enterococcus faecalis*, *Klebsiella pneumoniae*, and *Staphylococcus saprophyticus*. The high recurrence rates and evolving antibiotic resistance of uropathogens constitute a serious threat to the financial burden of these disorders. Pathogens in urine can be detected using a variety of diagnostic approaches, which are broadly divided into laboratory-based and point-of-care [POC] detection methods. Many research institutions and companies working in this subject have strived to establish a rapid and accurate pathogen identification because traditional approaches may be time-consuming. The indications and symptoms are further divided into the following three groups: General signs and symptoms like fever; lower urinary tract symptoms including urgency, frequency, and dysuria; and non-specific signs and symptoms such as nausea and malaise. Additionally, the prevalence of UTIs brought on by multidrug resistance [MDR] is rising, which has a considerable negative impact on the propagation of antibiotic resistance as well as the financial burden of these infections.

Keywords: UTI, Iraq, Different diagnostic, Multidrug resistance

* Correspondence author: shimaa.majeed@mu.edu.iq

¹ Ministry of Health, Baghdad, Iraq.

² Department of Microbiology, College of Medicine, Al-Muthanna University.

³ Department of Medical Laboratories, College of Health & Medical Technology, Sawa University.

Received 22 August 2023; revised 12 November 2023; accepted 29 November 2023, available online 19 December 2023.

Copyright © 2023 Zainulabdeen, et al. This is article distributed under the terms of the Creative Commons Attribution License <http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited

Introduction

Every year, 150 million individuals throughout the world suffer from urinary tract infections [UTIs], one of the most prevalent bacterial illnesses¹. According to reports, there were 10.5 million office visits for UTI symptoms in the United States alone in 2007 [0.9% of total ambulatory visits], and 2-3 million emergency room visits., according to Schappert and Rechtsteiner [1]. The same bacteria that cause UTIs in non-pregnant persons also cause UTIs in pregnant people. *E. coli* is the root cause of 70–80% of infections, according to Barr

et al. [2]. Two further typical GNBs are *Proteus mirabilis* and *Klebsiella* sp. [3]. Fastidious pathogens and anaerobic bacteria have been detected in large numbers of urine samples from pregnant women, but it is yet unknown how these germs may impact newborn outcomes. The pressure of the gravid uterus on the bladder, which restricts urine flow, as well as other hormonal and immunological changes that take place throughout a typical pregnancy, raise the risk of UTI in pregnant women [4]. As a result, throughout pregnancy, women should periodically check their urine [5]. Clinically, UTIs can be categorised as basic or complicated. Uncomplicated UTIs most frequently afflict individuals who are in general good health and do not have any anatomical or neurological abnormalities of the urinary system [6]. Pyelonephritis and cystitis are two different classifications of these illnesses. Some of the risk factors associated with cystitis include female gender, a history of UTI, sexual activity, obesity, diabetes vaginal infection, and genetic susceptibility [7]. According to Tien et al. [8], complex UTIs include the following: calculi, indwelling catheters, or other drainage devices; urinary obstruction; urinary retention brought on by neurological conditions; immunosuppression; renal failure; renal transplantation; pregnancy; and diseases that affect the host's capacity to protect the urinary tract. Sofyan et al.'s [9] research, indwelling catheters are to blame for 1 million cases of complex UTIs each year. Catheter-associated urinary tract infections [CAUTIs], which are the most common cause of future bloodstream infections, are linked to higher rates of morbidity and death. According to Rizwan et al. [10], long-term catheterization, being a woman, being older, and having diabetes are all risk factors for developing a CAUTI.

Both Gram-positive and Gram-negative bacteria, as well as some fungi, are responsible for UTIs. The most frequent cause of both simple and complex UTIs is Uropathogenic *Escherichia coli* [UPEC], which is highly widespread. According to Foxman [11], UPEC is followed in prevalence among the bacteria that cause simple UTIs by *K. pneumoniae*, *S. saprophyticus*, *E. faecalis*, group B streptococcus [GBS], *P. mirabilis*, *P. aeruginosa*, *S. aureus*, and *Candida spp*, *Enterococcus spp.* and *staphylococcus spp* are the most prevalent causal agents for complex UTIs, after UPEC, which is the most frequent [12].

Prevalence of UTI in Iraq and other Countries

Salman et al.[13] found that 14.7% of school-age children in Al Imamein Kadhimaein Medical City had urinary tract infections. This rate is lower than that found in a different Iraqi study conducted in Tikrit city in 2012, which found that the culture-positive rate in school-age children was 42.9% [14], but similar to those found in a descriptive cross-sectional study conducted in Iraq by Kareem and Issa [15], which discovered that 18.3% of kids from Basra

city's five hospitals had positive urine cultures, and the findings of a cross-sectional study conducted on 2,511 schoolchildren in Turkey by Zincir et al. [16], which revealed a prevalence of [7.1%], which was lower than the findings of a study carried out in Nigeria by Isa et al. [17], which revealed that 31.7% of primary school-aged kids had positive urine cultures for bacterial growth. There may be differences in socioeconomic level, culture, personal cleanliness, health infrastructure, and school health services among various communities that contribute to these variations in UTI rates among school-age children.. The rate found in the current study is also typical for symptomatic UTI, while the rate of asymptomatic UTI in our sample was 1.6%, which is lower than the results study in Iraq, which found a prevalence of [6.6%] for schoolchildren with an asymptomatic UTI [18]. Hasan and associates [19] discovered in 2022 While tigecycline, amikacin, ciprofloxacin trimethoprim/sulphomethoxizole, levofloxacin, meropenem, and tetracycline showed lower resistance to these illnesses' drug resistance, all gram-negative bacterial isolates were found to be highly resistant to ampicillin and cefotaxime. This is concerning because it limits clinical practitioners' options for treatment and forces them to choose drugs with low resistance ., Nahab et al.[20] found that E. coli was the most prevalent pathogenic bacteria. These results support those E. coli as the uropathogen in charge of UTI[21]. The findings of females with UTI in other Iraqi cities, such as Baghdad [22] and Al-Mosul [23], were supported by this discovery.

Pathogenesis of UTI

UTIs [urinary tract infections], which are brought on by uropathogens that reside in the gut, can travel to the urethra and bladder thanks to a group of adhesins. The bacteria start to multiply and produce poisons and enzymes that help them survive when the host's inflammatory response cannot entirely eradicate all of the bacteria. If the bacterium succeeds to get beyond the kidney epithelial barrier, further colonization of the kidneys could lead to bacteremia. Catheterization in cases of complete UTIs causes uropathogen infection, which is followed by bladder dysfunction. The strong immunological response brought on by catheterization frequently causes fibrinogen to build up on the catheter. Fibrinogen-binding proteins made by uropathogens enable them to attach to the catheter According to McLellan and Hunstad [24], UTIs are the most common bacterial infection in humans worldwide and the most common sickness picked up in a hospital. Infections can progress to pyelonephritis and bacteremia if they are left untreated because bacteria also thrive as a result of biofilm protection. According to Lewis et al. [25], the success of various uropathogen invasion and adhesion techniques is directly related to the spread of UTIs. The infection may not initially appear to be very severe, especially in the early stages, but if

aggravating conditions are present, it may get much worse [25] Biofilms, urinary stasis from blockage, and catheters are complicating variables that contribute to the development of UTI. UTIs are a diverse set of clinical illnesses that differ in their genesis and degree of severity. Numerous inherent and acquired variables, including urine retention, vesicoureteral reflux, frequent sexual activity, prostate gland enlargement, vulvovaginal atrophy, and familial history, might increase the risk of UTI. Women who take spermicides may get more UTIs [26]. Asymptomatic bacteriuria is defined as a urine culture with 10⁵ colony-forming units/mL without any particular UTI symptoms since it typically cures on its own and doesn't need to be treated [27]. Only some individuals, such as those who are pregnant, have neutropenia, or are having genitourinary surgery, should get therapy for asymptomatic UTIs since doing so increases the risk of bacterial resistance developing [28].. Contrarily, antibiotics are frequently used to treat symptomatic UTIs, which increases the likelihood of the development of multidrug-resistant microbes by altering the intestinal and vaginal microbiota [29].

Signs and Symptoms

Most UTI criteria used in the studies that were included required signs and symptoms to be present. It's interesting to note that 15% of research did not clarify which symptoms or indicators had to be present to identify UTIs, and even more studies did not indicate which symptoms or signs had to be present at all. The mainstay of UTI diagnosis continues to be symptoms and indications. By precisely identifying specific symptoms, the risk of classification mistakes may be decreased. In studies including older patients with UTI, symptom characterization is particularly crucial due to the high background prevalence of asymptomatic bacteriuria and pyuria [30]. Classic UTI-associated symptoms like dysuria, frequency, and urgency were commonly mentioned in studies that did not specify which symptoms were included in the UTI criteria. But we also found a wide range of non-specific symptoms, especially in trials that failed to isolate the specific UTI subtype under research. The fact that only a small percentage of the included studies employed the same collection of symptoms to diagnose UTI supports the theory that, despite the ambiguous clinical importance of nonspecific symptoms in UTI, this diversity of symptoms contributes to study heterogeneity. Furthermore, no minimum number of symptoms [for diagnosis] was specified in more than a third of the reports that were included. A minimum number of symptoms should be stated because even the normal lower urinary tract symptoms are not 100% specific for UTIs and because the likelihood of UTI increases when numerous symptoms are present [31].

Types of UTI and Definitions

Adult UTIs can be diagnosed as either pyelonephritis or cystitis. Men and women who have urethritis, a sexually transmitted disease, are included in the differential diagnosis. Males are susceptible to both acute and chronic bacterial prostatitis, which are all parts of the UTI spectrum. Men can suffer from the associated disorders epididymitis, orchitis, and epididymo-orchitis.

A-Uncomplicated cystitis: A healthy, premenopausal female with normal urinary system, non-pregnant has a bladder infection [32].

B-Complicated cystitis: a bladder infection accompanied with elements that either lessen the effectiveness of treatment or raise the danger of unfavorable effects. This includes vesicoureteral reflux, cystitis, a foreign item [such a catheter or urinary tract stone], and recent instrumentation. It also covers cystitis in men, pregnant women, those who have had kidney transplants, and those with various immunocompromised situations. These UTIs are brought on by unique organisms or bacteria that are resistant to numerous medications [33].

C-Uncomplicated pyelonephritis: A healthy, non-pregnant patient with a healthy urinary system develops a kidney infection [34].

D-Complicated pyelonephritis: An infection of the kidneys that develops during pregnancy or in a patient who has additional aggravating circumstances [as mentioned above under complex cystitis], and that often necessitates hospitalization [35].

E-Recurrent UTI [rUTI]- After the first infection, up to 25-32% of women may experience recurrent episodes, which are classified as 3 episodes within a year or 2 episodes within a 6-month period. The real rate, however, could be considerably higher given that a sizable portion of UTIs go undetected [36]. The most frequent organisms known to cause UTIs are *Escherichia coli* [*E. coli*], *Klebsiella pneumoniae*, *Proteus mirabilis*, *Enterococcus faecalis*, and *Staphylococcus saprophyticus* [37].

F-Asymptomatic bacteriuria [colonization]: > On clean catch urine culture, 100,000 cfu/mL of bacteria are found without any illness-related signs or symptoms [38].

Risk Factors

Females are more prone to get a UTI than men, perhaps as a result of the urethral opening's proximity to the bacteria-rich vagina and rectum, as well as the fact that there is less space

between it and the bladder. In healthy premenopausal females, the probability of periurethral *E. coli* colonisation is increased for both acute cystitis and recurrent UTI in the presence of recent or frequent sexual activity, the use of spermicide, or both [39].

Older age for both women and men, getting a UTI rises with age [especially after 65 and especially after 80]. According to Schaeffer and Nicolle [40], UTI is uncommon in men under the age of 60, but the rate rises significantly after that. By the age of 80, both men and women had comparable rates of UTI.

Decreased estrogen levels also in the postmenopausal women is a risk factor.

Incontinence: the urethral aperture is made vulnerable to germs by fecal incontinence, although most of these bacteria are quickly eliminated from the urinary system unless they are uropathogenic strains [41].

Family history and genetics also influence risk woman's chance of developing UTIs is 2-4 times higher if her mother had them [42]. In addition, it appears that genetic predisposition affects the severity of UTIs [43].

According to Li et al. [44], catheterization significantly raises the risk of a urinary tract infection. Over 70% of UTIs in patients being treated in hospitals are caused by catheters. An indwelling catheter is not as safe as clean intermittent catheterization. UTIs are more common when there are conditions that call for catheterization, such as neurogenic bladder, insufficient bladder emptying, and genitourinary tract structural abnormalities [45].

Procedure. The risk of UTI increases after a urinary tract operation, such as a flexible ureteroscopy for the treatment of bladder cancer or the management of kidney stones. According to a Cochrane analysis, giving patients undergoing cystoscopy antibiotic prophylaxis may lower their likelihood of developing a symptomatic UTI [46]. Urogynecologic surgery patients are more likely to get UTIs. According to Thomas-White et al. [47], between 7% and 24% of women who have surgery for pelvic organ prolapse or stress urine incontinence will get a postoperative UTI.

Recurrent UTIs can be brought on by kidney or bladder stones, which often harbour the same type of bacteria. According to Ripa et al. [48] ureteral stones enhance the probability of urosepsis in pyelonephritis patients.

UTI risk is increased by half in those with diabetes [49]. Although obesity has statistically been linked to a higher incidence of UTI, it is not known if obesity is the root of the problem

[50].

Immunocompromised status, one research found that solid organ transplants, namely kidney transplants, increase the risk of UTI by showing that 28% of 417 patients had a UTI within 13 days of getting the transplant [51]. Human immunodeficiency virus [HIV] infection is not thought to affect the frequency of symptomatic UTI even in those with low CD4 levels [52].

Laboratory detection methods

Midstream urine sampling is one of the most frequent ways to diagnose a UTI, followed by microbiological investigation [urine culture] for pathogen isolation, identification, and antibiotic susceptibility testing [AST] [53]. This type of diagnosis frequently takes two to three days, delaying treatment and promoting the spread of the infectious disease. Due to this restriction, many empirical decisions are made and unnecessary antibiotics are used. Microorganisms can be recovered and concentrated while testing times are decreased using separation and filtration techniques [chemical, physical, and antibody-based procedures]. These procedures are not optimal since they make use of potent chemicals that could harm cells and affect how well bacteria retain their surface characteristics during recovery and concentration [54]. The quantity of an antigen in a solution can be determined using a variety of immunoassays that depend on the precise interaction between an antibody and an antigen. According to Gan and Patel [55], the ELISA test is an enzyme-linked immunosorbent assay, is a biochemical test that identifies antigens, antibodies, and proteins in a sample using antibodies and enzyme-mediated colour change. The use of recently developed methods such as polymerase chain reaction [PCR], matrix-assisted laser desorption/ionization-time of flight [MALDI-TOF], fluorescence in situ hybridization [FISH], and forward light scattering has been beneficial for the diagnosis of UTIs [56]. These methods have significantly shorter detection times and better sensitivity and specificity than culture-based tests. Combining them also enables a precise identification and assessment of the pathogen's medication susceptibility. However, a clinical microbiology laboratory, as well as pricy reagents and equipment, are all necessary to carry out the [57].

Antimicrobial Resistance in UTIs

Due to their ability to obtain genes that code for extended-spectrum lactamases [ESBLs], which are located on transferable plasmids, Gram-negative bacteria, which are the main cause of UTIs, pose an increasing threat to the public's health [58]. These enzymes are unable to hydrolyze carbapenems, but can do so with third-generation cephalosporins and

monobactams [59]. ESBLs are also a public health problem since they are frequently found on plasmids that also carry additional resistance genes against several other classes of antibiotics [such as sulfonamides, aminoglycosides, and quinolones] [60]. Bacteria that take up these plasmids therefore acquire multidrug resistance. Although the mechanism by which all ESBLs function is the amide bond of the β -lactam ring, the genes that code for these enzymes are diverse and broken down into several groups [61]. These methods significantly shorten detection times while improving sensitivity and specificity when compared to culture-based assays. Combining them also makes it easier to accurately identify and assess the pathogen's drug susceptibility [62]. The majority of ESBLs are of the CTX-M type, which are found in *P. aeruginosa* and *Acinetobacter* species as well as other members of the order Enterobacterales [63]. Cefotaxime resistance is increased in isolated isolates harboring CTX-M, and they are less susceptible to ceftazidime [64]. Other significant drug resistance mechanisms include drug target alteration, drug efflux activity, and drug absorption restriction. Antimicrobials target certain bacterial proteins. One frequent method of resistance is to alter these bacterial proteins such that the antibiotic binds ineffectively or not at all. The efflux of medicines from cells via membrane transporters is the most frequent route of bacterial resistance. These transporters are proteins that are members of the ATP-binding cassette [ABC] gene superfamily. Because it enhances the efflux of many medications from cells and lowers the intracellular concentration of pharmaceuticals, overexpression of ABC transporters is a key factor in multidrug resistance [65]. Because uropathogens are becoming more and more resistant to medicines, there are more and more urinary tract infections that people get in hospitals and the general public. As a result of rising fluoroquinolone-resistant Enterobacteriaceae isolation rates, first-choice empiric treatment is no longer advised [66].

Antimicrobial Therapies

Medical practitioners are familiar with the medication nitrofurantoin, and its side effects are controllable. The well-known antimicrobial nitrofurantoin is less effective against *Klebsiella* and *Pseudomonas* than it is against other Gram-negative rods [67]. In especially in patients who are prone to these disorders, such as those with chronic obstructive pulmonary disease, ordering clinicians must be aware of the possibility for serious but relatively uncommon hazards of pulmonary fibrosis and interstitial pneumonitis. Although fosfomycin is suitable for treating a uncomplicated UTI suggest that it may not have as potent an antibacterial effect. Quinolones have a lengthy history of use in the management of UTIs. However, because to rising community resistance rates and the necessity to save it for more serious UTIs, experts recommend not to use it for simple UTIs [68].

Conclusions

Because of the closer proximity of the urethral opening to the bladder, females are more likely than males to get urinary tract infections [UTI]. UTIs have a negative effect on a person's quality of life and are associated with serious morbidity and mortality. The length of the ailment is shortened as a result of antibiotic therapy's efficacy. The development of antibiotic resistance, adverse effects, and other associated problems necessitated the development of alternative management strategies for UTIs. Natural treatments have been used to successfully manage a variety of illnesses to ease symptoms and improve general health.

Abbreviations

Not applicable

Declarations

Ethics approval and consent to participate

Funding

No funds from any institute

Authors' contributions

Ahmed Talib Abdulkarem: the research article proposal, preparing materials, design study, explaining the findings, and article writing. Shaimaa M.S. Zainulabdeen and Saif Mazeel Abed: Data curation, Statistical analysis, and review and editing.

Competing Interests

The authors declare that the research was conducted without any commercial or financial relationships that could be construed as a potential conflict of interest.

References

1. SCHAPPERT, Susan M. RECHTSTEINER, Elizabeth A. Ambulatory medical care utilization estimates for 2007. *Vital and Health Statistics. Series 13, Data from the National Health Survey*.2011; (169): 1-38.
2. Barr, J. G., et al. Microaerophilic/anaerobic bacteria as a cause of urinary tract infection in pregnancy. *BJOG: An International Journal of Obstetrics & Gynaecology*, 1985; 92(5): 506-510.
3. JUAREZ, Guillermo E.; MATEYCA, Celeste; GALVAN, Estela M. Proteus mirabilis outcompetes Klebsiella pneumoniae in artificial urine medium through secretion of ammonia and other volatile compounds. *Heliyon*, 2020; 6.2.
4. Abou Heidar, Nassib F., et al. Management of urinary tract infection in women: A practical approach for everyday practice. *Urology annals*, 2019; (11) 4: 339.
5. Tütüncü, Levent, et al. Urinary tract infection in pregnancy. *Perinatal journal*, 2005; 13[3] 114-121-114-121.
6. Hooton, Thomas M. Uncomplicated urinary tract infection. *New England Journal of Medicine*, 2012; 366(11): 1028-1037.
7. Tullus, Kjell; SHAIKH, Nader. Urinary tract infections in children. *The Lancet*, 2020; (395)10237: 1659-1668.
8. Tien, Brenda Yin Qi, et al. Enterococcus faecalis promotes innate immune suppression and polymicrobial catheter-associated urinary tract infection. *Infection and immunity*, 2017; (85)12: 10.1128/iai. 00378-17.
9. Sofyan, Miyayu Soneta, et al. Management of feline idiopathic cystitis (FIC) using probiotic combination treatment. *The Indian Veterinary Journal*, 2019;(96)12: 20-22.
10. Rizwan, Muhammad, et al. Diagnosis and Management of Catheter Associated Urinary Tract Infection. *Journal of Peoples University of Medical & Health Sciences Nawabshah.(JPUMHS)*, 2017;(7)4: 157-162..
11. Foxman, Betsy. Urinary tract infection syndromes: occurrence, recurrence, bacteriology, risk factors, and disease burden. *Infectious Disease Clinics*, 2014;(28)1: 1-13.
12. Shah, Kairav J., et al. Ampicillin for the treatment of complicated urinary tract infections caused by vancomycin-resistant Enterococcus spp (VRE): a single-center university hospital experience. *International Journal of Antimicrobial Agents*, 2018; (51)1: 57-61.
13. Ehlers, Sarah; Merrill, Stefan A. Staphylococcus saprophyticus. 2018.
14. Salman, Hamzah Abdulrahman, et al. Prevalence of multi-antibiotic resistant bacteria isolated from children with urinary tract infection from Baghdad, Iraq. *Microbiology and Biotechnology Letters*, 2022;(50)1: 147-156.
15. ISSA, Sajjad Salim. Prevalence of Urinary Tract Infection in Children from One to Fifteen Years Old in Basra City in 2014. *Journal of Health, Medicine and Nursing. An International Peer-reviewed Journal*, 2017; 36.
16. Al-Qaidi, Hussein S. Aswed; JASIM, Thamer M.; AL_MAZRUAI, Ibrahim A. Hussein. Urinary Tract Infection Among Children's Under (12) Years Old In Tikrit City. *Al Mustansiriyah Journal of Pharmaceutical Sciences*, 2012;(12)2: 171-177.
17. Zincir, Handan, et al. Prevalence of urinary tract infections and its risk factors in elementary school students. *Urologia internationalis*, 2012;(88)2: 194-197.
18. Isa, M. A., et al. Prevalence of urinary tract infection among primary school children in Maiduguri, Borno State, Nigeria. *International Journal of Environment*, 2013; (2)1: 9-15.
19. Jamal, R.; Rawi, Al. Prevalence of asymptomatic urinary tract infection among primary school children in Iraq [Hilla]. 2003;20-25.
20. Nahab, H. M.; Akeel Hamed Al-Oebady, M.; AQEEL ABDUL MUNEM, H. Bacteriological Study of Urinary Tract Infections among Pregnant Women in Al Samawa City of Iraq. *Archives of Razi Institute*, 2022; 77(1): 117-122.

21. Ateyah Al-Nasrawi, Adel, et al. Antibiotic sensitivity patterns of uropathogens isolated from females with urinary symptoms in Karbala. *Journal of kerbala university*, 2009;(7)2.
22. Tahir Jarjees, Yosra, et al. In vitro resistance to cephalosporins in women with bacterial urinary tract infections. *Iraqi Postgraduate Medical Journal*, 2012;(11)3: 321-325.
23. Kareem, Iman J.; RASHEED, Iman Y. Antibiotic susceptibilities of gram negative aerobic bacteria isolated from urinary tract infections in community. *Iraqi J Med Sci*, 2011;(9)4: 295-300.
24. McLellan, Lisa K.; HUNSTAD, David A. Urinary tract infection: pathogenesis and outlook. *Trends in molecular medicine*, 2016;(22)11: 946-957.
25. Lewis, Adam J.; Richards, Amanda C.; MULVEY, Matthew A. Invasion of host cells and tissues by uropathogenic bacteria. *Urinary Tract Infections: Molecular Pathogenesis and Clinical Management*, 2017; 359-381.
26. Zagaglia, Carlo, et al. Urinary tract infections caused by uropathogenic Escherichia coli strains—new strategies for an old pathogen. *Microorganisms*, 2022; (10)7: 1425.
27. Cabrera, Oscar Alfonso Storme, et al. Risk factors and predisposing conditions for urinary tract infection. 2019.
28. WILEY, Zanthia; JACOB, Jesse T.; BURD, Eileen M. Targeting asymptomatic bacteriuria in antimicrobial stewardship: the role of the microbiology laboratory. *Journal of clinical microbiology*, 2020;(58)5: 10.1128/jcm. 00518-18.
29. Luu, Tyler; ALBARILLO, Fritzie S. Asymptomatic bacteriuria: prevalence, diagnosis, management, and current antimicrobial stewardship implementations. *The American Journal of Medicine*, 2022.
30. Ourani, Mohammad, et al. Evaluation of evidence-based urinalysis reflex to culture criteria: impact on reducing antimicrobial usage. *International Journal of Infectious Diseases*, 2021; 102: 40-44.
31. Nicolle, Lindsay E., et al. Infectious Diseases Society of America guidelines for the diagnosis and treatment of asymptomatic bacteriuria in adults. *Clinical infectious diseases*, 2005;643-654.
32. Bent, Stephen, et al. Does this woman have an acute uncomplicated urinary tract infection?. *Jama*, 2002;(287)20: 2701-2710.
33. Colgan, Richard; WILLIAMS, Mozella. Diagnosis and treatment of acute uncomplicated cystitis. *American family physician*, 2011;(84) 7: 771-776.
34. TONY MAZZULLI, M. D. Diagnosis and management of simple and complicated urinary tract infections (UTIs). *Can J. Urol*, 2012;(19)1: 42-48.
35. Nicolle, Lindsay E. Uncomplicated urinary tract infection in adults including uncomplicated pyelonephritis. *Urologic Clinics of North America*, 2008;(35)1: 1-12.
36. Ciccamese, Federica, et al. Complicated pyelonephritis associated with chronic renal stone disease. *La radiologia medica*, 2021; 126: 505-516.
37. Abdelhamid, Ahmed G.; ESAAM, Aliaa; HAZAA, Mahmoud M. Cell free preparations of probiotics exerted antibacterial and antibiofilm activities against multidrug resistant E. coli. *Saudi pharmaceutical journal*, 2018;(26)5: 603-607.
38. Anderson, Gregory G.; MARTIN, Steven M.; HULTGREN, Scott J. Host subversion by formation of intracellular bacterial communities in the urinary tract. *Microbes and infection*, 2004;(6)12: 1094-1101..
39. Nicolle, Lindsay E. Asymptomatic bacteriuria. *Current opinion in infectious diseases*, 2014;(27)1: 90-96.
40. YOON, Byung Il, et al. Risk factors for recurrent cystitis following acute cystitis in female patients. *Journal of Infection and Chemotherapy*, 2013;(19)4: 727-731.
41. Schaeffer, Anthony J.; NICOLLE, Lindsay E. Urinary tract infections in older men. *New England Journal of Medicine*, 2016 ;(374) 6: 562-571.
42. Raz, Raul, et al. Recurrent urinary tract infections in postmenopausal women. *Clinical Infectious Diseases*, 2000;(30)1: 152-156.

43. Ao L, Q Zeng, DA Fullerton, X Meng, et al. Enhanced monocyte chemoattractant protein-1 production in aging mice exaggerates cardiac depression during endotoxemia. *Critical Care* 2014; 18:1-10.
44. LI, Fei, et al. Risk factors for catheter-associated urinary tract infection among hospitalized patients: A systematic review and meta-analysis of observational studies. *Journal of advanced nursing*, 2019;(75)3: 517-527.
45. Bettcher, Catherine M., et al. Urinary Tract Infection. 2021.
46. Christie, Janice. Antimicrobial agents for preventing urinary tract infections in adults undergoing cystoscopy: A Cochrane review summary. *International Journal of Nursing Studies*, 2020;121: 103646-103646.
47. Thomas-White, Krystal J., et al. Urinary microbes and postoperative urinary tract infection risk in urogynecologic surgical patients. *International urogynecology journal*, 2018; 29: 1797-1805.
48. Alamran FG. System dynamics analysis of mortality post allogeneic hematopoietic stem cell transplant. *American Journal of BioMedicine* 2023; 11(3):110-123.
49. De Lastours, Victoire; FOXMAN, Betsy. Urinary tract infection in diabetes: epidemiologic considerations. *Current infectious disease reports*, 2014, 16: 1-6.
50. Semins, Michelle J., et al. The impact of obesity on urinary tract infection risk. *Urology*, 2012, 79.2: 266-269.
51. Ooms, Liselotte, et al. Urinary Tract Infections After Kidney Transplantation: A Risk Factor Analysis of 417 Patients. *Annals of transplantation*, 2017; 22: 402-408.
52. Chandra, Harish, et al. Promising roles of alternative medicine and plant-based nanotechnology as remedies for urinary tract infections. *Molecules*, 2020;(25) 23: 5593.
53. Davenport, Michael, et al. New and developing diagnostic technologies for urinary tract infections. *Nature Reviews Urology*, 2017;(14)5: 296-310.
54. Kumar, Sandeep, et al. Point-of-care strategies for detection of waterborne pathogens. *Sensors*, 2019;(19)20: 4476.
55. Mo, Manni, et al. Rapid antimicrobial susceptibility testing of patient urine samples using large volume free-solution light scattering microscopy. *Analytical chemistry*, 2019;(91) 15: 10164-10171.
56. Tinôco M, Tupinambás dL. Neutropenic acute lymphocytic leukemia patients with different patterns of bacterial Infections. *American Journal of BioMedicine* 2023; 11(3):136-143. Bedenić, Branka; MEŠTROVIĆ, Tomislav. Mechanisms of resistance in gram-negative urinary pathogens: From country-specific molecular insights to global clinical relevance. *Diagnostics*, 2021; 11.5: 800.
57. Mancuso, Giuseppe, et al. Extended-spectrum β -lactamase & carbapenemase-producing fermentative Gram-negative bacilli in clinical isolates from a University Hospital in Southern Italy. *New Microbiol*, 2021; 44: 227-233.
58. Shaikh, Sibhghatulla, et al. Antibiotic resistance and extended spectrum beta-lactamases: Types, epidemiology and treatment. *Saudi journal of biological sciences*, 2015;(22)1: 90-101.
59. Abou Heidar, Nassib F., et al. Management of urinary tract infection in women: A practical approach for everyday practice. *Urology annals*, 2019; (11)4: 339.
60. Sawa, Teiji; Kooguchi, Kunihiko; Moriyama, Kiyoshi. Molecular diversity of extended-spectrum β -lactamases and carbapenemases, and antimicrobial resistance. *Journal of intensive care*, 2020; 8: 1-13.
61. Castanheira, Mariana; SIMNER, Patricia J.; BRADFORD, Patricia A. Extended-spectrum β -lactamases: an update on their characteristics, epidemiology and detection. *JAC-antimicrobial resistance*, 2021;(3) 3: dlab092.
62. Biondo, Carmelo. Bacterial Antibiotic Resistance: The Most Critical Pathogens. *Pathogens*, 2023;(12)1: 116.
63. Zeynudin, Ahmed, et al. Prevalence and antibiotic susceptibility pattern of CTX-M type

- extended-spectrum β -lactamases among clinical isolates of gram-negative bacilli in Jimma, Ethiopia. *BMC infectious diseases*, 2018;(18)1: 1-10.
64. Schaenzer, Adam J.; WRIGHT, Gerard D. Antibiotic resistance by enzymatic modification of antibiotic targets. *Trends in molecular medicine*, 2020;(26)8: 768-782.
 65. Hoang, Cuong Q., et al. Emergence of New Delhi Metallo-Beta-lactamase (NDM) and Klebsiella pneumoniae Carbapenemase (KPC) production by Escherichia coli and Klebsiella pneumoniae in southern Vietnam and appropriate methods of detection: a cross-sectional study. *BioMed research international*, 2019, 2019.
 66. Syed, Haamid, et al. Nitrofurantoin-induced interstitial pneumonitis: albeit rare, should not be missed. *Case Reports*, 2016, 2016: bcr2015213967.
 67. Cai, Tommaso, et al. Fosfomicin trometamol versus comparator antibiotics for the treatment of acute uncomplicated urinary tract infections in women: a systematic review and meta-analysis. *The Journal of Urology*, 2020; (203)3: 570-578.