

DOI: <http://doi.org/10.32792/utq.jceps.10.01.024>

Investigation and Determination of Further Algae Species in Aquatic Area Of Samawah Province

Athier Abdul Rasool Hassin¹, Dr. Taha Yaseen Farhan², Malath Ismael Yousif³

1. Director of Sawa Lake and Desert Studies Center, Muthanna University

2,3. Director of Sawa Lake and Desert Studies Center, Muthanna University, Muthanna, Iraq

Received 1/11/2018

Accepted 26/11/2018

Published 20/1/2020



This work is licensed under a [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/).

Abstract:

The great environmental importance of algae is formed of so many basis of food chains. This study aims to reveal about certain properties for further of algae in Sawa Lake. The study was the first of its kind on the waters of Sawa Lake, about the algae and its affected by the Physical and chemical properties of Sawa Lake water. Water samples were collected during winter and spring in 2017 from four sites around Sawa Lake. Algae samples were collected from the surface and then the qualitative and quantitative study of the algae found in the water was conducted. The results showed that diatoms or bacilli were predominant, followed by green and blue algae, and recorded a new type of algae which not previously recorded in Iraq (New Record). In conclusion the current study shows that algal growth is increasing in the spring season due to the increase in the activity of analyzers in this season.

Keywords: Sawa lake, Algae, chemical and physical properties, Iraq.

Introduction:

Algae are a various group of aquatic organisms that have the ability to conduct photosynthesis [1]. Algae are affected by the physical and chemical properties of water due to the amount of pollutants present in the water [2]. Algae can be defined as a thalocytes (plants lacking roots, stems and leaves) that are characterized by chlorophyll as a major dye of photosynthesis and lack the cover of sterile cells around their reproductive organs. This definition includes a number of plant forms that are not necessarily in close association with, for example, blue-green cyanobacteria that are closer in their evolution to the bacteria than the rest of the algae [3]. Dissolved oxygen is an important factor for the breathing of plant and animal aquatic organisms that has an important role in the metabolic processes of all living organisms [4]. It is a determining factor for the growth of many aquatic organisms [5]. There are many factors that control the rise of dissolved oxygen in water, such as photosynthesis and temperature. High temperatures reduce the concentration of dissolved oxygen in the water, and bacteria consume dissolved oxygen to dissolve organic matter [6]. In addition, nitrogen is a key element in algal growth. Also, it has an

important role in cell division and certain metabolic functions such as building fatty acids and proteins in algal cells [7]. Nitrogen is found in the nature in several non- organic forms, including NO^{-3} , NO^{-2} and NH^{+4} , as well as in organic forms such as algae and high-end aquatic plants [4]. Also, Phosphorus is an important element in the environment, as it is a determinant of the survival of the biological cycle in the nature because phosphorus is an important life component in the energy transmission devices in the living cells. In the event of a decrease in this element, it reduces the initial productivity [8].

Materials and Methods

Water samples were collected during winter and spring in 2017 from four sites in Sawa Lake (map 1). The physical and chemical properties of the lake water were included. Also, the algae samples were collected from the surface by using the means of a 45 micron algae collection network. These samples were placed in 25 cm³ bottles and then 10 cm³ distilled water with 4-5 drops of Lugols iodine Solution was added as a preservative from melting 15 gm of Iodine in 200 cm³ of distilled water, then adding 20 gm of potassium iodide and 20 cm³ of ice acetic acid were added. After that, the solution was left for several days before using [9]. The qualitative and quantitative study of the algae in the water was measured.

The number of non-diatomic algae cells were calculating one or more drops of the sample were applied after shaking well on the surface of each of the two slices of the counting slide and then placed the slide cover for a few minutes to settle the cells [10]. The final product is expressed by the number of cells in 1cm² of water sample in the field using the conversion coefficient as follows:

Number of cells calculated in 1 cm² = number of cells calculated in one cross-section ×
conversion factor of the concentrated sample


Conversion factor = number of sectors in 1cm³ of concentrated sample × sample concentration factor

Number of cross-sections in 1 cm³ of concentrated sample =1000 mm³

Concentrated sample size in one transverse sector 1mm³.

Original sample size in the cross-section mm³ = Area mm² × 0.1mm (Depth of the lobby)

The blue-green algae were calculated as one unit, while each cell of algae, green, and other algae one unit [11]. The micro transect method was used to calculate the number of diatomic algae cells by the preparation of the permanent slices [12].



Map (1) Algae collection sites in Sawa Lake

Results and Discussion:

1. Physical and Chemical Properties of Sawa Lake water.

1.1 In the current study the air and water temperature values showed clear seasonal changes, which reached the lowest values in winter as shown in Fig. (1). This is due to the nature of Iraqi climate, which is characterized by a large temperature difference according to the seasons of the year [13].




Figure (1). Shows the temperature of Sawa Lake water

1.2 pH: Rising of water levels, increase discharge rate and increase rainfall has led to lower pH values [14].

1.3 Electrical conductivity and salinity: the present study indicates that the salinity and conductivity values recorded in the spring are low, this possibly because of the high water levels in that period, as they reduce salts. The highest salinity and conductivity values during the winter may be caused by a large amounts of soil on the banks of the lake during the rainy season. These soils may be loaded with salts and thus affect the values of the electrical conductivity [15].

1.4 Dissolved Oxygen: the results of measuring dissolved oxygen showed seasonal variation, with the highest winter rates of 13.4 mg /l. This may be due to the fact that cold water has a greater ability to dissolve larger amounts of gases, such as oxygen compared to hot water [4]. The lowest concentration of dissolved oxygen in spring (2) may be due to higher temperatures leading to increased activity of microorganisms in organic matter decomposition, may be to increased oxygen consumption [16].




Figure (2) Shows the dissolved oxygen values of samples during in Saw Lale water

2- Plant Nutrients: seasonal and situational variations of total nitrogen values were recorded due to temperature variations and decomposition Figure (3).




Figure (3). Shows the values of nitrates during the two seasons in Sawa Lake

3- Phosphorus: the results showed that total phosphorus concentrations decreased during the spring season in Sawa Lake and may be due to increased phosphate and nutrient uptake by phytoplankton resulting in less concentrations of the phytoplankton (Figure 4).

3

Winter

2

spring

1

0

Site 1 site 2 site 3 site 4

Examined site




Figure (4). Shows the values of phosphorus during the two seasons in Sawa Lake

3- Phytoplankton: species are belonging to the bacterial algae which have predominance compare with other species of algae in various study sites. This is due to the ability of bactericidal algae to grow and reproduce in a wide range of environmental changes such as temperature, light intensity, plant nutrients and the rapid changes in chemical and biological factors [17]. As well as, their ability to grow in different types of aquatic environments [18]. The class of blue green algae came second only to the diatom category in this study, and the predominance of this variety on the green algae species is consistent with many studies in the water bodies [19]. Followed by blue-green algae (Fig. 5).




Figure (5). Shows the algae ratios in Sawa Lake

In conclusion the current study shows that algal growth is increasing in the spring season due to the increase in the activity of analyzers in this season, which analyze the complex organic compounds into simple compounds (Table 1).

Table (1) shows the types of phytoplankton in Lake Sawa during the period of study (+exists) / (_ does not exist)

TAXA	Station 1	Station 2	Station 3	Station 4
Cyanophyceae				
<i>Chroococciopsis indica</i> (New record)	-	-	+	-
<i>Oscillatoria anguina</i> (Bory) Gomont	-	-	+	+
<i>Spirulina laxissima</i> G. S. West	-	-	+	-
Euglenophyceae				
<i>Phacus</i> sp.	-	+	+	-
Dinophyceae				
<i>Ceratium</i> sp.	-	+	+	+
<i>Peridinium</i> sp.	+	-	+	-
Bacillariophyceae				
Centrales				
<i>Aulacoseira ambigua</i> O.Muller	+	+	+	+
<i>Coscinodiscus lacustris</i> Grunow	+	-	+	+
<i>Cyclotella comta</i> (Ehr.) Kuetzing	+	+	+	-
<i>Stephanodiscus astrea</i> (Ehr.) Gran.	+	-	-	-
Pennales				
<i>Achnanthes affinis</i> Grunow	+	+	-	+
<i>Amphora normanni</i> Rab.	+	-	-	-
<i>Anomoeneis vitrea</i> (Grum.) Ross	-	+	-	+
<i>Bacillaria paxilifer</i> (Muell.) Hendey	+	-	-	+
<i>Caloneis permagana</i> (Bail.) Cleve	-	+	-	+
<i>Cocconeis diminuta</i> Pantocsek	+	+	-	+
<i>Denticula elegans</i> Kutz.	+	-	-	-
<i>Diatoma elongatum</i> (Lyngb) Agardh	+	+	+	+

Fragilaria brevistriata Grounw	+	+	-	-
Gomphonema olivacea (Lyngbye) Dawson Ehrenberg	+	-	-	+
Navicula americana (Ehr.)	+	-	-	+
Pinnularia appendiculata (Ag.) Cleve	-	-	+	-
Rhoicosphenia curvata (Ktz.) Grunow	-	-	+	-
Chlorophyceae				
Chlamydomonas sp	-	+	+	-
Chlorella ellipsoidea Gerneck	+	-	-	-
Scenedesmus abundans (Kirch.) Chodat	+	-	-	-
S. bijuga (Turp.) Lagerheim	-	-	+	-
S.quadricauda (Turp.) de Brebisson	-	+	+	-
Spirogyra crassa Kuetzing	+	-	+	-
Tetraedron minimum (A. Branum) Hansgirg	+	+	+	+

References:

- [1] Lili, B. Z. and Lusan, L. (2010). Biomonitoring and bioindicators used for river Ecosystems: Definitions, Approaches and Trends. Procedia Environmental Sciences 2 (2010) 1510–1524.
- [2] Al-Tamimi, A.A. M (2006). Use of algae Biological evidence to pollute the lower part of theDiyala River organic materials, PhD thesis, University of Baghdad.
- [3] Lee, R.E. (2008). Phycology. Cambridge University Press. Cambridge, UK.
- [4] Wetzel, R.G. (2001) Limnology, lake and river ecology. 3rd ed. Academic Press, 1006 pp.

- [5] Smith, R. (2004). Current methods in aquatic Science. Waterloo, Canada: University of Waterloo.382p.
- [6] Hassani, J.S. (2010). Study of the environment and diversity of algae attached to some aquatic plants in Hawr al-Hawizah in southern Iraq, PhD thesis, University of Baghdad.
- [7] Boney, A. D. (1975). Phytoplankton. Camlot Press. Ltd. Southampton. 116pp.
- [8] Saadi, H. A. (2006). The basics of ecology and the Arabic edition. Amman - Jordan: Dar Al Yazouri Scientific Publishing and Distribution.
- [9] Prescott, G.W. (1979). How to know the freshwater algae? 3rd ed. William C. Brown Co., Publishers, Dubuque, Iowa. 348 pp.
- [10] Martinez, M.R.; Chakroff, R. P. and Pantastico, J.B. (1975). Note on direct phytoplankton counting technique using the hemocytometer. Phil. Agric. 59:1-12.
- [11] APHA, AWWA and WFF (2005). Standard methods for the examination of water and wastewater, 21th ed. American Water Work Association and Environment Federation, USA.
- [12] Hadi, R. A.M. (1981). Algal studies of the river USK. Ph.D. thesis, Univ. College Cardiff. 364 pp.
- [13] Al-Lami, A. A.Z, Salman, S.K.A and Louay Mohamed (2002). The diversity of benthic algae in four gradient rivers salinity in central Iraq, Journal of Fisheries, No. (21): 67-77.
- [14] Kassim, T. I. and Al-Saadi, H. A. (1995). Seasonal variation of epiphytic algae in a marsh area (southern Iraq). Acta Hydrobiol. 37(3): 153-161.
- [15] Jubouri, M. H.and Saleh, S (2009). An environmental and diagnostic study of algae in cross sections of the Tigris River within the governorate of Salah al-Din. Master Thesis, Tikrit University.
- [16] Naser, N.H.A. (2012). Environmental study of algae attached to mud in the Euphrates Eastern Province of Iraq. Master Thesis - University of Qadisiyah / Diwaniyah / (Hafar).
- [17] Kasim, M. and Mukai, H. (2006). Contribution of benthic and epiphytic diatoms to Clam and Oyster production in the Akkeshi-Ko estuary. J. Oceanogr., 62: 267-281.
- [18] Leelahakrie, K.P. and Peerapornpisal, Y. (2010). Diversity of benthic diatoms and water quality of the Ping river Northern Thailand. the international Journal published by the Thai Society of High Education Institutes on Environment. Environment Asia 3(1) ;82-94.
- [19] Saadallah, H.A.A. (1988). Environmental study on the effect of the façade of Saqlawiyah on the Tigris River in the city of Baghdad. Master Thesis . Baghdad University . P. 182.