

## *Zizyphus spina christi* Effect on Pentylenetetrazole-Induced Kindling

Abeer M. Al-Humaidhi<sup>1</sup> MSc, Abdulkareem H. Abd<sup>2</sup> PhD, Haider F. Ghazi<sup>3</sup> PhD

<sup>1</sup>Dept. of Pharmacology, College of Pharmacy, Thi-Qar University, Thi-Qar, Iraq, <sup>2</sup>Dept. of Pharmacology, College of Medicine, Al-Nahrain University, Baghdad, Iraq, <sup>3</sup>Dept. of Microbiology, College of Medicine, Al-Nahrain University, Baghdad, Iraq,

### Abstract

**Background** Epilepsy is a common disorder of brain that is not well controlled via drugs available for it. Gephyrin, a well-known inhibitory synaptic regulator, has a role in epilepsy. Additionally, significant development of oxidative stresses tends to be involved in epileptic seizure pathophysiology that giving rise to neuronal cell death. *Zizyphus spina christi* is a vastly available Iraqi tree and its leaves have several beneficial effects on central nervous system level.

**Objective** To investigate the possible neuroprotective effect of crude *Zizyphus spina christi* extract in kindling model induced.

**Methods** The removed brains of forty albino male mice from four separate groups were exposed to immunohistochemical test for gephyrin, glutathione, and NeuN expressions.

**Results** The data found that pretreatment with *Zizyphus spina christi* crude extract shows highly significant difference (P value <0.001) in immunohistochemical scores for all studied parameters as compared with those being of pentylenetetrazole-kindled group.

**Conclusion** These findings suggest that crude *Zizyphus spina christi* extract of leaves is effective in protection against kindling that induced via pentylenetetrazole when applied orally for male mice.

**Keywords** Gephyrin; glutathione; kindling; NeuN; pentylenetetrazole; *Zizyphus spina christi*

**Citation** Al-Humaidhi AM, Abd AH, Ghazi HF. *Zizyphus spina christi* effect on pentylenetetrazole-induced kindling. Iraqi JMS. 2021; 19(1): 49-55. doi: 10.22578/IJMS.19.1.7

**List of abbreviations:** AEDs = Antiepileptic drugs, CNS = Central nervous system, DPX = Dibutylphthalate polystyrene xylene, GABA = Gamma aminobutyric acid, GSH = Glutathione, HPLC = High performance liquid chromatography, IHC = Immunohistochemistry, NeuN = Neuronal Nuclei, NMDA = N-methyl-D-aspartate, PTZ = Pentylenetetrazole, ZSC = *Zizyphus spina christi*

### Introduction

Epilepsy is a complex etiology disorder of brain that makes individual more susceptible to having unprovoked and repeated seizures. Gephyrin is the base scaffolding protein present in miscellaneous brain regions and a master regulator of neuronal activity at inhibitory sites<sup>(1)</sup>. Notably, one of pathological mechanisms of epileptic

seizure is production of massive reactive oxygen species, which is responsible for death of neuronal cells<sup>(2)</sup>.

The antagonist of gamma aminobutyric acid (GABA) receptor, pentylenetetrazole (PTZ), is utilized experimentally as a chemoconvulsant where its frequent sub-convulsant doses<sup>(3)</sup> resulting in kindling, which is an animal model applied for investigation of the pathophysiology of epilepsy and inspection of anti-epileptic drugs<sup>(4)</sup>.

*Zizyphus spina christi* "Sidr" is a significant plant species and one of well-known Iraqi

trees. Different portions of this plant have been widely utilized in folk medicine.

On the central nervous system (CNS) level, some of laboratory studies have shown that the extract of *Zizyphus spina christi* has antianxiety, neuroprotective as well as central inhibitory effect on rodents<sup>(5-8)</sup>.

Thus, current study aimed to investigate the neuroprotective effect of crude *Zizyphus spina christi* extract against PTZ-induced kindling.

## **Methods**

### **Animals and experimental design**

Forty albino male mice weighing (27.6-34.8) grams were involved in the current study. Mice were housed in standard cages in a room with controlled environment and had free access to food and water. The forty male mice utilized in this study were randomly allocated into the following four groups of ten mice each: Group I served as normal control and got only distilled water. Group II, III, and IV were exposed to induction of kindling via intraperitoneal PTZ injection of 35 mg/kg dose every second day for 23 days. Thirty minutes post each PTZ injection group III and IV were received diazepam (2 mg/kg, orally) and crude extract of *Zizyphus spina christi* leaves (50 mg/kg, orally) respectively.

### **PTZ induced-kindling**

PTZ were purchased from Sigma-Aldrich, USA; its solution was freshly prepared at the day of administration by dissolving 2 mg of it in 1 ml of 0.9% saline<sup>(4,9)</sup>. For kindling induction, a 35 mg/kg of PTZ was applied intraperitoneally every second day. After each PTZ injection, mice were observed for thirty minutes to record the seizure score<sup>(4)</sup>. In the present study, a total of 12 injections (23 days) were needed to acquire kindling.

### **Preparation of brain tissues**

At 24<sup>th</sup> day of study, the intact brain of mice was rapidly removed under deep anesthesia and kept in (10%) formalin containing cup for immunohistochemical evaluation.

### **Immunohistochemical evaluation of gephyrin, glutathione, and NeuN expression in brain tissues**

The 5 µm thicknesses of tissue blocks were affixed on adhesive positively charged slide, dewaxed; rehydrated; put in retrieval solution bath; heated; and treated with peroxidase block for 20 minutes. The slides were rinsed; with wash buffer; treated with protein block; incubated; exposed to diluted antibody; and incubated again for overnight. At the second day, the slides were washed; incubated with horse radish peroxidase polymer; washed with washing buffer. Then, the slides subjected to 3, 3'-diaminobenzidine (DAB) chromogen/substrate reagent; incubated; washed with buffer wash; immersed in Meyers' hematoxylin; washed again; dehydration using ascending grades of 50%, 70%, 90%, and absolute ethanol; immersed in two changes of xylene for ten minutes; mounted with DPX and then covered slipped and left to dry. Immunostaining for expression of each of gephyrin, glutathione, and NeuN were evaluated were based on the percentage of positively stained cells per section 10: Score 0: less than 5%; score 1: 5-25%; score 2: 26-50%; score 3: more than 50%.

### **Statistical analysis**

Data were presented as median, 25% percentile, 75% percentile, mean as well as standard deviation and analyzed using SPSS 20.0. Mann-Whitney U test utilized and the statistical significance was considered as  $P \leq 0.05$ .

## **Results**

### **Effect of crude *Zizyphus spina christi* extract on immunohistochemical expression of gephyrin**

As showed in table 1 and figure 1, IHC scores and brain expression in PTZ-kindled group declined significantly in median and mean  $\pm$  SD (1.00,  $0.57 \pm 0.53$ ) as compared with normal group. In contrast, IHC scores of diazepam (2.00,  $2.43 \pm 0.53$ ) and crude extract of *Zizyphus spina christi* (2.00,  $2.29 \pm 0.48$ )

previewed highly significant increment ( $P \leq 0.001$ ) in relation to those being PTZ-kindled group.


**Table 1: Immunohistochemical scores of gephyrin**

Statistics	Normal control (N=10)	Crude extract (N=10)	Diazepam (N=10)	PTZ (N=10)
Median	2.00	2.00 <sup>NS*</sup>	2.00	1.00
25% Percentile	1	2	2	0
75% Percentile	3	3	3	1
Mean	2.00	2.29	2.43	0.57
SD	0.57	0.48	0.53	0.53

Mann-Whitney test

NS, not significantly different from corresponding diazepam score,  $P > 0.05$

\*, Highly significant different from corresponding PTZ score,  $P \leq 0.001$



**Figure 1. Immunohistochemical expression (40X) of gephyrin for (A) distilled water, (B) crude extract of *Zizyphus spina Christi*, (C) diazepam, (D) pentylentetrazole-kindled tissue**

#### **Effect of crude *Zizyphus spina christi* extract on immunohistochemical expression of glutathione**

IHC scores and brain expression in PTZ-kindled group declined with great statistically significant ( $P \leq 0.001$ ) in median and mean  $\pm$  SD (1.00,  $0.57 \pm 0.53$ ) as compared with apparently healthy group. Contrariwise, IHC

scores of diazepam (3.00,  $2.71 \pm 0.48$ ) and crude *Zizyphus spina christi* extract (3.00,  $2.86 \pm 0.37$ ) previewed highly marked increment ( $P \leq 0.001$ ) in median and mean  $\pm$  SD of glutathione when compared with those of kindled group. All data are presented in table 2 and illustrated in figure 2.

Table 2: Immunohistochemical scores of glutathione

Statistics	Normal control (N=10)	Crude extract (N=10)	Diazepam (N=10)	PTZ (N=10)
Median	2.00	3.00 <sup>NS*</sup>	3.00	1.00
25% Percentile	2	2	2	0
75% Percentile	3	3	3	1
Mean	2.43	2.86	2.71	0.57
SD	0.53	0.37	0.48	0.53

Mann-Whitney test

NS, not significantly different from corresponding diazepam score,  $P > 0.05$

\*, Highly significant different from corresponding PTZ score,  $P \leq 0.05$




Figure 2. Immunohistochemical expression (40X) of glutathione for (A) distilled water, (B) crude extract of *Zizyphus spina Christi*, (C) diazepam, (D) pentylenetetrazole-kindled tissue


**Effect of crude *Zizyphus spina christi* extract on immunohistochemical expression of NeuN**  
IHC scores and staining are shown in table 3 and figure 3 respectively. There was highly significant suppression ( $P \leq 0.001$ ) of NeuN expression in brain tissues of PTZ-kindled group with median and mean  $\pm$  SD of (1.00,  $0.67 \pm$

0.50) in relation to normal group. In the opposite manner, the IHC scores of diazepam group (3.00,  $2.57 \pm 0.53$ ) and crude *Zizyphus spina christi* extract group (2.00,  $1.71 \pm 0.48$ ) were increased remarkably ( $P \leq 0.001$ ) as compared with corresponding in PTZ-kindled group.

**Table 3. Immunohistochemical scores of NeuN**

Statistics	Normal control (N=10)	Crude extract (N=10)	Diazepam (N=10)	PTZ (N=10)
Median	3.00	3.00 <sup>NS*</sup>	3.00	1.00
25% Percentile	2	2	2	0
75% Percentile	3	3	3	1
Mean	2.89	2.71	2.57	0.67
SD	0.33	0.48	0.53	0.5

Mann-Whitney test

NS, not significantly different from corresponding diazepam score,  $P > 0.05$ \*, Highly significant different from corresponding PTZ score,  $P \leq 0.001$ **Figure 3. Immunohistochemical expression (40X) of NeuN for (A) distilled water, (B) crude extract of *Zizyphus spina Christi*, (C) diazepam, (D) pentylentetrazole-kindled tissue**

### Discussion

The kindling models are more look alike human epilepsy syndromes and have been widely utilized to reveal the disease modifying agents<sup>(11)</sup>. In PTZ-induced kindling, repeated administration of sub-convulsive doses of PTZ, a pro-convulsant, resulting in development of seizure activity and ultimately leads to occurrence of primary generalized tonic-clonic seizures<sup>(12,13)</sup>. PTZ originally acts as an antagonist on GABA receptor. Owing to presence of miscellaneous bioactive compounds such as alkaloids, flavonoids, steroids, phenols and saponins in *Zizyphus*

*spina christi*<sup>(14,15)</sup>, it was disclosed that its extract possess diverse pharmacological effects against the brain disorders such as anti-anxiety effect<sup>(5)</sup>, attenuation of memory impairment induced via scopolamine<sup>(16)</sup>, and neuroprotection against cerebral ischemia<sup>(6)</sup>. The present work found that the effects of pretreatment with crude extract exhibited a remarkable gephyrin expression in brain tissues similar to those of diazepam group as compared to those of PTZ-kindled mice. The finding reinforced the sedative effect of *Zizyphus spina christi* leaf extract and it is in harmony with that of previous study<sup>(5)</sup>. This

sedative nature of *Zizyphus crude extract* may be related to its peptide, cyclopeptide alkaloids, and saponin contents. The possible mechanism of alkaloids action may be through modulation of systems of neurotransmitter involving GABA receptors<sup>(17)</sup>. Meanwhile, the underlying mechanism of saponins might be related to its targeting GABAA receptors<sup>(18)</sup>. In the present work, the glutathione and NeuN expression witnessed a remarkable increment in mice pretreated with crude extract of *Zizyphus spina christi* leaf in comparison with that being in PTZ-induced kindled mice. This might be attributed to the presence of the alkaloids, saponins and flavonoids compounds in crude extract. In fact, epilepsy associated with N-methyl-D-aspartate (NMDA) receptors activation that giving rise to oxidative stress production and consequently damage of neurons and even death. The possible mechanism of action of phyto-constituents of plant extract could be due to modulation of neurotransmitter system as well as their anti-oxidant effects<sup>(16,17,19,20)</sup>.

In conclusion, hydro-alcoholic extract of *Zizyphus spina christi* exhibited protective effects against PTZ-induced kindling via upregulation of gephyrin, glutathione as well as NeuN expression. However, further studies are required to investigate its therapeutic effects and to detect its underlying mechanisms.

### **Acknowledgement**

The authors would like to acknowledge all staff members in Department of Pharmacology, College of Medicine, Al-Nahrain University and all staff members in Department of Pharmacognosy, College of Pharmacy, Baghdad University.

### **Author contribution**

Dr. Al-Humaidhi: designed and conducted this research; Dr. Abd and Dr. Ghazi supervised the study and participated in its interpretation.

### **Conflict of interest**

Authors declare no conflict of interest.

### **Funding**

Self-funding.

### **References**

1. Tyagarajan SK, Fritschy JM. Gephyrin: a master regulator of neuronal function? *Nat Rev Neurosci*. 2014; 15(3): 141-56. doi: 10.1038/nrn3670. PMID: 24552784.
2. Mao XY, Zhou HH, Jin WL. Ferroptosis Induction in Pentylentetrazole Kindling and Pilocarpine-Induced Epileptic Seizures in Mice. *Front Neurosci*. 2019; 13: 721. doi: 10.3389/fnins.2019.00721.
3. Cremer CM, Palomero-Gallagher N, Bidmon HJ, et al. Pentylentetrazole-induced seizures affect binding site densities for GABA, glutamate and adenosine receptors in the rat brain. *Neuroscience*. 2009; 163(1): 490-9. doi: 10.1016/j.neuroscience.2009.03.068.
4. Shimada T, Yamagata K. Pentylentetrazole-induced kindling mouse model. *J Vis Exp*. 2018; (136): 56573. doi: 10.3791/56573.
5. Setorki, M. Effect of hydro-alcoholic extract of *Zizyphus spina-christi* against scopolamine-induced anxiety in rats. *Bangladesh J Pharmacol*. 2016; 11: 421-7. doi: 10.3329/bjp.v11i2.26505.
6. Setorki M, Hooshmandi Z. Neuroprotective effect of *Zizyphus spina-christi* on brain injury induced by transient global cerebral ischemia and reperfusion in rat. *Bangladesh J Pharmacol*. 2017; 12: 69-76. doi: 10.3329/bjp.v12i1.29964.
7. Adzu B, Amos S, Dzarma S, et al. Effect of *Zizyphus spina-christi* Willd aqueous extract on the central nervous system in mice. *J Ethnopharmacol*. 2002; 79(1): 13-6. doi: 10.1016/s0378-8741(01)00348-8.
8. Adzu B, Haruna A, Ilyas M, et al. CNS activity of ZS-1A: a phytochemical from *Zizyphus spina-christi* root bark. *Int J Biol Chem Sci*. 2009; 2: 456-61. doi: 10.4314/ijbcs.v2i4.39766.
9. Erkec OE, Arihan O. Pentylentetrazol kindling epilepsy model. *J Turkish Epilepsi Soc*. 2015; 21: 6-12. doi: 10.5505/epilepsi.2015.08108.
10. Hou X, Wang X, Zhang L. Conditional downregulation of brain- derived neurotrophic factor and tyrosine kinase receptor B blocks epileptogenesis in the human temporal lobe epilepsy hippocampus. *Neurol India*. 2010; 58(1): 29-34. doi: 10.4103/0028-3886.60392.
11. Hui Yin Y, Ahmad N, Makmor-Bakry M. Pathogenesis of epilepsy: challenges in animal models. *Iran J Basic Med Sci*. 2013; 16(11): 1119-32.
12. Ammon-Treiber S, Grecksch G, Angelidis C, et al. Pentylentetrazol-kindling in mice overexpressing heat shock protein 70. *Naunyn Schmiedebergs Arch Pharmacol*. 2007; 375(2): 115-21. doi: 10.1007/s00210-007-0143-0.
13. Bialer M, White HS. Key factors in the discovery and development of new antiepileptic drugs. *Nat Rev Drug Discov*. 2010; 9(1): 68-82. doi: 10.1038/nrd2997.
14. Alhassan KA, Indabawa AS, Shah M. Phytochemical analysis, proximate composition and antibacterial activities of *Zizyphus* Species (*Z. jujube* and *Z. spina*

- christi). *J Appl Adv Res.* 2019; 4: 42. doi: 10.21839/jaar.2019.v4i1.262.
15. Ahmed J, Salim KA, Lim LB, et al. Evaluation of antioxidant activity and phytochemical screening of leaves, barks, stems and fruits of *Alphitonia philippinensis* (Rhamnaceae) from Brunei Darussalam. *Polymer J.* 2019; 11, 951-61. doi: 10.5530/pj.2019.11.151.
  16. Notarki M, Setorki M, Hooshmandi Z. The antioxidant effect of Sidr (*Zizyphusspina-christi*) leaf extract helping to improve the scopolamine induced memory impairment in male rats. *Iran J Pharmaceut Sci.* 2017; 13, 13-24. doi: 10.22034/IJPS.2017.31129.
  17. Hussain G, Rasul A, Anwar H, et al. Role of plant derived alkaloids and their mechanism in neurodegenerative disorders. *Int J Biol Sci.* 2018; 14(3): 341-57. doi: 10.7150/ijbs.23247.
  18. Shen CY, Wan L, Zhu JJ, et al. Targets and underlying mechanisms related to the sedative and hypnotic activities of saponin extracts from semen *Ziziphus jujube*. *Food Funct.* 2020; 11(5): 3895-903. doi: 10.1039/d0fo00098a.
  19. Citraro R, Navarra M, Leo A, et al. The Anticonvulsant Activity of a Flavonoid-Rich Extract from Orange Juice Involves both NMDA and GABA-Benzodiazepine Receptor Complexes. *Molecules.* 2016; 21(9): 1261. doi: 10.3390/molecules21091261.
  20. Zhou YJ, Chen JM, Sapkota K, et al. Panax notoginseng saponins attenuate CCL2-induced cognitive deficits in rats via anti-inflammation and anti-apoptosis effects that involve suppressing over-activation of NMDA receptors. *Biomed Pharmacother.* 2020; 127: 110139. doi: 10.1016/j.biopha.2020.110139.

---

**Correspondence to Abeer M. Al-Humaidhi**

**E-mail: [abeer-m@utq.edu.iq](mailto:abeer-m@utq.edu.iq)**

**Received Nov. 2<sup>nd</sup> 2020**

**Accepted Dec. 23<sup>rd</sup> 2020**