



Tikrit Journal of Pure Science

ISSN: 1813 – 1662 (Print) --- E-ISSN: 2415 – 1726 (Online)

Journal Homepage: <http://tjps.tu.edu.iq/index.php/j>



Prevalence of *Helicobacter pylori* infected in the Kirkuk native population and associated with serum ferritin and Iron levels/Iraq

Nadia M. Mohammad¹, Sabah M. Salih²

¹ Department of Biology , College of Education for Pure Sciences , Tikrit University , Tikrit , Iraq

² College of Nursing, University of Kirkuk, Kirkuk, Iraq

DOI: <http://dx.doi.org/10.25130/tjps.24.2019.029>

ARTICLE INFO.

Article history:

-Received: 2 / 9 / 2018

-Accepted: 9 / 10 / 2018

-Available online: / / 2019

Keywords: Iron deficiency, *H.pylori* infection, ferritin.

Corresponding Author:

Name: Nadia M. Mohammad

E-mail:

Naduasultan33@yahoo.com

Tel:

Introduction

Helicobacter pylori are urease producing organisms gram-negative that are found throughout the world and produce chronic gastritis, it may also predispose patients to develop duodenal ulcers, gastric lymphoma or carcinoma [1,2]. Current evidences suggest that *Helicobacter pylori* is associated with iron deficiency. Many studies have found an association bacteria and low body iron stores and iron deficiency anemia and a reduced response to iron supplementation[3,4]. Iron deficiency, defined as decreased total body iron content, is among the most common nutritional deficiencies in the world. Iron deficiency results in impairments in immune, cognitive, and reproductive functions as well as decreased work performance[5]. Iron deficiency develops through three phases: 1-iron depletion, 2-iron-deficient erythropoiesis, and 3-iron-deficiency anemia (IDA)[1], clinical and epidemiologic studies suggest that infection with bacterial is associated with iron deficiency and IDA[6]. Barabino hypothesized that gastritis levels of neutrophil-derived lactoferrin, the infection would result in the rise increased iron losses related to bacterial turnover[7]. A study in Alaska showed that anemia responded to oral iron replacement but recurred when iron therapy was discontinued suggesting that mild chronic bleeding was involved[8]. The pathogenesis is multifactorial, including combinations of reduced iron absorption

ABSTRACT

An association of *Helicobacter pylori* and iron deficiency (ID) was conducted in Kirkuk city on 176 patients attending private laboratories in Kirkuk for the period from 1/8/2016 to 1/4/2017, blood sampling and a data collection survey were performed in 176 patients suffering from digestive problems and disorder, 120 of them with *H.pylori*, serum ferritin and iron were measured by ELISA and direct enzymatic method techniques respectively. the result showed that 67 (55.83%) of the patients have serum ferritin concentrations below the normal range indicating iron deficiency, 69 (57.5%) of the patients have iron concentrations below the normal range indicating iron deficiency.

related to decreased acid secretion, increased iron loss from microbleeding, and utilization by bacteria. The purpose of this study is to investigate the infection with *Helicobacter pylori* gastritis association with iron-deficiency (ID) in some Kirkuk patients.

Materials and methods

The patients were referred to private laboratories in Kirkuk city in period from 1/8/2016 to 30/4/2017 *H.pylori* infection was diagnosed serologically by using anti-*H. Pylori* IgG, antibodies Elisa kit (CALBIOTECH IgG.USA). Totally 176 patients 120 were seropositive *H. pylori*-IgG and 56 anti-*H. pylori*-IgG seronegative. Some factors such as age, sex, smoking, drug treatment and Chronic diseases (gastric lymphoma, duodenal ulcers or carcinoma) were recorded for all of the patients

Sample Collections

from the subjects 5 ml venous blood was obtained. All samples were dispensed into dry glass test tubes for clotting and retraction to take place. Sera was obtained after samples were centrifuged at 2000 g for ten minutes and stored at -20°C until assayed for laboratory investigations.

Ferritin was detected by ELISA., this assay system utilizes one rabbit anti-ferritin antibody for solid phase immobilization and a mouse monoclonal anti-ferritin antibody in the antibody-enzyme (horseradish peroxidase) conjugate solution. The normal range of

ferritin as recommended by BioCheck is 20-250ng/ml for male and 10- 120ng/ml for female, serum iron concentration was investigated by direct enzymatic method., after dissociation of iron- transferring bound in acid medium, ascorbic acid reduces Fe+3 iron into Fe+2 iron. The absorbance measured at 600 nm is directly proportional to the amount of iron in the specimen. The normal rang of iron as recommended by BIOLABO is 11.6-31.1 $\mu\text{mol/L}$ for male and 9.0-30.4 $\mu\text{mol/L}$ for female.

Statistical Analysis

All values were expressed as mean $\pm$ SD. Statistical analyses were done using the Student's t-test to assess differences among patients. The level of significance was set at $P < 0.05$.

Results

A total of 120 patients (79 women and 41 men) with infected *H.pylori* were enrolled in this study. Mean $\pm$ SD age of the 79 female/male patients was $44.2 \pm 14.4 / 45.4 \pm 14.9$ years, range 15-74/18- 73 years,

respectively. Table 1 shows the lifestyle and clinical characteristics of subjects (women and men).

Table 1. A number of patients infected by *Helicobacter pylori* bacteria according to age group (year), smoking and presence of chronic disease

Varaibl	Men (n=41)	Women (n=79)
Age	(18-75)	(15-74)
Mean $\pm$ SD	43.4 ± 14.2	44.2 ± 14.4
Range		
Smoking%	32 (78.04)	11 (13.92)
Chronic diseases %	15 (36.58)	37 (46.83)

Of the 120 *H.pylori* patients (67) 55.83%, showed low concentration of ferritin, (69) 57.5% showed low concentration of iron. In general median serum ferritin and iron levels were significantly lower than the normal ranges in *H.pylori* infected patients than in anti- *H.pylori*-IgG seronegative as it showed in table 2,3.

Table 2 .shows serum concentration of ferritin (ng/ml) according to prevalence of *H .pylori* under study

Serum ferritin conc (ng/ml)	H.Pylori (+ve) %	H.pylori (-ve) %	Total %
Low ferritin conc.	67 (55.83)	11 (19.64)	78 (44.31)
Normal ferritin conc.	53 (44.16)	45 (80.35)	(55.68) 98
Total	120 (100)	56 (100)	176 (100)
Chi-Square=**20.265		P value=0.0006	

There was no significant difference in median concentrations of both iron and ferritin between women and men.

Table 3. presents serum concentration of iron ($\mu\text{mol/L}$) according to prevalence of *H .pylori* under study

Serum Iron conc.	H.Pylori (+ve) %	H.pylori (-ve) %	Total %
Low Iron conc.	69 (57.5)	12 (21.42)	81 (46.02)
Normal Iron conc.	51 (42.5)	44 (78.57)	95 (53.97)
Total	120 (100)	56 (100)	176 (100)
Chi-Square =**19.999		P value=0.0005	

Discussion

iron deficiency and IDA are conditions with important health consequences regarding reproduction, work performance, immunity and possibly cognitive development; it is a simple result of an imbalance between absorption and iron loss. It is unclear why some patients with *H. pylori* infection develop IDA. Postulates include decreased iron absorption secondary to atrophic gastritis, reduced gastric acid production, decreased ascorbic acid concentration in stomach, and uptake of iron by bacteria. [6,9] as *Helicobacter pylori* sociated gastritis has emerged as a cause of IDA that is unresponsive to iron therapy.[3,10] Laboratory Investigations: Ferritin is a very good marker for iron deficiency but iron is measured too as another indicator for iron deficiency due to ferritin which is an acute phase protein, it can be elevated in inflammation conditions that a normal serum ferritin may not always exclude iron deficiency in this results the proposal that *Helicobacter.pylori* infection is associated with ID, this proposal is achieved by many studies with different explanation of the mechanisms *abaout Helicobacter. pylori* affect

iron absorbance. Baysoy, *et al.* have investigated *Helicobacter. pylori* related-changes in gastric physiology and histology. They have reported that *H.pylori* infection is associated with low serum iron levels and with a decrease in gastric juice ascorbic acid concentration.[10] Capurso, *et al.* has suggested that *H.pylori* infection may be the cause of atrophic gastritis leading to achlorhydria and gastric hypoacidity.[11] A study in Alaska including 2080 adult patients, where there is a high prevalence of *H.pylori*, have suggested a significant correlation between *H.pylori*-IgG positivity and low serum ferritin levels. [12] They have suggested that ulceration causes bleeding which leads to IDA and this result was confirmed in this study. The blood loss in chronic gastritis, and bleeding from duodenal or gastric ulcers related to *H pylori* infection, which plays an important role in the development of iron deficiency in adults. In response to. *Helicobacter. pylori* chronic gastric inflammation, the epithelial cells in the mucosa are damaged, leading to detachment and apoptosis. In the absence of bleeding lesions, the possible mechanisms by which.

Helicobacter pylori is involved in the development of IDA remain not understading. studies suggest that the growth and proliferation of. *Helicobacter pylori* requires iron from the host and that some *Helicobacter. pylori* strains have an ability to interfere with iron metabolism by binding iron to their outer membrane proteins[13].

Moreover, Boggs reviewed that eradication of *Helicobacter. pylori* with a triple therapy consisting of clarithromycin, amoxicillin and lansoprazole for

References

- [1] Bui, D.; Brown HE.; Harris RB. and Oren E (2016). Serologic Evidence for fecal oral Transmission of *Helicobacter pylori*. Am J Trop Med Hyg 94:82-88.
- [2] Blaser, M.J and Atherton, J.C. (2004). *Helicobacter pylori* persistence: biology and disease. J Clin Invest ; 113:321-333.
- [3] Tan, H.J;Goh KL. (2012): Extragastrintestinal Mani festations of *Helicobacter pylori* infection fact or myth? Acritical review. J. Dig Dis: 13(7):342-9.
- [4] Mahalanabis, D.; Islam MA. and Shaikh, S. (2005). Haematological response to iron supplementation is reduced in children with asymptomatic *Helicobacter pylori* infection. Br J Nutr; 94: 969-975.
- [5] Akodu OS.; Disu EA.; Njokanma O. and Kehinde OA. (2016). Iron deficiency anaemia among apparently healthy pre-school children in lagos Nigeria AFR Health sci 16:61-68.
- [6] Dubois, S. and Kearney, DJ. (2005). Iron-deficiency anemia and *Helicobacter pylori* infection: a review of the evidence. Am J Gastroenterol; 100:453-9.
- [7] Short MW. and Domagalski JE. (2013). Iron deficiency anemia: evaluation and management. Am fam physician 87:98-104.
- [8] Yip, R.; Limburg, PJ. and Ahlquist, DA. (1997). Pervasive occult gastrointestinal bleeding in an Alaska native population with prevalent iron deficiency. Role of *Helicobacter pylori* gastritis. JAMA; 277:1135-9.
- [9] Stoltzfus, RJ. (2003). Iron deficiency: global prevalence and consequences. Food Nutr Bull; 24: S97-S101.
- [10] Baysoy, G.; Ertem, D. and Ademoglu, E.(2004). Gastric histopathology, iron status and iron deficiency anemia in children with *Helicobacter pylori* infection. J Pediatr Gastroenterol Nutr; 38: 146-151.
- [11] Capurso, G.; Lahner, E. And Marcheggiano ,A. (2001). Involvement of the corporal mucosa and related changes in gastric acid secretion characterize patients with iron deficiency anaemia associated with *Helicobacter pylori* infection. Aliment Pharmacol Ther; 15: 1753-1761.
- [12] Yip, R.; Limburg, PJ and Carpenter, H.A (1997). Pervasive occult gastrointestinal bleeding in an Alaska native population with prevalent iron deficiency. Role of *Helicobacter pylori* gastritis. JAMA; 277:1135-1139.
- [13] Perez-Perez, GI. and Israel, DA. (2000). Role of iron in *Helicobacter pylori*: its influence in outer membrane protein expression and in pathogenicity. Eur J Gastroenterol Hepatol; 12: 1263- 1265.
- [14] Boggs, Well. (2009) Eradication of *H.pylori* reduces iron deficiency anemia prevalence. J Infect Dis; 199:652-660.

الإصابة باللولبيات البوابية المصاحبة لنقص الحديد لدى المرضى في مدينة كركوك

نادية محمود محمد¹، صباح محمد صالح²

¹قسم علوم الحياة ، كلية التربية للعلوم الصرفة ، جامعة تكريت ، تكريت ، العراق

²كلية التمريض ، جامعة كركوك ، كركوك ، العراق

الملخص

ثمة علاقة بين الإصابة باللولبيات البوابية ونقص الحديد، في هذه الدراسة تم الحصول على نماذج دم من 176 مريضاً من المراجعين للمختبرات الخاصة في مدينة كركوك يعانون من سوء الهضم واضطرابات معدية ومعدية للفترة من 1/ 8/ 2016 ولغاية 1/ 4/ 2017 تبين إصابة 120 مريضاً ببكتريا اللولبيات البوابية، وعند قياس تراكيز الحديد والفريت في مصل الدم لجميع المرضى تبين ان 67 (55.83%) مريضاً كان مستوى الفريت منخفض عن المستوى الطبيعي و69 (57.5%) مريضاً نسبة الحديد منخفضة عن المستوى الطبيعي مما يؤكد إصابتهم بمرض فقر الدم الناتج عن نقص الحديد، وجدت فروقات معنوية في الإصابة بين الذكور والإناث إذ بلغت 41 (34.3%) و79 (65.83%) على التوالي.