

## **Generalization of Fuzzy Laplace Transforms for Fuzzy Derivatives**

**تعميم تحويلات لابلاس الضبابية للمشتقات الضبابية**

Amal K. Haydar\* and Hawrra F. Mohammad Ali

\*Department of Mathematics, Faculty of Education for Girls, Kufa University, Iraq.

[\\*amalkh.hayder@uokufa.edu.iq](mailto:amalkh.hayder@uokufa.edu.iq)

### **Abstract**

The main aim of this paper is to find the general formula to the fuzzy derivative of the  $n$ -th order and the general formula for fuzzy Laplace transforms to the fuzzy derivative of the  $n$ -th order by using generalized H-differentiability.

### **المستخلص**

الهدف الرئيسي من هذا البحث ايجاد الصيغة العامة للمشتقة الضبابية من الرتبة  $n$  وايضا الصيغة العامة الى تحويلات لابلاس الضبابية العامة للمشتقة الضبابية من الرتبة  $n$  باستخدام قابلية الاشتقاق  $H$  - المعممة .

### **1. Introduction**

The concept of the fuzzy derivative was first introduced by Chang and Zadeh in [1], it was followed up by Dubois and Prade in 1982 [2] and Puri and Ralescu in 1983 [3]. Abbasbandy et al. [4] studied a numerical method for  $n$ -th order fuzzy differential equations based on Seikkala derivative with initial value conditions, Allahviranloo et al. [5] considered eigenvalue-eigenvector method for solving  $n$ -th order fuzzy differential equations with fuzzy initial conditions, Bede et al. [6] provided solution of first order linear fuzzy differential equations by variation of constant formula. . Recently, many researchers worked on solving fuzzy differential equations by using fuzzy Laplace transforms, especially in [7,8,9].

This paper is arranged as follows: Basic concepts are given in Section 2. In Section 3, the general formula for the fuzzy derivative of the  $n$ -th order and fuzzy Laplace transform for the fuzzy derivative of the  $n$ -th order are found. In Section 4, an example of the third order is solved. In Section 5, conclusions are drawn.

### **2. Basic Concepts**

In this section, some necessary definitions and concepts are introduced:

**Definition 2.1 [8]** A fuzzy number  $u$  in parametric form is a pair  $(\underline{u}, \bar{u})$  of functions  $\underline{u}(\alpha)$  and  $\bar{u}(\alpha)$ ,  $0 \leq \alpha \leq 1$  which satisfy the following requirements:

1.  $\underline{u}(\alpha)$  is a bounded non-decreasing left continuous function in  $(0,1]$ , and right continuous at 0,
2.  $\bar{u}(\alpha)$  is a bounded non-increasing left continuous function in  $(0,1]$ , and right continuous at 0,
3.  $\underline{u}(\alpha) \leq \bar{u}(\alpha)$ ,  $0 \leq \alpha \leq 1$ .

**Definition 2.2 [8]** Let  $x, y \in E$ . If there exists  $z \in E$  such that  $x = y + z$ , then  $z$  is called the H-difference of  $x$  and  $y$ , and it is denoted by  $x \ominus y$ . In this paper, the sign “ $\ominus$ ” always stands for H-difference, and also note that  $x \ominus y \neq x + (-1)y$ .

**Definition 2.3 [8]** Let  $f : (a,b) \rightarrow E$  and  $x_0 \in (a,b)$ . We say that  $f$  is strongly generalized differential at  $x_0$  if there exists an element  $f'(x_0) \in E$ , such that

i. For all  $h > 0$  sufficiently small,  $\exists f(x_0 + h) \ominus f(x_0)$ ,  $\exists f(x_0) \ominus f(x_0 - h)$  and the limits (in the metric  $d$ )

$$\lim_{h \rightarrow 0} [(f(x_0 + h) \ominus f(x_0)) / h] = \lim_{h \rightarrow 0} [(f(x_0) \ominus f(x_0 - h)) / h] = f'(x_0) \text{ or}$$

ii. For all  $h > 0$  sufficiently small,  $\exists f(x_0) \ominus f(x_0 + h)$ ,  $\exists f(x_0 - h) \ominus f(x_0)$  and the limits (in the metric  $d$ )

$$\lim_{h \rightarrow 0} [(f(x_0) \ominus f(x_0 + h)) / (-h)] = \lim_{h \rightarrow 0} [(f(x_0 - h) \ominus f(x_0)) / (-h)] = f'(x_0) \text{ or}$$

iii. For all  $h > 0$  sufficiently small,  $\exists f(x_0 + h) \ominus f(x_0)$ ,  $\exists f(x_0 - h) \ominus f(x_0)$  and the limits (in the metric  $d$ )

$$\lim_{h \rightarrow 0} [(f(x_0 + h) \ominus f(x_0)) / h] = \lim_{h \rightarrow 0} [(f(x_0 - h) \ominus f(x_0)) / (-h)] = f'(x_0) \text{ or}$$

iv. For all  $h > 0$  sufficiently small,  $\exists f(x_0) \ominus f(x_0 + h)$ ,  $\exists f(x_0) \ominus f(x_0 - h)$  and the limits (in the metric  $d$ )

$$\lim_{h \rightarrow 0} [(f(x_0) \ominus f(x_0 + h)) / (-h)] = \lim_{h \rightarrow 0} [(f(x_0) \ominus f(x_0 - h)) / h] = f'(x_0)$$

**Definition 2.4 [7]** Let  $f(t)$  be continuous fuzzy-valued function. Suppose that  $f(t) e^{-st}$  is

improper fuzzy Riemann-integrable on  $[0, \infty)$ , then  $\int_0^\infty f(t) e^{-st} dt$  is called fuzzy Laplace

transforms and is denoted as  $L(f(t)) = \int_0^\infty f(t) \cdot e^{-st} dt$ , ( $s > 0$ ). We have

$$\int_0^\infty f(t) e^{-st} dt = \left( \int_0^\infty \underline{f}(t, \alpha) e^{-st} dt, \int_0^\infty \bar{f}(t, \alpha) e^{-st} dt \right),$$

also by using the definition of classical Laplace transform

$$l[\underline{f}(t, \alpha)] = \int_0^\infty \underline{f}(t, \alpha) \cdot e^{-st} dt \text{ and } l[\bar{f}(t, \alpha)] = \int_0^\infty \bar{f}(t, \alpha) \cdot e^{-st} dt,$$

then, we follow:

$$L(f(t)) = (l(\underline{f}(t, \alpha)), l(\bar{f}(t, \alpha))).$$

### 3. Fuzzy Laplace Transforms for the $n$ -th Derivative

In this section, we find a general form for the fuzzy derivative of any order  $n$ ,  $n \in \mathbb{Z}^+$ . Also, we find, a generalization for fuzzy Laplace transforms for  $n$ -th derivative.

**Theorem 3.1** Suppose that  $F(t), F'(t), \dots, F^{(n-1)}(t)$  are differentiable fuzzy valued functions such that  $F^{(i_1)}(t), F^{(i_2)}(t), \dots, F^{(i_m)}(t)$  are (ii)-differentiable functions for  $0 \leq i_1 < i_2 < \dots < i_m \leq n-1$ ,  $0 \leq m \leq n$  and  $F^{(p)}(t)$  is (i)-differentiable for  $p \neq i_j, j = 1, 2, \dots, m$ , and if  $\alpha$ -cut representation of fuzzy-valued function  $F(t)$  is denoted by  $[F(t)]^\alpha = [f_\alpha(t), g_\alpha(t)]$ , then:

(a) If  $m$  is an even number then  $[F^{(n)}(t)]^\alpha = [f_\alpha^{(n)}(t), g_\alpha^{(n)}(t)]$ .

(b) If  $m$  is an odd number then  $[F^{(n)}(t)]^\alpha = [g_\alpha^{(n)}(t), f_\alpha^{(n)}(t)]$ .

**Proof** We shall prove by mathematical induction on  $n$ . We have the relations (a) and (b) are true for  $n = 1$  [7].

We suppose that the relations (a) and (b) are true for  $n = k$ , i.e., if  $F, F', \dots, F^{(k-1)}$  are differentiable fuzzy-valued functions such that  $F^{(i_1)}, F^{(i_2)}, \dots, F^{(i_m)}$  are (ii)-differentiable functions for  $0 \leq i_1 < i_2 < \dots < i_m \leq k - 1$ ,  $0 \leq m \leq k$ , then

If  $m$  is an even number we have:

$$[F^{(k)}(t)]^\alpha = [f_\alpha^{(k)}(t), g_\alpha^{(k)}(t)], \tag{3.1}$$

and if  $m$  is an odd number we have:

$$[F^{(k)}(t)]^\alpha = [g_\alpha^{(k)}(t), f_\alpha^{(k)}(t)]. \tag{3.2}$$

Now, we prove that the relations (a) and (b) are true for  $n = k + 1$  as follows:

(a) Let  $F, F', \dots, F^{(k)}$  be differentiable fuzzy valued functions such that  $F^{(i_1)}, F^{(i_2)}, \dots, F^{(i_m)}$  are (ii)-differentiable functions for  $0 \leq i_1 < i_2 < \dots < i_m \leq k$ ,  $0 \leq m \leq k + 1$  and  $m$  is an even number. If  $m = 0$  then there is no (ii)-differentiable functions, i.e.,  $F(t), F'(t), \dots, F^{(k)}(t)$  are (i)-differentiable. Since  $F(t), F'(t), \dots, F^{(k-1)}(t)$  are (i)-differentiable then by relation (3.1) of induction hypothesis we have:

$$[F^{(k)}(t)]^\alpha = [f_\alpha^{(k)}(t), g_\alpha^{(k)}(t)].$$

Since  $F^{(k)}$  is (i)-differentiable then from definition 2.3, we have:

$$[F^{(k)}(t+h) \Theta F^{(k)}(t)]^\alpha = [f_\alpha^{(k)}(t+h) - f_\alpha^{(k)}(t), g_\alpha^{(k)}(t+h) - g_\alpha^{(k)}(t)],$$

$$[F^{(k)}(t) \Theta F^{(k)}(t-h)]^\alpha = [f_\alpha^{(k)}(t) - f_\alpha^{(k)}(t-h), g_\alpha^{(k)}(t) - g_\alpha^{(k)}(t-h)].$$

and, multiplying by  $\frac{1}{h}, h > 0$  we get:

$$\frac{1}{h} [F^{(k)}(t+h) \Theta F^{(k)}(t)]^\alpha = \left[ \frac{f_\alpha^{(k)}(t+h) - f_\alpha^{(k)}(t)}{h}, \frac{g_\alpha^{(k)}(t+h) - g_\alpha^{(k)}(t)}{h} \right],$$

and

$$\frac{1}{h} [F^{(k)}(t) \Theta F^{(k)}(t-h)]^\alpha = \left[ \frac{f_\alpha^{(k)}(t) - f_\alpha^{(k)}(t-h)}{h}, \frac{g_\alpha^{(k)}(t) - g_\alpha^{(k)}(t-h)}{h} \right].$$

Using  $h \longrightarrow 0$  on both sides of aforementioned relation, we get

$$[F^{(k+1)}(t)]^\alpha = [f_\alpha^{(k+1)}(t), g_\alpha^{(k+1)}(t)]. \tag{3.3}$$

Then the relation (a) is true if  $m = 0$ .

Now, if  $m$  is an even number such that  $2 \leq m \leq k + 1$ , then we have two possibilities for the functions  $F, F', \dots, F^{(k-1)}$  as follows:

**Either**  $F, F', \dots, F^{(k)}$  contain an even number  $m$  and  $F, F', \dots, F^{(k-1)}$  contain an even number  $m$  of (ii)-differentiable functions  $F^{(i_1)}, F^{(i_2)}, \dots, F^{(i_m)}$  then  $F^{(k)}$  is (i)-differentiable. By relation (3.1) of induction hypothesis, we have  $[F^{(k)}(t)]^\alpha = [f_\alpha^{(k)}(t), g_\alpha^{(k)}(t)]$ .

Since  $F^{(k)}$  is (i)-differentiable then from definition 2.3, we have:

$$[F^{(k)}(t+h) \Theta F^{(k)}(t)]^\alpha = [f_\alpha^{(k)}(t+h) - f_\alpha^{(k)}(t), g_\alpha^{(k)}(t+h) - g_\alpha^{(k)}(t)],$$

$$[F^{(k)}(t) \Theta F^{(k)}(t-h)]^\alpha = [f_\alpha^{(k)}(t) - f_\alpha^{(k)}(t-h), g_\alpha^{(k)}(t) - g_\alpha^{(k)}(t-h)].$$

and, multiplying by  $\frac{1}{h}, h > 0$  we get:

$$\frac{1}{h} [F^{(k)}(t+h) \Theta F^{(k)}(t)]^\alpha = \left[ \frac{f_\alpha^{(k)}(t+h) - f_\alpha^{(k)}(t)}{h}, \frac{g_\alpha^{(k)}(t+h) - g_\alpha^{(k)}(t)}{h} \right],$$

and

$$\frac{1}{h}[F^{(k)}(t) \Theta F^{(k)}(t-h)]^\alpha = \left[ \frac{f_\alpha^{(k)}(t) - f_\alpha^{(k)}(t-h)}{h}, \frac{g_\alpha^{(k)}(t) - g_\alpha^{(k)}(t-h)}{h} \right].$$

Using  $h \rightarrow 0$  on both sides of aforementioned relation, we get the relation (3.3).

**Or**  $F, F', \dots, F^{(k)}$  contain an even number  $m$  and  $F, F', \dots, F^{(k-1)}$  contain an odd number  $(m-1)$  of (ii)-differentiable functions  $F^{(i_1)}, F^{(i_2)}, \dots, F^{(i_m)}$  then  $F^{(k)}$  is (ii)-differentiable. By (3.2) of induction hypothesis, we have:

$$[F^{(k)}(t)]^\alpha = [g_\alpha^{(k)}(t), f_\alpha^{(k)}(t)].$$

Since  $F^{(k)}(t)$  is (ii)-differentiable then from definition 2.3, we have:

$$[F^{(k)}(t) \Theta F^{(k)}(t+h)]^\alpha = [g_\alpha^{(k)}(t) - g_\alpha^{(k)}(t+h), f_\alpha^{(k)}(t) - f_\alpha^{(k)}(t+h)],$$

$$[F^{(k)}(t-h) \Theta F^{(k)}(t)]^\alpha = [g_\alpha^{(k)}(t-h) - g_\alpha^{(k)}(t), f_\alpha^{(k)}(t-h) - f_\alpha^{(k)}(t)].$$

and, multiplying by  $\frac{1}{-h}$ ,  $h > 0$  we get:

$$\frac{1}{-h}[F^{(k)}(t) \Theta F^{(k)}(t+h)]^\alpha = \left[ \frac{f_\alpha^{(k)}(t+h) - f_\alpha^{(k)}(t)}{h}, \frac{g_\alpha^{(k)}(t+h) - g_\alpha^{(k)}(t)}{h} \right],$$

and

$$\frac{1}{-h}[F^{(k)}(t-h) \Theta F^{(k)}(t)]^\alpha = \left[ \frac{f_\alpha^{(k)}(t) - f_\alpha^{(k)}(t-h)}{h}, \frac{g_\alpha^{(k)}(t) - g_\alpha^{(k)}(t-h)}{h} \right].$$

Using  $h \rightarrow 0$  on both sides of aforementioned relation, we get the relation (3.3). Then relation (a) is true if  $m$  is an even number such that  $2 \leq m \leq k+1$ . Then (a) is true if  $n = k+1$  and  $m$  is an even number such that  $0 \leq m \leq k+1$ .

Then relation (a) is true for any positive integer  $n$ .

(b) Let  $F, F', \dots, F^{(k)}$  be differentiable fuzzy-valued functions such that  $F^{(i_1)}, F^{(i_2)}, \dots, F^{(i_m)}$  are (ii)-differentiable functions for  $0 \leq i_1 < i_2 < \dots \leq k$ ,  $1 \leq m \leq k+1$  and  $m$  is an odd number. Also, we have two possibilities for the functions  $F, F', \dots, F^{(k-1)}$  as follows:

**Either**  $F, F', \dots, F^{(k)}$  contain an odd number  $m$  and  $F, F', \dots, F^{(k-1)}$  contain an even number  $(m-1)$  of (ii)-differentiable functions  $F^{(i_1)}, F^{(i_2)}, \dots, F^{(i_m)}$  then  $F^{(k)}$  is (ii)-differentiable. By relation (3.1), we get

$$[F^{(k)}(t)]^\alpha = [f_\alpha^{(k)}(t), g_\alpha^{(k)}(t)].$$

Since  $F^{(k)}$  is (ii)-differentiable then from definition 2.3, we have:

$$[F^{(k)}(t) \Theta F^{(k)}(t+h)]^\alpha = [f_\alpha^{(k)}(t) - f_\alpha^{(k)}(t+h), g_\alpha^{(k)}(t) - g_\alpha^{(k)}(t+h)],$$

$$[F^{(k)}(t-h) \Theta F^{(k)}(t)]^\alpha = [f_\alpha^{(k)}(t-h) - f_\alpha^{(k)}(t), g_\alpha^{(k)}(t-h) - g_\alpha^{(k)}(t)].$$

and, multiplying by  $\frac{1}{-h}$ ,  $h > 0$  we get:

$$\frac{1}{-h}[F^{(k)}(t) \Theta F^{(k)}(t+h)]^\alpha = \left[ \frac{g_\alpha^{(k)}(t+h) - g_\alpha^{(k)}(t)}{h}, \frac{f_\alpha^{(k)}(t+h) - f_\alpha^{(k)}(t)}{h} \right],$$

and

$$\frac{1}{-h}[F^{(k)}(t-h) \Theta F^{(k)}(t)]^\alpha = \left[ \frac{g_\alpha^{(k)}(t) - g_\alpha^{(k)}(t-h)}{h}, \frac{f_\alpha^{(k)}(t) - f_\alpha^{(k)}(t-h)}{h} \right].$$

Using  $h \longrightarrow 0$  on both sides of aforementioned relation, we get

$$[F^{(k+1)}(t)]^\alpha = [g_\alpha^{(k+1)}(t), f_\alpha^{(k+1)}(t)] \tag{3.4}$$

**Or**  $F, F', \dots, F^{(k)}$  contain an odd number  $m$  and  $F, F', \dots, F^{(k-1)}$  contain an odd number  $m$  of (ii)-differentiable functions  $F^{(i_1)}, F^{(i_2)}, \dots, F^{(i_m)}$  then  $F^{(k)}$  is (i)-differentiable. By relation (3. 2), we get:

$$[F^{(k)}(t)]^\alpha = [g_\alpha^{(k)}(t), f_\alpha^{(k)}(t)].$$

Since  $F^{(k)}(t)$  is (i)-differentiable then from definition 2.3, we have:

$$[F^{(k)}(t+h) \Theta F^{(k)}(t)]^\alpha = [g_\alpha^{(k)}(t+h) - g_\alpha^{(k)}(t), f_\alpha^{(k)}(t+h) - f_\alpha^{(k)}(t)],$$

$$[F^{(k)}(t) \Theta F^{(k)}(t-h)]^\alpha = [g_\alpha^{(k)}(t) - g_\alpha^{(k)}(t-h), f_\alpha^{(k)}(t) - f_\alpha^{(k)}(t-h)].$$

and , multiplying by  $\frac{1}{h}$ ,  $h > 0$  we get:

$$\frac{1}{h} [F^{(k)}(t+h) \Theta F^{(k)}(t)]^\alpha = \left[ \frac{g_\alpha^{(k)}(t+h) - g_\alpha^{(k)}(t)}{h}, \frac{f_\alpha^{(k)}(t+h) - f_\alpha^{(k)}(t)}{h} \right],$$

and

$$\frac{1}{h} [F^{(k)}(t) \Theta F^{(k)}(t-h)]^\alpha = \left[ \frac{g_\alpha^{(k)}(t) - g_\alpha^{(k)}(t-h)}{h}, \frac{f_\alpha^{(k)}(t) - f_\alpha^{(k)}(t-h)}{h} \right].$$

Finally, using  $h \longrightarrow 0$  on both sides of aforementioned relation, we get the relation (3.4), then the relation (b) is true when  $n = k + 1$  and  $m$  is an odd such that  $1 \leq m \leq k + 1$ .

Then the relation (b) is true for any positive integer  $n$ .

Thus, the theorem is true for any positive integer  $n$ .

**Remark 3.2** If we put  $n=1,2,3,4$  in theorem 3.1, we get the same results given in [7,8,9,9] respectively.

**Theorem 3.3** Suppose that  $g(t), g'(t), \dots, g^{(n-1)}(t)$  be continuous fuzzy-valued functions on  $[0, \infty)$  and of exponential order and that  $g^{(n)}(t)$  is piecewise continuous fuzzy-valued function on  $[0, \infty)$ .

Let  $g^{(i_1)}(t), g^{(i_2)}(t), \dots, g^{(i_m)}(t)$  be (ii)-differentiable functions for  $0 \leq i_1 < i_2 < \dots < i_m \leq n-1$  and  $g^{(p)}$  be (i)-differentiable function for  $p \neq i_j, j = 1, 2, \dots, m$  and  $g(t) = (\underline{g}(t, \alpha), \bar{g}(t, \alpha))$ ; then

(1) If  $m$  is an even number, we have

$$L(g^{(n)}(t)) = s^n L(g(t)) \Theta s^{n-1} g(0) \otimes \sum_{k=1}^{n-1} s^{n-(k+1)} g^{(k)}(0), \tag{3.5}$$

such that

$$\otimes = \begin{cases} \Theta, & \text{if the number of (ii) - differentiable functions } g^{(i)}, \text{ provided} \\ & \text{that } i < k \text{ is an even number} \\ - , & \text{if the number of (ii) - differentiable functions } g^{(i)}, \text{ provided} \\ & \text{that } i < k \text{ is an odd number} \end{cases} \tag{3.6}$$

(2) If  $m$  is an odd number, we have

$$L(g^{(n)}(t)) = -s^{n-1}g(0)\Theta(-s^n)L(g(t)) \otimes \sum_{k=1}^{n-1} s^{n-(k+1)}g^{(k)}(0), \tag{3.7}$$

such that

$$\otimes = \begin{cases} \Theta, & \text{if the number of (ii) - differentiable functions } g^{(i)}, \text{ provided} \\ & \text{that } i < k \text{ is an odd number} \\ - , & \text{if the number of (ii) - differentiable functions } g^{(i)}, \text{ provided} \\ & \text{that } i < k \text{ is an even number} \end{cases} \tag{3.8}$$

**Proof (1)** Let  $g^{(i_1)}(t), g^{(i_2)}(t), \dots, g^{(i_m)}(t)$  be (ii)-differentiable fuzzy valued functions for  $0 \leq i_1 < i_2 < \dots < i_m \leq n-1$  and  $m$  be an even number such that  $2 \leq m \leq n$ , then by theorem 3.1(a), we get

$$g^{(n)}(t) = (\underline{g}^{(n)}(t, \alpha), \overline{g}^{(n)}(t, \alpha)),$$

where  $f_\alpha(t) = \underline{g}(t, \alpha)$  and  $g_\alpha(t) = \overline{g}(t, \alpha)$ . Therefore, we get:

$$\underline{g}^{(n)}(t, \alpha) = \underline{g}^{(n)}(t, \alpha), \overline{g}^{(n)}(t, \alpha) = \overline{g}^{(n)}(t, \alpha). \tag{3.9}$$

Then from (3.9), we get

$$\begin{aligned} L(g^{(n)}(t)) &= L(\underline{g}^{(n)}(t, \alpha), \overline{g}^{(n)}(t, \alpha)) \\ &= (l(\underline{g}^{(n)}(t, \alpha)), l(\overline{g}^{(n)}(t, \alpha))). \end{aligned} \tag{3.10}$$

We know from the ordinary differential equations that:

$$l(\underline{g}^{(n)}(t, \alpha)) = s^n l(\underline{g}(t, \alpha)) - \sum_{i=0}^{n-1} s^{n-(i+1)} \underline{g}^{(i)}(0, \alpha). \tag{3.11}$$

Equation (3.11) can be written as:

$$\begin{aligned} l(\underline{g}^{(n)}(t, \alpha)) &= s^n l(\underline{g}(t, \alpha)) - s^{n-1} \underline{g}(0, \alpha) - \sum_{k=1}^{i_1} s^{n-(k+1)} \underline{g}^{(k)}(0, \alpha) - \sum_{k=i_1+1}^{i_2} s^{n-(k+1)} \underline{g}^{(k)}(0, \alpha) \\ &\quad - \sum_{k=i_2+1}^{i_3} s^{n-(k+1)} \underline{g}^{(k)}(0, \alpha) - \dots - \sum_{k=i_m+1}^{n-1} s^{n-(k+1)} \underline{g}^{(k)}(0, \alpha). \end{aligned} \tag{3.12}$$

In a similar manner, we can get

$$\begin{aligned} l(\overline{g}^{(n)}(t, \alpha)) &= s^n l(\overline{g}(t, \alpha)) - s^{n-1} \overline{g}(0, \alpha) - \sum_{k=1}^{i_1} s^{n-(k+1)} \overline{g}^{(k)}(0, \alpha) - \sum_{k=i_1+1}^{i_2} s^{n-(k+1)} \overline{g}^{(k)}(0, \alpha) \\ &\quad - \sum_{k=i_2+1}^{i_3} s^{n-(k+1)} \overline{g}^{(k)}(0, \alpha) - \dots - \sum_{k=i_m+1}^{n-1} s^{n-(k+1)} \overline{g}^{(k)}(0, \alpha). \end{aligned} \tag{3.13}$$

Since  $0 \leq i_1 < i_2 < \dots < i_m \leq n-1$  we can apply theorem 3.1 for each  $g^{(k)}(t)$  where  $1 \leq k \leq n-1$  as follows:

$$\begin{aligned} \underline{g}^{(k)}(0, \alpha) &= \underline{g}^{(k)}(0, \alpha), \overline{g}^{(k)}(0, \alpha) = \overline{g}^{(k)}(0, \alpha), \quad 1 \leq k \leq i_1, \\ \overline{g}^{(k)}(0, \alpha) &= \underline{g}^{(k)}(0, \alpha), \underline{g}^{(k)}(0, \alpha) = \overline{g}^{(k)}(0, \alpha), \quad i_1+1 \leq k \leq i_2, \\ \underline{g}^{(k)}(0, \alpha) &= \underline{g}^{(k)}(0, \alpha), \overline{g}^{(k)}(0, \alpha) = \overline{g}^{(k)}(0, \alpha), \quad i_2+1 \leq k \leq i_3, \\ &\vdots \\ \underline{g}^{(k)}(0, \alpha) &= \underline{g}^{(k)}(0, \alpha), \overline{g}^{(k)}(0, \alpha) = \overline{g}^{(k)}(0, \alpha), \quad i_m+1 \leq k \leq n-1. \end{aligned} \tag{3.14}$$

The last one of the equations in (3.14) yields from theorem 3.1(a) because  $m$  is an even number. Using (3.12), (3.13) and (3.14), equation (3.10) becomes

$$\begin{aligned}
 L(g^{(n)}(t)) &= (s^n l(\underline{g}(t, \alpha)) - s^{n-1} \underline{g}(0, \alpha) - \sum_{k=1}^{i_1} s^{n-(k+1)} \underline{g}^{(k)}(0, \alpha) - \sum_{k=i_1+1}^{i_2} s^{n-(k+1)} \overline{g}^{(k)}(0, \alpha) - \\
 &\quad \sum_{k=i_2+1}^{i_3} s^{n-(k+1)} \underline{g}^{(k)}(0, \alpha) - \dots - \sum_{k=i_m+1}^{n-1} s^{n-(k+1)} \underline{g}^{(k)}(0, \alpha), s^n l(\overline{g}(t, \alpha)) - s^{n-1} \overline{g}(0, \alpha) \\
 &\quad - \sum_{k=1}^{i_1} s^{n-(k+1)} \overline{g}^{(k)}(0, \alpha) - \sum_{k=i_1+1}^{i_2} s^{n-(k+1)} \underline{g}^{(k)}(0, \alpha) - \sum_{k=i_2+1}^{i_3} s^{n-(k+1)} \overline{g}^{(k)}(0, \alpha) - \dots - \\
 &\quad \sum_{k=i_m+1}^{n-1} s^{n-(k+1)} \overline{g}^{(k)}(0, \alpha)). \\
 &= s^n L(g(t)) \Theta s^{n-1} g(0) \Theta \sum_{k=1}^{i_1} s^{n-(k+1)} g^{(k)}(0) - \sum_{k=i_1+1}^{i_2} s^{n-(k+1)} g^{(k)}(0) \\
 &\quad \Theta \sum_{k=i_2+1}^{i_3} s^{n-(k+1)} g^{(k)}(0) - \dots - \Theta \sum_{k=i_m+1}^{n-1} s^{n-(k+1)} g^{(k)}(0) \\
 &= s^n L(g(t)) \Theta s^{n-1} g(0) \otimes \sum_{k=1}^{n-1} s^{n-(k+1)} g^{(k)}(0),
 \end{aligned}$$

where  $\otimes$  is defined as in (3.6). Then, the theorem is true for any even number  $m$  such that  $2 \leq m \leq n$

We note that if  $m = 0$  we have  $g(t), g'(t), \dots, g^{(n-1)}(t)$  are (i)-differentiable functions, then equation (3.10) becomes:

$$\begin{aligned}
 L(g^{(n)}(t)) &= (s^n l(\underline{g}(t, \alpha)) - \sum_{k=0}^{n-1} s^{n-(k+1)} \underline{g}^{(k)}(0, \alpha), s^n l(\overline{g}(t, \alpha)) - \sum_{k=0}^{n-1} s^{n-(k+1)} \overline{g}^{(k)}(0, \alpha)) \\
 &= s^n L(g(t)) \Theta \sum_{k=0}^{n-1} s^{n-(k+1)} g^{(k)}(0).
 \end{aligned}$$

It is clear that the above relation reconciles with the relations (3.5) and (3.6).

Therefore, the theorem is true for any even number  $m$  such that  $0 \leq m \leq n$

**Proof (2)** Let  $g^{(i_1)}(t), g^{(i_2)}(t), \dots, g^{(i_m)}(t)$  be (ii)-differentiable fuzzy valued functions for  $0 \leq i_1 < i_2 < \dots < i_m \leq n-1$  and  $m$  be an odd number such that  $1 \leq m \leq n$  and, then by theorem 3.1(b) we get

$$g^{(n)}(t) = (\overline{g}^{(n)}(t, \alpha), \underline{g}^{(n)}(t, \alpha)).$$

Therefore, we get:

$$\underline{g}^{(n)}(t, \alpha) = \overline{g}^{(n)}(t, \alpha), \overline{g}^{(n)}(t, \alpha) = \underline{g}^{(n)}(t, \alpha). \tag{3.15}$$

Therefore, from (3.15) we get

$$\begin{aligned}
 L(g^{(n)}(t)) &= L(\underline{g}^{(n)}(t, \alpha), \overline{g}^{(n)}(t, \alpha)) \\
 &= (l(\overline{g}^{(n)}(t, \alpha)), l(\underline{g}^{(n)}(t, \alpha))).
 \end{aligned} \tag{3.16}$$

Since  $0 \leq i_1 < i_2 < \dots < i_m \leq n-1$  we can apply theorem 3.1 for each  $g^{(k)}(t)$  where  $1 \leq k \leq n-1$  as follows:

$$\begin{aligned}
 \underline{g}^{(k)}(0, \alpha) &= \underline{g}^{(k)}(0, \alpha), \overline{g}^{(k)}(0, \alpha) = \overline{g}^{(k)}(0, \alpha), \quad 1 \leq k \leq i_1, \\
 \overline{g}^{(k)}(0, \alpha) &= \underline{g}^{(k)}(0, \alpha), \underline{g}^{(k)}(0, \alpha) = \overline{g}^{(k)}(0, \alpha), \quad i_1 + 1 \leq k \leq i_2, \\
 \underline{g}^{(k)}(0, \alpha) &= \underline{g}^{(k)}(0, \alpha), \overline{g}^{(k)}(0, \alpha) = \overline{g}^{(k)}(0, \alpha), \quad i_2 + 1 \leq k \leq i_3, \\
 &\vdots
 \end{aligned} \tag{3.17}$$

$$\bar{g}^{(k)}(0, \alpha) = \underline{g}^{(k)}(0, \alpha), \underline{g}^{(k)}(0, \alpha) = \overline{g}^{(k)}(0, \alpha), \quad i_m + 1 \leq k \leq n - 1.$$

The last one of the equations in (3.17) yields from theorem 3.1(b) because  $m$  is an odd number. Using (3. 12), (3.13) and (3. 17), equation (3. 16) becomes

$$\begin{aligned} L(g^{(n)}(t)) = & s^n l(\bar{g}(t, \alpha)) - s^{n-1} \bar{g}(0, \alpha) - \sum_{k=1}^{i_1} s^{n-(k+1)} \overline{g}^{(k)}(0, \alpha) - \sum_{k=i_1+1}^{i_2} s^{n-(k+1)} \underline{g}^{(k)}(0, \alpha) - \\ & \sum_{k=i_2+1}^{i_3} s^{n-(k+1)} \overline{g}^{(k)}(0, \alpha) - \dots - \sum_{k=i_m+1}^{n-1} s^{n-(k+1)} \underline{g}^{(k)}(0, \alpha), s^n l(\underline{g}(t, \alpha)) - s^{n-1} \underline{g}(0, \alpha) \\ & - \sum_{k=1}^{i_1} s^{n-(k+1)} \underline{g}^{(k)}(0, \alpha) - \sum_{k=i_1+1}^{i_2} s^{n-(k+1)} \overline{g}^{(k)}(0, \alpha) - \sum_{k=i_2+1}^{i_3} s^{n-(k+1)} \underline{g}^{(k)}(0, \alpha) - \dots - \\ & \sum_{k=i_m+1}^{n-1} s^{n-(k+1)} \overline{g}^{(k)}(0, \alpha). \end{aligned}$$

Thus

$$\begin{aligned} L(g^{(n)}(t)) = & -s^{n-1} g(0) \Theta (-s^n) L(g(t)) - \sum_{k=1}^{i_1} s^{n-(k+1)} g^{(k)}(0) \Theta \sum_{k=i_1+1}^{i_2} s^{n-(k+1)} g^{(k)}(0) \\ & - \sum_{k=i_2+1}^{i_3} s^{n-(k+1)} g^{(k)}(0) \Theta \dots \Theta \sum_{k=i_m+1}^{n-1} s^{n-(k+1)} g^{(k)}(0) \\ = & -s^{n-1} g(0) \Theta (-s^n) L(g(t)) \otimes \sum_{k=1}^{n-1} s^{n-(1+k)} g^{(k)}(0), \end{aligned}$$

where  $\otimes$  is defined as in (3. 8)

**Remark 3.4** If we put  $n=1,2,3,4$  in theorem 3.3, we get the same results given in [7,8,9,9] respectively.

**Remark 3.5** By applying theorem 3.3, we note that: There are  $2^n$  cases for fuzzy Laplace transforms for  $g^{(n)}(t), n \in \mathbb{Z}^+$ . Also, we have

$$\sum_{k=0}^n \binom{n}{k} = 2^n,$$

where  $\binom{n}{k}$  is the number of cases that contain  $k$  functions of the type (ii)-differentiable functions among the functions  $g(t), g'(t), \dots, g^{(n-1)}(t)$ .

#### 4. Application of the Generalization

In this section, we present an example of the third order to show the validity of the generalization given in theorem 3.3.



**Example 4.1** Consider the following third-order FIVP:

$$y'''(t) = -y''(t) - 3y'(t) + 5y(t), \tag{4.1}$$

$$y(0) = \left(\frac{3}{4} + \frac{r}{4}, \frac{5}{4} - \frac{r}{4}\right), \quad y'(0) = \left(\frac{3}{2} + \frac{r}{2}, \frac{5}{2} - \frac{r}{2}\right), \quad y''(0) = \left(\frac{15}{4} + \frac{r}{4}, \frac{17}{4} - \frac{r}{4}\right).$$

We note that:

$$\underline{y}(0, r) = \frac{3}{4} + \frac{r}{4}, \bar{y}(0, r) = \frac{5}{4} - \frac{r}{4}, \underline{y}'(0, r) = \frac{3}{2} + \frac{r}{2}, \bar{y}'(0, r) = \frac{5}{2} - \frac{r}{2}, \underline{y}''(0, r) = \frac{15}{4} + \frac{r}{4}, \bar{y}''(0, r) = \frac{17}{4} - \frac{r}{4}$$

By taking fuzzy Laplace transform for both sides of equation (4.1), we get:

$$L[y'''(t)] = L[-y''(t) - 3y'(t) + 5y(t)] \tag{4.2}$$

Now, we shall use theorem 3.3 to find Laplace transform for each term in equation (4.2), therefore we have  $2^3 = 8$  cases as follows:

**Case 1** Let us consider  $y(t), y'(t)$  and  $y''(t)$  be (i)-differentiable. Then equation (4.2) becomes:

$$s^3 L(y(t)) \ominus s^2 y(0) \ominus s y'(0) \ominus y''(0) = -[s^2 L(y(t)) \ominus s y(0) \ominus y'(0)] - 3 [s L(y(t)) \ominus y(0)] + 5L(y(t)),$$

then, we get the system:

$$\begin{aligned} (s^3 - 5)l(\underline{y}(t, r)) + (s^2 + 3s)l(\bar{y}(t, r)) &= \left(\frac{3}{4} + \frac{r}{4}\right)s^2 + \left(\frac{11}{4} + \frac{r}{4}\right)s + 10 - r, \\ (s^3 - 5)l(\bar{y}(t, r)) + (s^2 + 3s)l(\underline{y}(t, r)) &= \left(\frac{5}{4} - \frac{r}{4}\right)s^2 + \left(\frac{13}{4} - \frac{r}{4}\right)s + 8 + r. \end{aligned} \tag{4.3}$$

The solution of system (4.3) is as follows:

$$l(\underline{y}(t, r)) = \frac{(r + 3)s^5 + (2r + 6)s^4 + 12s^3 - (6r + 86)s^2 - (17r + 151)s + 20r - 200}{4(s^3 + s^2 + 3s - 5)(s^3 - s^2 - 3s - 5)},$$

$$l(\bar{y}(t, r)) = \frac{(5 - r)s^5 + (10 - 2r)s^4 + 12s^3 + (6r - 98)s^2 + (17r - 185)s - 20r - 160}{4(s^3 + s^2 + 3s - 5)(s^3 - s^2 - 3s - 5)}.$$

To make a cubic partition for  $l(\underline{y}(t, r))$  and  $l(\bar{y}(t, r))$ , we suppose that:

$$\frac{a_5 s^5 + a_4 s^4 + a_3 s^3 + a_2 s^2 + a_1 s + a_0}{4(s^3 + s^2 + 3s - 5)(s^3 + b_1 s^2 + b_2 s + b_3)} = \frac{1}{4} \left[ \frac{As^2 + Bs + C}{s^3 + s^2 + 3s - 5} + \frac{Ds^2 + Es + F}{s^3 + b_1 s^2 + b_2 s + b_3} \right], \tag{4.4}$$

yields the system:

$$\begin{aligned} A + D &= a_5, \\ b_1 A + B + D + E &= a_4, \\ b_2 A + b_1 B + C + 3D + E + F &= a_3, \\ b_3 A + b_2 B + b_1 C - 5D + 3E + F &= a_2, \\ b_3 B + b_2 C - 5E + 3F &= a_1, \\ b_3 C - 5F &= a_0. \end{aligned} \tag{4.5}$$

To get the cubic partition for  $l(\underline{y}(t, r))$ , we put  $a_0 = 20r - 200, a_1 = -151 - 17r, a_2 = -86 - 6r, a_3 = 12, a_4 = 2r + 6, a_5 = r + 3, b_1 = -1, b_2 = -3$  and  $b_3 = -5$  in the system (4.5), then we get:

$$l(\underline{y}(t, r)) = \frac{s^2 + 3s + 9}{(s - 1)(s^2 + 2s + 5)} + \frac{(r - 1)(s^2 + s - 4)}{4(s^3 - s^2 - 3s - 5)}. \quad (4.6)$$

Similarly, to get the partition for  $l(\bar{y}(t, r))$ , we put  $a_0 = -20r - 160, a_1 = 17r - 185, a_2 = 6r - 98, a_3 = 12, a_4 = 10 - 2r, a_5 = 5 - r, b_1 = -1, b_2 = -3$  and  $b_3 = -5$  in the system (4.5), then:

$$l(\bar{y}(t, r)) = \frac{s^2 + 3s + 9}{(s - 1)(s^2 + 2s + 5)} - \frac{(r - 1)(s^2 + s - 4)}{4(s^3 - s^2 - 3s - 5)}. \quad (4.7)$$

Now, by finding a root for the cubic polynomial given in (4.6) and (4.7), we get:

$$\begin{aligned} \underline{y}(t, r) &= l^{-1}\left(\frac{s^2 + 3s + 9}{(s - 1)(s^2 + 2s + 5)}\right) + \frac{r - 1}{4} l^{-1}\left(\frac{s^2 + s - 4}{(s - a)(s^2 + bs + c)}\right) \\ \bar{y}(t, r) &= l^{-1}\left(\frac{s^2 + 3s + 9}{(s - 1)(s^2 + 2s + 5)}\right) - \frac{r - 1}{4} l^{-1}\left(\frac{s^2 + s - 4}{(s - a)(s^2 + bs + c)}\right) \end{aligned} \quad (4.8)$$

where

$$\begin{aligned} a &= \frac{1}{3} + \frac{1}{3} \sqrt[3]{82 + 6\sqrt{159}} + \frac{1}{3} \sqrt[3]{82 - 6\sqrt{159}}, \\ b &= \frac{-2}{3} + \frac{1}{3} \sqrt[3]{82 + 6\sqrt{159}} + \frac{1}{3} \sqrt[3]{82 - 6\sqrt{159}}, \\ c &= -1 - \frac{1}{9} \sqrt[3]{82 + 6\sqrt{159}} - \frac{1}{9} \sqrt[3]{82 - 6\sqrt{159}} + \frac{1}{9} (82 + 6\sqrt{159})^{\frac{2}{3}} + \frac{1}{9} (82 - 6\sqrt{159})^{\frac{2}{3}}. \end{aligned} \quad (4.9)$$

To find  $l^{-1}$  in the system (4.8), we put:

$$\frac{d_2 s^2 + d_1 s + d_0}{(s - a)(s^2 + bs + c)} = \frac{(-d_0 - ad_1 + cd_2 + abd_2)s + (d_1 c - d_0 a - d_0 b + ad_2 c)}{(a^2 + ab + c)(s^2 + bs + c)} + \frac{d_2 a^2 + d_1 a + d_0}{a^2 + ab + c} \frac{1}{s - a}.$$

Then:

$$\begin{aligned} l^{-1}\left(\frac{d_2 s^2 + d_1 s + d_0}{(s - a)(s^2 + bs + c)}\right) &= \frac{d_2 a^2 + d_1 a + d_0}{a^2 + ab + c} e^{at} - \frac{d_0 + ad_1 - cd_2 - abd_2}{a^2 + ab + c} e^{-\frac{b}{2}t} \cosh \sqrt{\frac{b^2}{4} - ct} \\ &+ \frac{d_0 + ad_1 - cd_2 - abd_2}{(a^2 + ab + c) \sqrt{\frac{b^2}{4} - c}} \left(\frac{b}{2} - \frac{ad_0 + bd_0 - cd_1 - acd_2}{d_0 + ad_1 - cd_2 - abd_2}\right) e^{-\frac{b}{2}t} \sinh \sqrt{\frac{b^2}{4} - ct}. \end{aligned} \quad (4.10)$$

By using relation (4.10), then  $r$  - cut representation of solution given in equation (4.8) becomes:

$$\underline{y}(t, r) = \frac{13}{8} e^t - \frac{5}{8} e^{-t} \cos 2t - \frac{1}{8} e^{-t} \sin 2t + (r - 1)(c_1 e^{at} - c_2 e^{\frac{-b}{2}t} \cosh c_4 t + c_3 e^{\frac{-b}{2}t} \sinh c_4 t),$$

$$\bar{y}(t, r) = \frac{13}{8}e^t - \frac{5}{8}e^{-t} \cos 2t - \frac{1}{8}e^{-t} \sin 2t - (r-1)(c_1 e^{a t} - c_2 e^{\frac{-b}{2} t} \cosh c_4 t + c_3 e^{\frac{-b}{2} t} \sinh c_4 t),$$

where

$$c_1 = \frac{-12 + 5\sqrt[3]{82 + 6\sqrt{159}} + 5\sqrt[3]{82 - 6\sqrt{159}} + (82 + 6\sqrt{159})^{\frac{2}{3}} + (82 - 6\sqrt{159})^{\frac{2}{3}}}{4 [30 + 3(82 + 6\sqrt{159})^{\frac{2}{3}} + 3(82 - 6\sqrt{159})^{\frac{2}{3}}]},$$

$$c_2 = \frac{-42 + 5\sqrt[3]{82 + 6\sqrt{159}} + 5\sqrt[3]{82 - 6\sqrt{159}} - 2(82 + 6\sqrt{159})^{\frac{2}{3}} - 2(82 - 6\sqrt{159})^{\frac{2}{3}}}{4 [30 + 3(82 + 6\sqrt{159})^{\frac{2}{3}} + 3(82 - 6\sqrt{159})^{\frac{2}{3}}]},$$

$$c_3 = \frac{6[22\sqrt[3]{82 + 6\sqrt{159}} + 22\sqrt[3]{82 - 6\sqrt{159}} + 5(82 + 6\sqrt{159})^{\frac{2}{3}} + 5(82 - 6\sqrt{159})^{\frac{2}{3}}] c_4}{4 [30 + 3(82 + 6\sqrt{159})^{\frac{2}{3}} + 3(82 - 6\sqrt{159})^{\frac{2}{3}}][20 - (82 + 6\sqrt{159})^{\frac{2}{3}} - (82 - 6\sqrt{159})^{\frac{2}{3}}]}$$

$$c_4 = \frac{1}{6} \sqrt{60 - 3(82 + 6\sqrt{159})^{\frac{2}{3}} - 3(82 - 6\sqrt{159})^{\frac{2}{3}}}.$$

and  $a, b$  and  $c$  are defined as in (4.5).

**Case 2** Let  $y'(t)$  and  $y''(t)$  be (i)-differentiable and  $y(t)$  be (ii)-differentiable. Then equation (4.2) becomes:

$$-s^2 y(0) \Theta(-s^3) L(y(t)) \Theta s y'(t) \Theta y''(0) = -[-s y(0) \Theta(-s^2) L(y(t)) \Theta y'(t)] - 3[-y(0) \Theta(-s) L(y(t))] + 5L(y(t)),$$

then, we get the system:

$$\begin{aligned} s^3 l(y(t, r)) + (s^2 + 3s - 5) l(\bar{y}(t, r)) &= \left(\frac{3}{4} + \frac{r}{4}\right) s^2 + \left(\frac{15}{4} - \frac{3r}{4}\right) s + \frac{19}{2} - \frac{r}{2}, \\ s^3 l(\bar{y}(t, r)) + (s^2 + 3s - 5) l(y(t, r)) &= \left(\frac{5}{4} - \frac{r}{4}\right) s^2 + \left(\frac{9}{4} + \frac{3r}{4}\right) s + \frac{17}{2} + \frac{r}{2}. \end{aligned} \tag{4.11}$$

By achieving the same steps given in case 1, we get:

$$\begin{aligned} \underline{y}(t, r) &= l^{-1}\left(\frac{s^2 + 3s + 9}{(s-1)(s^2 + 2s + 5)}\right) + \frac{r-1}{4} l^{-1}\left(\frac{s^2 - 3s - 2}{(s-a)(s^2 + bs + c)}\right), \\ \bar{y}(t, r) &= l^{-1}\left(\frac{s^2 + 3s + 9}{(s-1)(s^2 + 2s + 5)}\right) - \frac{r-1}{4} l^{-1}\left(\frac{s^2 - 3s - 2}{(s-a)(s^2 + bs + c)}\right). \end{aligned} \tag{4.12}$$

where

$$\begin{aligned}
 a &= \frac{1}{3} + \frac{1}{3} \sqrt[3]{-53 + 3\sqrt{201}} + \frac{1}{3} \sqrt[3]{-53 - 3\sqrt{201}}, \\
 b &= \frac{-2}{3} + \frac{1}{3} \sqrt[3]{-53 + 3\sqrt{201}} + \frac{1}{3} \sqrt[3]{-53 - 3\sqrt{201}}, \\
 c &= -1 - \frac{1}{9} \sqrt[3]{-53 + 3\sqrt{201}} - \frac{1}{9} \sqrt[3]{-53 - 3\sqrt{201}} + \frac{1}{9} (-53 + 3\sqrt{201})^{\frac{2}{3}} + \frac{1}{9} (-53 - 3\sqrt{201})^{\frac{2}{3}}.
 \end{aligned}
 \tag{4.13}$$

By using relation (4.10), then  $r$  – cut representation of solution given in equation (4.12) becomes:

$$\begin{aligned}
 \underline{y}(t, r) &= \frac{13}{8} e^t - \frac{5}{8} e^{-t} \cos 2t - \frac{1}{8} e^{-t} \sin 2t + (r-1)(c_1 e^{at} - c_2 e^{\frac{-b}{2}t} \cosh c_4 t + c_3 e^{\frac{-b}{2}t} \sinh c_4 t), \\
 \bar{y}(t, r) &= \frac{13}{8} e^t - \frac{5}{8} e^{-t} \cos 2t - \frac{1}{8} e^{-t} \sin 2t - (r-1)(c_1 e^{at} - c_2 e^{\frac{-b}{2}t} \cosh c_4 t + c_3 e^{\frac{-b}{2}t} \sinh c_4 t),
 \end{aligned}$$

where

$$\begin{aligned}
 c_1 &= \frac{-6 - 7\sqrt[3]{-53 + 3\sqrt{201}} - 7\sqrt[3]{-53 - 3\sqrt{201}} + (-53 + 3\sqrt{201})^{\frac{2}{3}} + (-53 - 3\sqrt{201})^{\frac{2}{3}}}{4[30 + 3(-53 + 3\sqrt{201})^{\frac{2}{3}} + 3(-53 - 3\sqrt{201})^{\frac{2}{3}}]}, \\
 c_2 &= \frac{-36 - 7\sqrt[3]{-53 + 3\sqrt{201}} - 7\sqrt[3]{-53 - 3\sqrt{201}} - 2(-53 + 3\sqrt{201})^{\frac{2}{3}} - 2(-53 - 3\sqrt{201})^{\frac{2}{3}}}{4 [30 + 3(-53 + 3\sqrt{201})^{\frac{2}{3}} + 3(-53 - 3\sqrt{201})^{\frac{2}{3}}]}, \\
 c_3 &= \frac{6[16\sqrt[3]{-53 + 3\sqrt{201}} + 16\sqrt[3]{-53 - 3\sqrt{201}} - 7(-53 + 3\sqrt{201})^{\frac{2}{3}} - 7(-53 - 3\sqrt{201})^{\frac{2}{3}}] c_4}{4 [30 + 3(-53 + 3\sqrt{201})^{\frac{2}{3}} + 3(-53 + 3\sqrt{201})^{\frac{2}{3}}][20 - (-53 + 3\sqrt{201})^{\frac{2}{3}} - (-53 - 3\sqrt{201})^{\frac{2}{3}}]}, \\
 c_4 &= \frac{1}{6} \sqrt{60 - 3(-53 + 3\sqrt{201})^{\frac{2}{3}} - 3(-53 - 3\sqrt{201})^{\frac{2}{3}}}.
 \end{aligned}$$

and  $a, b$  and  $c$  are defined as in (4.13).

**Case 3** Let  $y(t)$  and  $y''(t)$  be (i)-differentiable and  $y'(t)$  be (ii)- differentiable. Then equation (4.2) becomes:

$$-s^2 y(0) \Theta(-s^3) L(y(t)) - s y'(t) \Theta y''(0) = [-s y(0) \Theta(-s^2) L(y(t)) - y'(t)] - 3 [s L(y(t)) \Theta y(0)] + 5 L(y(t)),$$

then, we get the system:

$$\begin{aligned}
 (s^3 + 3s)l(\underline{y}(t, r)) + (s^2 - 5)l(\bar{y}(t, r)) &= \left(\frac{3}{4} + \frac{r}{4}\right)s^2 + \left(\frac{11}{4} + \frac{r}{4}\right)s + 9, \\
 (s^3 + 3s)l(\bar{y}(t, r)) + (s^2 - 5)l(\underline{y}(t, r)) &= \left(\frac{5}{4} - \frac{r}{4}\right)s^2 + \left(\frac{13}{4} - \frac{r}{4}\right)s + 9.
 \end{aligned}
 \tag{4.14}$$

The solution of system (4.14) is as follows:

$$l(\underline{y}(t, r)) = \frac{13}{8(s-1)} - \frac{5s+7}{8(s^2+2s+5)} + \frac{(r-1)s}{4(s^2-2s+5)},$$

$$l(\bar{y}(t, r)) = \frac{13}{8(s-1)} - \frac{5s+7}{8(s^2+2s+5)} - \frac{(r-1)s}{4(s^2-2s+5)}.$$

Then, we get  $r$  – cut representation of solution as follows:

$$\underline{y}(t, r) = \frac{13}{8}e^t - \frac{5}{8}e^{-t} \cos 2t - \frac{1}{8}e^{-t} \sin 2t + (r-1)\left(\frac{1}{4}e^t \cos 2t + \frac{1}{8}e^t \sin 2t\right),$$

$$\bar{y}(t, r) = \frac{13}{8}e^t - \frac{5}{8}e^{-t} \cos 2t - \frac{1}{8}e^{-t} \sin 2t - (r-1)\left(\frac{1}{4}e^t \cos 2t + \frac{1}{8}e^t \sin 2t\right).$$

**Case 4** Let  $y(t)$  and  $y'(t)$  be (i)-differentiable and  $y''(t)$  be (ii)- differentiable. Then equation (4.2) becomes:

$$-s^2y(0)\Theta(-s^3)L(y(t)) - sy'(t) - y''(0) = -[s^2L(y(t))\Theta sy(0)\Theta y'(t)] - 3[sL(y(t))\Theta y(0)] + 5L(y(t)),$$

then, we get the system:

$$(s^3 + s^2 + 3s)l(\underline{y}(t, r)) - 5l(\bar{y}(t, r)) = \left(\frac{3}{4} + \frac{r}{4}\right)s^2 + \left(\frac{9}{4} + \frac{3r}{4}\right)s + \frac{15}{2} + \frac{3r}{2}, \tag{4.15}$$

$$(s^3 + s^2 + 3s)l(\bar{y}(t, r)) - 5l(\underline{y}(t, r)) = \left(\frac{5}{4} - \frac{r}{4}\right)s^2 + \left(\frac{15}{4} - \frac{3r}{4}\right)s + \frac{21}{2} - \frac{3r}{2}.$$

By achieving the same steps given in case 1, we get:

$$\underline{y}(t, r) = l^{-1}\left(\frac{s^2 + 3s + 9}{(s-1)(s^2 + 2s + 5)}\right) + \frac{r-1}{4}l^{-1}\left(\frac{s^2 + 3s + 6}{(s-a)(s^2 + bs + c)}\right), \tag{4.16}$$

$$\bar{y}(t, r) = l^{-1}\left(\frac{s^2 + 3s + 9}{(s-1)(s^2 + 2s + 5)}\right) - \frac{r-1}{4}l^{-1}\left(\frac{s^2 + 3s + 6}{(s-a)(s^2 + bs + c)}\right).$$

where

$$a = \frac{-1}{3} + \frac{1}{3}\sqrt[3]{-55 + 3\sqrt{393}} + \frac{1}{3}\sqrt[3]{-55 - 3\sqrt{393}},$$

$$b = \frac{2}{3} + \frac{1}{3}\sqrt[3]{-55 + 3\sqrt{393}} + \frac{1}{3}\sqrt[3]{-55 - 3\sqrt{393}}, \tag{4.17}$$

$$c = 1 + \frac{1}{9}\sqrt[3]{-55 + 3\sqrt{393}} + \frac{1}{9}\sqrt[3]{-55 - 3\sqrt{393}} + \frac{1}{9}(-55 + 3\sqrt{393})^{\frac{2}{3}} + \frac{1}{9}(-55 - 3\sqrt{393})^{\frac{2}{3}}.$$

By using relation (4.10), then  $r$  – cut representation of solution given in equation (4.16) becomes:

$$\underline{y}(t, r) = \frac{13}{8}e^t - \frac{5}{8}e^{-t} \cos 2t - \frac{1}{8}e^{-t} \sin 2t + (r-1)(c_1e^{at} - c_2e^{\frac{-b}{2}t} \cosh c_4 t + c_3e^{\frac{-b}{2}t} \sinh c_4 t),$$

$$\bar{y}(t, r) = \frac{13}{8}e^t - \frac{5}{8}e^{-t} \cos 2t - \frac{1}{8}e^{-t} \sin 2t - (r-1)(c_1e^{at} - c_2e^{\frac{-b}{2}t} \cosh c_4 t + c_3e^{\frac{-b}{2}t} \sinh c_4 t),$$

where

$$c_1 = \frac{30 + 7\sqrt[3]{-55 + 3\sqrt{393}} + 7\sqrt[3]{-55 - 3\sqrt{393}} + (-55 + 3\sqrt{393})^{\frac{2}{3}} + (-55 - 3\sqrt{393})^{\frac{2}{3}}}{4[-24 + 3(-55 + 3\sqrt{393})^{\frac{2}{3}} + 3(-55 - 3\sqrt{393})^{\frac{2}{3}}]},$$

$$c_2 = \frac{54 + 7\sqrt[3]{-55 + 3\sqrt{393}} + 7\sqrt[3]{-55 - 3\sqrt{393}} - 2(-55 + 3\sqrt{393})^{\frac{2}{3}} - 2(-55 - 3\sqrt{393})^{\frac{2}{3}}}{4[-24 + 3(-55 + 3\sqrt{393})^{\frac{2}{3}} + 3(-55 - 3\sqrt{393})^{\frac{2}{3}}]},$$

$$c_3 = \frac{6[-38\sqrt[3]{-55 + 3\sqrt{393}} - 38\sqrt[3]{-55 - 3\sqrt{393}} + 7(-55 + 3\sqrt{393})^{\frac{2}{3}} + 7(-55 - 3\sqrt{393})^{\frac{2}{3}}] c_4}{4[-24 + 3(-55 + 3\sqrt{393})^{\frac{2}{3}} + 3(-55 - 3\sqrt{393})^{\frac{2}{3}}][ -16 - (-55 + 3\sqrt{393})^{\frac{2}{3}} - (-55 - 3\sqrt{393})^{\frac{2}{3}}]},$$

$$c_4 = \frac{1}{6} \sqrt{-48 - 3(-55 + 3\sqrt{393})^{\frac{2}{3}} - 3(-55 - 3\sqrt{393})^{\frac{2}{3}}}.$$

and  $a, b$  and  $c$  are defined as in (4.17).

**Case 5** Let  $y''(t)$  be (i)-differentiable and  $y(t)$  and  $y'(t)$  be (ii)- differentiable. Then equation (4.2) becomes:

$$s^3 L(y(t)) \Theta s^2 y(0) - s y'(t) \Theta y''(0) = -[s^2 L(y(t)) \Theta s y(0) - y'(t)] - 3[-y(0) \Theta (-s)L(y(t))] + 5L(y(t)),$$

then, we get the system:

$$(s^3 + 3s - 5)l(\underline{y}(t, r)) + s^2 l(\bar{y}(t, r)) = \left(\frac{3}{4} + \frac{r}{4}\right)s^2 + \left(\frac{15}{4} - \frac{3r}{4}\right)s + \frac{15}{2} + \frac{3r}{2},$$

$$(s^3 + 3s - 5)l(\bar{y}(t, r)) + s^2 l(\underline{y}(t, r)) = \left(\frac{5}{4} - \frac{r}{4}\right)s^2 + \left(\frac{9}{4} + \frac{3r}{4}\right)s + \frac{21}{2} - \frac{3r}{2}.$$
(4.18)

By achieving the same steps given in case 1, we get:

$$\underline{y}(t, r) = l^{-1}\left(\frac{s^2 + 3s + 9}{(s-1)(s^2 + 2s + 5)}\right) + \frac{r-1}{4} l^{-1}\left(\frac{s^2 - 3s + 6}{(s-a)(s^2 + bs + c)}\right),$$

$$\bar{y}(t, r) = l^{-1}\left(\frac{s^2 + 3s + 9}{(s-1)(s^2 + 2s + 5)}\right) - \frac{r-1}{4} l^{-1}\left(\frac{s^2 - 3s + 6}{(s-a)(s^2 + bs + c)}\right).$$
(4.19)

where

$$a = \frac{1}{3} + \frac{1}{3}\sqrt[3]{55 + 3\sqrt{393}} + \frac{1}{3}\sqrt[3]{55 - 3\sqrt{393}},$$

$$b = \frac{-2}{3} + \frac{1}{3}\sqrt[3]{55 + 3\sqrt{393}} + \frac{1}{3}\sqrt[3]{55 - 3\sqrt{393}},$$

$$c = 1 - \frac{1}{9}\sqrt[3]{55 + 3\sqrt{393}} - \frac{1}{9}\sqrt[3]{55 - 3\sqrt{393}} + \frac{1}{9}(55 + 3\sqrt{393})^{\frac{2}{3}} + \frac{1}{9}(55 - 3\sqrt{393})^{\frac{2}{3}}.$$
(4.20)

By using relation (4.10), then  $r$  – cut representation of solution given in equation (4.19) becomes:

$$\underline{y}(t, r) = \frac{13}{8}e^t - \frac{5}{8}e^{-t} \cos 2t - \frac{1}{8}e^{-t} \sin 2t + (r-1)(c_1 e^{at} - c_2 e^{\frac{-b}{2}t} \cosh c_4 t + c_3 e^{\frac{-b}{2}t} \sinh c_4 t),$$

$$\bar{y}(t, r) = \frac{13}{8}e^t - \frac{5}{8}e^{-t} \cos 2t - \frac{1}{8}e^{-t} \sin 2t - (r-1)(c_1 e^{at} - c_2 e^{\frac{-b}{2}t} \cosh c_4 t + c_3 e^{\frac{-b}{2}t} \sinh c_4 t),$$

where

$$c_1 = \frac{30 - 7\sqrt[3]{55 + 3\sqrt{393}} - 7\sqrt[3]{55 - 3\sqrt{393}} + (55 + 3\sqrt{393})^{\frac{2}{3}} + (55 - 3\sqrt{393})^{\frac{2}{3}}}{4[-24 + 3(55 + 3\sqrt{393})^{\frac{2}{3}} + 3(55 - 3\sqrt{393})^{\frac{2}{3}}]},$$

$$c_2 = \frac{54 - 7\sqrt[3]{55 + 3\sqrt{393}} - 7\sqrt[3]{55 - 3\sqrt{393}} - 2(55 + 3\sqrt{393})^{\frac{2}{3}} - 2(55 - 3\sqrt{393})^{\frac{2}{3}}}{4[-24 + 3(55 + 3\sqrt{393})^{\frac{2}{3}} + 3(55 - 3\sqrt{393})^{\frac{2}{3}}]},$$

$$c_3 = \frac{6[-38\sqrt[3]{55 + 3\sqrt{393}} - 38\sqrt[3]{55 - 3\sqrt{393}} - 7(55 + 3\sqrt{393})^{\frac{2}{3}} - 7(55 - 3\sqrt{393})^{\frac{2}{3}}] c_4}{4[-24 + 3(55 + 3\sqrt{393})^{\frac{2}{3}} + 3(55 - 3\sqrt{393})^{\frac{2}{3}}][-12 - (55 + 3\sqrt{393})^{\frac{2}{3}} - (55 - 3\sqrt{393})^{\frac{2}{3}}]},$$

$$c_4 = \frac{1}{6} \sqrt{-36 - 3(55 + 3\sqrt{393})^{\frac{2}{3}} - 3(55 - 3\sqrt{393})^{\frac{2}{3}}}.$$

and  $a, b$  and  $c$  are defined as in (4.20).

**Case 6** Let  $y'(t)$  be (i)-differentiable and  $y(t)$  and  $y''(t)$  be (ii)- differentiable. Then equation (4.2) becomes:

$$s^3 L(y(t)) \Theta s^2 y(0) - s y'(t) - y''(0) = -[-s y(0) \Theta (-s^2) L(y(t)) \Theta y'(t)] - 3[-y(0) \Theta (-s) L(y(t))] + 5L(y(t)),$$

then, we get:

$$l(\underline{y}(t, r)) = \frac{13}{8(s-1)} + \frac{(2r-7)s-7}{8(s^2+2s+5)},$$

$$l(\bar{y}(t, r)) = \frac{13}{8(s-1)} - \frac{(2r+3)s+7}{8(s^2+2s+5)}.$$

Then, we get  $r$  – cut representation of solution as follows:

$$\underline{y}(t, r) = \frac{13}{8}e^t - \frac{7}{8}e^{-t} \cos 2t + r\left(\frac{1}{4}e^{-t} \cos 2t - \frac{1}{8}e^{-t} \sin 2t\right),$$

$$\bar{y}(t, r) = \frac{13}{8}e^t - \frac{3}{8}e^{-t} \cos 2t - \frac{1}{4}e^{-t} \sin 2t - r\left(\frac{1}{4}e^{-t} \cos 2t - \frac{1}{8}e^{-t} \sin 2t\right).$$

**Case 7** Let  $y(t)$  be (i)-differentiable and  $y'(t)$  and  $y''(t)$  be (ii)- differentiable. Then equation (4.2) becomes:

$$s^3 L(y(t)) \Theta s^2 y(0) \Theta s y'(t) - y''(0) = -[-s y(0) \Theta (-s^2) L(y(t)) - y'(t)] - 3[s L(y(t)) \Theta y(0)] + 5L(y(t)),$$

then, we get the system:

$$\begin{aligned} (s^3 + s^2 - 5)l(\underline{y}(t, r)) + 3s l(\bar{y}(t, r)) &= \left(\frac{3}{4} + \frac{r}{4}\right)s^2 + \left(\frac{9}{4} + \frac{3r}{4}\right)s + \frac{19}{2} - \frac{r}{2}, \\ (s^3 + s^2 - 5)l(\bar{y}(t, r)) + 3s l(\underline{y}(t, r)) &= \left(\frac{5}{4} - \frac{r}{4}\right)s^2 + \left(\frac{15}{4} - \frac{3r}{4}\right)s + \frac{17}{2} + \frac{r}{2}. \end{aligned} \tag{4.21}$$

By achieving the same steps given in case 1, we get:

$$\begin{aligned} \underline{y}(t, r) &= l^{-1}\left(\frac{s^2 + 3s + 9}{(s-1)(s^2 + 2s + 5)}\right) + \frac{r-1}{4} l^{-1}\left(\frac{s^2 + 3s - 2}{(s-a)(s^2 + bs + c)}\right), \\ \bar{y}(t, r) &= l^{-1}\left(\frac{s^2 + 3s + 9}{(s-1)(s^2 + 2s + 5)}\right) - \frac{r-1}{4} l^{-1}\left(\frac{s^2 + 3s - 2}{(s-a)(s^2 + bs + c)}\right). \end{aligned} \tag{4.22}$$

where

$$\begin{aligned} a &= \frac{-1}{3} + \frac{1}{3}\sqrt[3]{53 + 3\sqrt{201}} + \frac{1}{3}\sqrt[3]{53 - 3\sqrt{201}}, \\ b &= \frac{2}{3} + \frac{1}{3}\sqrt[3]{53 + 3\sqrt{201}} + \frac{1}{3}\sqrt[3]{53 - 3\sqrt{201}}, \\ c &= -1 + \frac{1}{9}\sqrt[3]{53 + 3\sqrt{201}} + \frac{1}{9}\sqrt[3]{53 - 3\sqrt{201}} + \frac{1}{9}(53 + 3\sqrt{201})^{\frac{2}{3}} + \frac{1}{9}(53 - 3\sqrt{201})^{\frac{2}{3}}. \end{aligned} \tag{4.23}$$

By using relation (4.10), then  $r$  – cut representation of solution given in equation (4.22) becomes:

$$\begin{aligned} \underline{y}(t, r) &= \frac{13}{8}e^t - \frac{5}{8}e^{-t} \cos 2t - \frac{1}{8}e^{-t} \sin 2t + (r-1)(c_1 e^{a t} - c_2 e^{\frac{-b}{2}t} \cosh c_4 t + c_3 e^{\frac{-b}{2}t} \sinh c_4 t), \\ \bar{y}(t, r) &= \frac{13}{8}e^t - \frac{5}{8}e^{-t} \cos 2t - \frac{1}{8}e^{-t} \sin 2t - (r-1)(c_1 e^{a t} - c_2 e^{\frac{-b}{2}t} \cosh c_4 t + c_3 e^{\frac{-b}{2}t} \sinh c_4 t), \end{aligned}$$

where

$$\begin{aligned} c_1 &= \frac{-6 + 7\sqrt[3]{53 + 3\sqrt{201}} + 7\sqrt[3]{53 - 3\sqrt{201}} + (53 + 3\sqrt{201})^{\frac{2}{3}} + (53 - 3\sqrt{201})^{\frac{2}{3}}}{4[30 + 3(53 + 3\sqrt{201})^{\frac{2}{3}} + 3(53 - 3\sqrt{201})^{\frac{2}{3}}]}, \\ c_2 &= \frac{-36 + 7\sqrt[3]{53 + 3\sqrt{201}} + 7\sqrt[3]{53 - 3\sqrt{201}} - 2(53 + 3\sqrt{201})^{\frac{2}{3}} - 2(53 - 3\sqrt{201})^{\frac{2}{3}}}{4[30 + 3(53 + 3\sqrt{201})^{\frac{2}{3}} + 3(53 - 3\sqrt{201})^{\frac{2}{3}}]}, \\ c_3 &= \frac{6[16\sqrt[3]{53 + 3\sqrt{201}} + 16\sqrt[3]{53 - 3\sqrt{201}} + 7(53 + 3\sqrt{201})^{\frac{2}{3}} + 7(53 - 3\sqrt{201})^{\frac{2}{3}}] c_4}{4 [30 + 3(53 + 3\sqrt{201})^{\frac{2}{3}} + 3(53 - 3\sqrt{201})^{\frac{2}{3}}][20 - (53 + 3\sqrt{201})^{\frac{2}{3}} - (53 - 3\sqrt{201})^{\frac{2}{3}}]}, \\ c_4 &= \frac{1}{6} \sqrt{20 - 3(53 + 3\sqrt{201})^{\frac{2}{3}} - 3(53 - 3\sqrt{201})^{\frac{2}{3}}}. \end{aligned}$$



and  $a, b$  and  $c$  are defined as in (4.23).

**Case 8** Let  $y(t)$ ,  $y'(t)$  and  $y''(t)$  be (ii)- differentiable. Then equation (4.2) becomes:

$$-s^2 y(0) \Theta s^3 L(y(t)) \Theta s y'(t) - y''(0) = -[s^2 L(y(t)) \Theta s y(0) - y'(t)] - 3 [-y(0) \Theta (-s) L(y(t))] + 5L(y(t)),$$

then, we get the system:

$$\begin{aligned} (s^3 + s^2)l(\underline{y}(t, r)) + (3s - 5)l(\bar{y}(t, r)) &= \left(\frac{3}{4} + \frac{r}{4}\right)s^2 + \left(\frac{13}{4} - \frac{r}{4}\right)s + 10 - r, \\ (s^3 + s^2)l(\bar{y}(t, r)) + (3s - 5)l(\underline{y}(t, r)) &= \left(\frac{5}{4} - \frac{r}{4}\right)s^2 + \left(\frac{11}{4} + \frac{r}{4}\right)s + 8 + r. \end{aligned} \tag{4.24}$$

By achieving the same steps given in case 1, we get:

$$\begin{aligned} \underline{y}(t, r) &= l^{-1}\left(\frac{s^2 + 3s + 9}{(s-1)(s^2 + 2s + 5)}\right) + \frac{r-1}{4} l^{-1}\left(\frac{s^2 - s - 4}{(s-a)(s^2 + bs + c)}\right), \\ \bar{y}(t, r) &= l^{-1}\left(\frac{s^2 + 3s + 9}{(s-1)(s^2 + 2s + 5)}\right) - \frac{r-1}{4} l^{-1}\left(\frac{s^2 - s - 4}{(s-a)(s^2 + bs + c)}\right). \end{aligned} \tag{4.25}$$

where

$$\begin{aligned} a &= \frac{-1}{3} + \frac{1}{3} \sqrt[3]{-82 + 12\sqrt{159}} + \frac{1}{3} \sqrt[3]{-82 - 12\sqrt{159}}, \\ b &= \frac{2}{3} + \frac{1}{3} \sqrt[3]{-82 + 12\sqrt{159}} + \frac{1}{3} \sqrt[3]{-82 - 12\sqrt{159}}, \\ c &= -1 + \frac{1}{9} \sqrt[3]{-82 + 12\sqrt{159}} + \frac{1}{9} \sqrt[3]{-82 - 12\sqrt{159}} + \frac{1}{9} (-82 + 12\sqrt{159})^{\frac{2}{3}} + \frac{1}{9} (-82 - 12\sqrt{159})^{\frac{2}{3}}. \end{aligned} \tag{4.26}$$

By using relation (4.10), then  $r$  – cut representation of solution given in equation (4.25) becomes:

$$\begin{aligned} \underline{y}(t, r) &= \frac{13}{8} e^t - \frac{5}{8} e^{-t} \cos 2t - \frac{1}{8} e^{-t} \sin 2t + (r-1)(c_1 e^{at} - c_2 e^{\frac{-b}{2}t} \cosh c_4 t + c_3 e^{\frac{-b}{2}t} \sinh c_4 t), \\ \bar{y}(t, r) &= \frac{13}{8} e^t - \frac{5}{8} e^{-t} \cos 2t - \frac{1}{8} e^{-t} \sin 2t - (r-1)(c_1 e^{at} - c_2 e^{\frac{-b}{2}t} \cosh c_4 t + c_3 e^{\frac{-b}{2}t} \sinh c_4 t), \end{aligned}$$

where

$$c_1 = \frac{-12 - 5\sqrt[3]{-82 + 12\sqrt{159}} - 5\sqrt[3]{-82 - 12\sqrt{159}} + (-82 + 12\sqrt{159})^{\frac{2}{3}} + (-82 - 12\sqrt{159})^{\frac{2}{3}}}{4[30 + 3(-82 + 12\sqrt{159})^{\frac{2}{3}} + 3(-82 - 12\sqrt{159})^{\frac{2}{3}}]},$$

$$c_2 = \frac{-42 - 5\sqrt[3]{-82 + 12\sqrt{159}} - 5\sqrt[3]{-82 - 12\sqrt{159}} - 2(-82 + 12\sqrt{159})^{\frac{2}{3}} - 2(-82 - 12\sqrt{159})^{\frac{2}{3}}}{4[30 + 3(-82 + 12\sqrt{159})^{\frac{2}{3}} + 3(-82 - 12\sqrt{159})^{\frac{2}{3}}]},$$
$$c_3 = \frac{6[22\sqrt[3]{-82 + 12\sqrt{159}} + 22\sqrt[3]{-82 - 12\sqrt{159}} - 5(-82 + 12\sqrt{159})^{\frac{2}{3}} - 5(-82 - 12\sqrt{159})^{\frac{2}{3}}] c_4}{4 [30 + 3(-82 + 12\sqrt{159})^{\frac{2}{3}} + 3(-82 - 12\sqrt{159})^{\frac{2}{3}}][20 - (-82 + 12\sqrt{159})^{\frac{2}{3}} - (-82 - 12\sqrt{159})^{\frac{2}{3}}]}$$
$$c_4 = \frac{1}{6} \sqrt{60 - 3(-82 + 12\sqrt{159})^{\frac{2}{3}} - 3(-82 - 12\sqrt{159})^{\frac{2}{3}}}.$$

and  $a, b$  and  $c$  are defined as in (4.26).

## 5. Conclusions

The formula of fuzzy derivatives of any order  $n, n \in \mathbb{Z}^+$  and the formula of fuzzy Laplace transforms of fuzzy derivatives of any order  $n, n \in \mathbb{Z}^+$  are found under generalized H-differentiability. To show the validity of the two above generalizations, solutions to FIVP of the third order are provided.

## References

- [1] Chang SSL, Zadeh L, "On Fuzzy Mapping and Control". IEEE Trans SystCybern, 2: 30-34, 1972.
- [2] Dubios D, Prade H, "To Wards Fuzzy Differential Calculus", Fuzzy Set Syst, 8(1-7):105-116, 225-233, 1982.
- [3] Puri ML, Ralescu DA, "Differentials for Fuzzy Functions", Journal of Mathematical Analysis and Applications, 91:552-558, 1983.
- [4] Abbasbandy S, Allahviranloo T, Darabi P, "Numerical Solution of n-Order Fuzzy Differential Equations ByRunge-Kutta Method", Mathematical and Computational Applications, 16(4):935-946, 2011.
- [5] Allahviranloo T, Ahmady E, Ahmady N, "A Method for Solving nth Order Fuzzy Linear Differential Equations", International Journal of Computer Mathematics, 86(4):730-742, 2009.
- [6] Bede B, Rudas IJ, Bencsik AL, "First Order Linear Fuzzy Differential Equations Under Generalized Differentiability". InfSci, 177:1648-1662, 2006.
- [7] Allahviranloo T, Ahmadi MB, "Fuzzy Laplace Transforms", Soft Comput, 14: 235-243, 2010.
- [8] Salahshour S, Allahviranloo T, "Applications of Fuzzy Laplace transforms", Soft comput, 17: 145-158, 2013.
- [9] Haydar A.K., Mohammad Ali H.F., " Fuzzy Laplace Transforms for Derivatives of Higher Orders", Mathematical Theory and Modeling, 4: 24-31, 2014.