

The Effect of titanium dioxide nanoparticles on the activity of salivary peroxidase in periodontitis patients

Mays A. Talib¹, Basima G. Ali², Eaman A. Al-Rubae³, Maysaa Mahdy⁴

1. Al Anbar Health Directorate, Ministry of Health, Iraq.

2. Department of Periodontics, College of Dentistry, University of Baghdad, Iraq.

3. Department of Basic Science, College of Dentistry, University of Baghdad, Iraq.

4. Al-Sader dental specialized center, Ministry of Health, Iraq.

* Correspondence email; Dr.basimaali@gmail.com

Received date: 20-04-2022

Accepted date: 29-05-2022

Published date: 15-06-2023



Copyright: © 2022 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license

<https://creativecommons.org/licenses/by/4.0/>.



Abstract: Background: The technology of nanoparticles has been expanded to many aspects of modern life. Titanium dioxide nanoparticles were of many nanomaterials utilized in biomedical applications. The interactions between nanoparticles and proteins are believed to be the base for the biological effect of the nanoparticles. The oxidation reaction of many substances is catalyzed by oxidizing enzymes called peroxidases. The activity of salivary peroxidase is elevated with periodontal diseases. The aim of this study is to examine the action of titanium dioxide nanoparticles on salivary peroxidase activity. Material and method: 75 participants were enrolled in this study – Periodontitis group with 44 participants and the non-periodontitis group with 31 participants. The participants' age range was 35 to 50 years for both groups. The clinical parameters of plaque index, gingival index, probing pocket depth and clinical attachment level were used in this study to determine the presence or absence of the periodontal disease. Unstimulated saliva was collected from all participants and analyzed for the activity of peroxidase enzyme under the effect of titanium dioxide nanoparticles. Results: The Periodontitis group showed higher peroxidase enzyme activity than the non-periodontitis group and the activity of salivary peroxidase showed no correlation with the clinical parameters. Titanium dioxide nanoparticles increased salivary peroxidase activity. Conclusion: This study demonstrated that the solid surface of nanoparticles could induce changes in the attached protein molecule which in turn causes changes in the effect of the nanoparticles on living tissue or organism. The titanium dioxide nanoparticles play a role in increasing the activity of salivary peroxidase within the saliva of chronic periodontitis patients.

Keywords: *TiO₂ nanoparticles, salivary peroxidase, periodontitis, saliva.*

Introduction

Periodontitis is a wide-spreading disease characterized by the pain-free and sluggish spread of the condition. This disease starts with a plaque under the gingiva and is altered by the immune response. The disease can occur in different age groups but is most common in adults. It causes loss of the periodontium, which in time leads to tooth mobility and loss^(1,2).

Periodontitis and other inflammatory diseases are related to what is known as oxidative stress. In periodontitis, an imbalance or a shift occurs between antioxidant and oxidant enzymatic and non-enzymatic defense systems. During inflammation, the polymorph nuclear leukocytes make reactive oxygen species (ROS) which are very disastrous to the tissue. The damage caused by these ROS is removed or repaired by antioxidants (AO)⁽³⁾.

Peroxidases are oxidizing enzymes that trigger oxidation reactions for different materials. Hydrogen peroxidase (H₂O₂) helps the reaction to continue then it's reduced to water⁽⁴⁾. Peroxidase can be found in

bodily fluids such as saliva and tears. In addition, it can be found in cells, removing free radicals with the assistance of H₂O₂. The two structures of peroxidases, which are salivary peroxidase and myeloperoxidase, are detected in the whole saliva and play an important role in the defense mechanism⁽⁵⁾.

Nanomaterials consist of components less than 100 nm in at least one dimension⁶. Titanium dioxide nanoparticles (TiO₂ NPs) have many distinctive qualities and properties such as compatibility with living tissue and optical properties⁷ and many studies showed its antibacterial activity^(8,9) when added to dental materials, Therefore TiO₂ NPs have been investigated profusely in recent years to make advances in dentistry. The aim of this study was to analyze the action of TiO₂ NPs on peroxidase enzyme activity in patients suffering from chronic periodontitis.

Subjects and Methods

1. Sample selection:

In this study, two study groups were recruited. The first group (group 1) consisted of 44 participants (periodontitis group), while the second group (group 2) consisted of 31 participants (non-periodontitis group). The total number of participants was 75 and the age range was from 35 to 50 years. The collection of the samples started in October and finished in December of 2017. The participants were patients seeking treatment in the Department of Periodontics at the College of Dentistry, University of Baghdad. Both consent forms and ethical approval were acquired for this study.

2. Clinical examination:

The periodontal parameters (gingival index⁽¹⁰⁾, plaque index⁽¹¹⁾, bleeding on probing, clinical attachment level, and probing pocket depth) were used for the assessment and diagnosis of the periodontal condition of the participants.

3. Collection of Saliva

The spitting method for the collection of whole unstimulated saliva was used. Patients were asked to rinse their mouth first, then wait a few minutes before spitting into a plain tube to collect 5 ml of saliva. This was done at least one hour after the participant's last meal. The time of sample collection was between 9 a.m.- 11 a.m. After the collection of the sample; it was centrifuged for 15 minutes at 2500 rpm. A layer called supernatant is formed and then collected and stored in an Eppendorf tube at -20°C.

4. Laboratory procedures:

a. Saliva sample volume determination:

To determine the best saliva volume to obtain the optimum activity of peroxidase, different volumes of saliva samples were tested (20, 40, 60, 80, and 100 µl).

In this experiment, different volumes of saliva were collected (20, 40, 60, 80 and 100 µl) to determine the best saliva volume to measure the activity of salivary peroxidase. The optimum enzyme activity for this experiment was found to be (100 µl).

b. Characterization of Titanium dioxide nanoparticles:

TiO₂ Nanopowder was obtained from Hongwu international group Ltd, Guangdong, China. An ultraviolet-visible spectrophotometer (PG Instruments Limited/ United Kingdom) was used to determine the absorption spectra for the TiO₂ NPs solution used in this study. The measurement was done at room temperature. The size and structure of the TiO₂ NPs in the samples were determined using a transmission electron microscope TEM (Philips CM10).

c. Salivary Peroxidase Assay:

The colorimetric method was used to determine the activity of the peroxidase enzyme in saliva. In this study, 4-aminoantipyrine was used as a hydrogen donor. The H₂O₂ decomposes through the incubation period causing an elevation in absorption at $\lambda = 510$ nm, which helps in determining the activity of the enzyme, after adding 1.4 ml of 4-aminoantipyrine (2.5 mM) with phenol (0.17 M) solution to 1.5 ml of H₂O₂ (1.7 mM) in a buffer of phosphate (0.2 M) of pH 7.0 solution. The initiation of the reaction was done by adding saliva (100 μ l). To obtain ($\Delta A/\text{min}$) the elevation in absorbance at 510 nm was calculated for 5 minutes. One unit of enzyme activity represents one μ mole decomposition of H₂O₂ through the period of one minute at pH =7.0 under certain conditions.

d. Preparation of TiO₂ NPs solution:

The first step was to prepare a stock solution of TiO₂ NPs (300 $\mu\text{g}/\text{ml}$). A solvent of 3:1 water to ethanol was used to dilute the stock solution to different concentrations (20, 40, 60, 80, 100, and 120 $\mu\text{g}/\text{ml}$). The best concentration was found to be 120 $\mu\text{g}/\text{ml}$.

e. Determination of the action of TiO₂ NPs on the activity of peroxidase enzyme:

A detection kit was used to measure the action of TiO₂ NPs on the activity of peroxidase enzyme by adding 20 μ l of 120 $\mu\text{g}/\text{ml}$ TiO₂ NPs to 100 μ l of saliva and then using the detection kit. To measure the absorbance at a wavelength of 510 nm and establish the peroxidase activity in the sample. A percentage of activation equation was used to calculate the effect percentage. The activity of the enzyme with the presence of TiO₂ NPs and without the nanoparticles was compared according to the percentage of activation equation to calculate the percentage of effect on the peroxidase enzyme activity:

$$\% \text{ activation} = 100 - 100 \times [\text{Activity in the presence of nanoparticles} / \text{Activity without the nanoparticles}].$$

5. Statistical analysis:

Statistical package for social sciences (SPSS) and Microsoft Office Excel were used to analyze the collected data. The significant difference was assessed using the student t-test ($P < 0.05$). The Pearson correlation coefficient was also used.

Results

1. Clinical findings

The mean values and standard deviations (SD) of Peroxidase activity for group 1 were highly significant compared to group 2 (p< 0.001).

The mean values and SD of Peroxidase activity for both groups are shown in table1:

Table 1: mean, standard division, t-test and p-value of Peroxidase activity for group1 and group2:

| Groups | N | Mean ±SD | t-test | p-value |
|--------|----|---------------|--------|--------------------|
| Group1 | 44 | 242.38±137.31 | 2.68 | 0.001 ^s |
| Group2 | 31 | 175.67±76.08 | | |

N: Number, SD: Standard Deviation, t-test: student's t-test, p-value: Probability value, S: significant p-value (p< 0.001)

The correlation was non-significant when comparing the activity of peroxidase and both plaque index (PLI) and gingival index (GI) in group 1. A weak negative and non-significant correlation was determined for GI and a non-significant correlation for PLI in group 2. Results are as illustrated in the table below:

Table 2: correlation and p-value between periodontal parameters and peroxidase activity in group1 and group 2:

| Periodontal parameters | Study groups | | | |
|------------------------|--------------|---------------------|---------------|---------------------|
| | Group1, N=44 | | Group 2, N=31 | |
| | r | p_value | r | p_value |
| PLI | 0.044 | 0.777 _{NS} | 0.360 | 0.05 _{NS} |
| GI | 0.261 | 0.086 _{NS} | -0.011 | 0.951 _{NS} |

r: Coefficient of correlation, p_value: Probability value, PL: Plaque index, GI: Gingival index, NS: Non - Significant p_value ≥ 0.05, N: Number.

2. TiO2 NPS characterization:

Spectra of UV-VIS absorption were used to determine TiO2 NPs absorption qualities. The peak was around 200 nm which points to the intensity of absorption of TiO2 NPs dispersion at <300 nm in the ultraviolet area of the spectrum, as shown in Figure (1):




Figure 1: Absorbance spectra of TiO₂ nanoparticles.

TEM was used to identify the structure and Nano size measurement of TiO₂ NPs in the samples (figure 2). The average diameter of the particle size was found to be < 30nm.




Figure 2: The TEM pictures and size distribution of TiO₂ NPs. The arrow shows a nanoparticle size of 28.5nm (less than 30 nm).

3. The effect of TiO₂ NPs on Peroxidase enzyme in saliva

The differences in the activities of peroxidase enzyme (mean ± SD) for the studied groups are shown in Table (3) below. The results conveyed that the activity of peroxidase in saliva samples from group 1 with TiO₂ NPs was greater than the enzyme activity in samples without TiO₂ NPs with a statistically high significance difference (p-value< 0.001), as shown in the table below. Similarly, a highly significant difference was also found After comparing the enzyme activity in group 2 with and without TiO₂ NPs.

Table 3: intragroup comparison of the effect of TiO2 NPs on peroxidase activity in both groups.

| | Study groups | | | | | |
|-------------------------|----------------|--------|----------------------|---------------|--------|----------------------|
| | Group1, N=44 | | | Group1, N=31 | | |
| | MEAN± SD | t-test | P_value | MEAN± SD | t-test | P_value |
| with TiO2 NPs | 304.81± 153.43 | 7.230 | 0.0001 ^{HS} | 213.48± 88.65 | 5.036 | 0.0001 ^{HS} |
| without TiO2 NPs | 242.38± 137.31 | | | 175.67± 76.08 | | |

TiO2 NPs: Titanium Dioxide Nanoparticles, HS: Highly Significant (p_value< 0.001)

The activity of peroxidase enzyme in the presence of TiO2 NPs was compared between both groups. The results showed a high statistically significant difference (p-value< 0.001), as stated in Table (4) below:

Table 4: Intergroup comparison of the effect of TiO2 NPs on peroxidase activity in both groups and significance level:

| Groups | peroxidase Activity in U/L Mean ±SD | t-test | P_value |
|-------------------------------|-------------------------------------|--------|---------------------|
| Group1(With TiO2 NPs) | 304.81± 153.43 | 3.252 | 0.001 ^{HS} |
| Group2 (With TiO2 NPs) | 213.48± 88.65 | | |

Figure (3) shows the action of TiO2 NPs (µg/ml) on the peroxidase activity (U/L) in the reaction mix of a total volume of (3020). It was established that TiO2 NPs at a concentration of 0.79 µg/ml have a higher activation effect on the enzyme activity in the mixture.

At a concentration of 0.79 µg/ml of TiO2 NPs, peroxidase activity reached its highest activation percentage (65.93%), making this concentration of the nanoparticles the most effective concentration in this experiment as shown in Figure (4).




Figure 3: effect of different concentrations of TiO2 NPs on salivary peroxidase activity.

V= volume, U/L= units per liter, µg/ml= microgram per millilitre, TiO2 NPs= titanium dioxide nanoparticles




Figure 4: Percentage of activation of salivary peroxidase activity in different concentrations of TiO₂ NPs.
TiO₂ NPs= titanium dioxide nanoparticles

Discussion

Nowadays, TiO₂ NPs were incorporated in many products used in daily life which made it necessary to evaluate the action of this type of nanoparticle on living systems⁽¹²⁾.

The chemical evaluation revealed that TiO₂ NPs caused an elevation in peroxidase activity. Al-Rubaei et al. in 2016 showed that Total salivary peroxidase was activated significantly ($p < 0.001$) by TiO₂ NPs¹³. The surface of the nanoparticles can cause changes in the form of the protein when it attaches itself to the nanoparticle surface. This change in the protein form can lead to a change in the protein's function. This changes the bio reactivity of the nanoparticles⁽¹⁴⁾.

Many studies showed evidence of conformational changes when enzyme interacts with NPs, such as a study on Zinc oxide NPs in which it was found that the NPs modify the secondary structure of lysozyme. The enzyme keeps its catalytic activity and resists denaturation in the presence of these NPs⁽¹⁵⁾.

In another study to determine the effect of zinc oxide nanoparticles on peroxidase enzyme activity, an inhibition action on peroxidase enzyme activity was noticed for this type of nanoparticles⁽¹⁶⁾. The activity of peroxidase was found to be elevated in periodontitis compared to non-periodontitis patients, with a significant difference. Similar results were obtained in another study by Al-Rassam et al. in 2017⁽¹⁷⁾ in which a comparison was made between salivary peroxidase activity in the chronic periodontitis group and a control group, the enzyme activity was elevated in the chronic periodontitis group compared to the control group and with a significant difference. Another study found that salivary peroxidase activity significantly increases with inflammation and reduces after oral hygiene measures⁽¹⁸⁾.

In another study, Glutathione peroxidase in saliva and gingival tissues of subjects with and without chronic periodontitis was evaluated and it was found that there was an increase in Glutathione peroxidase level in saliva and inflamed gingival tissue⁽¹⁹⁾.

The elevated peroxidase level in the saliva and tissue of the gingiva of patients suffering from periodontitis can be accredited to the scavenging of the redundant lipid peroxidation products at the inflammatory sites⁽²⁰⁾.

Also, the elevated level of reactive oxygen species (ROS) formed may have caused oxidative stress, which lead to an increased need for peroxidase generation to establish the ROS–AO balance to prevent tissue damage⁽¹⁹⁾.

In this study, a non-significant correlation between the gingival index and the activity of salivary peroxidase in group1 and group2, a similarly non-significant correlation was found between plaque index and the action of salivary peroxidase in both groups ($P > 0.05$). This is, in contrast, to a study by Dagar et al. in 2015 ⁽²¹⁾ in which a significant correlation was found between the activity of peroxidase and PLI. This difference in results could be accredited to the variation in saliva-collecting methods, the number of participants in each study, and analysis methods.

In this study, TiO₂ NPs caused an increase in the action of the peroxides enzyme. Peroxidase enzyme level in the saliva of patients suffering from periodontitis was significantly higher than patients without periodontitis disease.

However, more research are needed to overcome the limitations of this study. Some of these limitations, such as the limited study sample size, the time constraints, and the technique-sensitive method for preparing the nanoparticle solution, can be addressed in future studies for more precise results.

Conclusion

The solid surface of nanoparticles can induce changes in the attached protein molecule which in turn causes changes in the effect of the nanoparticles on living tissue. The titanium dioxide nanoparticles play a role in increasing the activity of salivary peroxidase within the saliva of periodontitis patients.

Conflict of interest: None

References

1. Shaddox M, Walker CB. Treating chronic periodontitis: current status, challenges, and future directions. *Clin Cosmet Investig Dent.* 2010;11:79-91. [\(Crossref\)](#)
2. Trivedi S, Lal N. Antioxidant enzymes in periodontitis. *J Oral Biol Craniofac Res.* 2017;7:54-57. [\(Crossref\)](#)
3. Vincent RR, Appukkuttan D, Prakash PSG, Victor J, Vincent RR. Oxidative Stress: Role in Patho-genesis of Periodontal Disease. *Int J Pharma Bio Sci.* 2017;8:1033-1041. [\(Crossref\)](#)
4. Totan A, Greabu M, Totan C, Spinu T. Salivary aspartate aminotransferase, alanine aminotransferase and alkaline phosphatase: possible markers in periodontal diseases?. *Clinical Chemistry and Laboratory Medi-cine (CCLM).* 2006 May 1;44(5):612-5. [\(Cross-ref\)](#)
5. Sakamoto W, Fujii Y, Kanehira T, Asano K, Izumi H. A novel assay system for myeloperoxidase activ-ity in whole saliva. *Clin Biochem.* 2008;41:584-590. [\(Crossref\)](#)
6. Verma SK, Prabhat KC, Goyal L, Rani M, Jain A. A critical review of the implication of nano-technology in modern dental practice. *Natl J Maxillofac Surg.* 2010;1:41-44. [\(Crossref\)](#)

7. Yin ZF, Wu L, Yang HG, Su YH. Recent progress in biomedical applications of titanium dioxide. *Phys Chem Chem Phys*. 2013;15:4844-4858. ([Crossref](#))
8. Hailan SY, Al-khatieeb M. Antimicrobial efficacy of silver, zinc oxide, and titanium dioxide na-noparticles incorporated in orthodontic bonding agent. *J Bagh Coll Dent*. 2019;31:10-16. ([Crossref](#))
9. Sodagar A, Akhouni MSA, Bahador A, Jalali YF, Behzadi Z, Elhaminejad F, et al. Effect of TiO₂ nanoparticles incorporation on antibacterial properties and shear bond strength of dental composite used in Orthodontics. *Dental Press J Orthod*. 2017;22:67-74. ([Crossref](#))
10. Loe H. The Gingival Index, the Plaque Index and the Retention Index Systems. *J Periodontol*. 1967;38:610-616.. ([crossref](#))
11. Silness J, Loe H. Periodontal disease in pregnancy II. Correlation between oral hygiene and periodontal condition. *Acta Odontol Scand*. 1964;22:121-35. ([Crossref](#))
12. Parashar A, Chakraborty D, Alex SA, Dan P, Chandrasekaran N, Mukherjee A. Effects of titanium dioxide nanoparticles on horseradish peroxidase-mediated peroxidation reactions. *J Mol Liq*. 2017;241:852-60. ([Crossref](#))
13. Al-Rubae EAS, Salman ZA, Abd ST, Aziz RA. Kinetic study of the effect of titanium dioxide nanoparticles on salivary peroxidase activity. *Int J Biol Res*. 2016;1:56-9.
14. Saptarshi SR, Duschl A, Lopata AL. Interaction of nanoparticles with proteins: relation to bio-reactivity of the nanoparticle. *J Nanobiotechnology*. 2013;11:26. ([Crossref](#))
15. Chakraborti S, Chatterjee T, Joshi P, Poddar A, Bhattacharyya B, Singh SP, et al. Structure and activity of lysozyme on binding to ZnO nanoparticles. *Langmuir*. 2010;26:3506-13. ([Crossref](#))
16. Abd ST, Ali AF. Effect of zinc oxide nanoparticles on total salivary peroxidase activity of human saliva (in vitro study). *J Bagh Coll Dent*. 2015;27:178.
17. Al-Rassam ZT, Taha MY, Taha MYM. Serum cytokines profiles and some salivary parameters in chronic periodontitis patients in Mosul, Iraq. *Int J Sci*. 2014;16:339-50.
18. Reddahi S, Bouziane A, Rida S, Tligui H, Ennibi O. Salivary biomarkers in periodontitis patients: A pilot study. *International Journal of Dentistry*. 2022 Mar 24;2022. ([Crossref](#))
19. Smitha S, Biju T, Rahul B, Nina S. Comparative evaluation of salivary and gingival tissue glutathione peroxidase levels in subjects with healthy periodontium and chronic periodontitis. *Sch J Dent Sc*. 2017;4:211-5. ([Crossref](#))
20. Guentsch A, Preshaw PM, Bremer-Streck S, Klinger G, Glockmann E, Sigusch BW. Lipid peroxidation and antioxidant activity in saliva of periodontitis patients: effect of smoking and periodontal treatment. *Clinical oral investigations*. 2008 Dec;12:345-52.. ([Crossref](#))
21. Dagar M, Deepa DK, Molly M, Sharma A, Khattak BP. Effect of nonsurgical periodontal therapy on salivary myeloperoxidase levels: a biochemical study. *J Indian Soc Periodontol*. 2015;19:531-6. ([Crossref](#))

**تأثير جزيئات ثاني اوكسيد التيتانيوم النانوية على فعالية انزيم البيروكسيديس في لعاب المرضى المصابين بالتهاب دواعم السن
الباحثون: طبيب الاسنان الاختصاص ميس علاء ,أستاذ باسمة غفوري ,أستاذ دكتور ايمان الربيعي وطبيب الاسنان الاختصاص ميساء مهدي
الخلاصة**

الخلفية: لقد تم توسيع تقنية الجسيمات النانوية لتشمل العديد من جوانب الحياة الحديثة. كانت الجسيمات النانوية لثاني أكسيد التيتانيوم واحدة من العديد من المواد النانوية التي تم استخدامها في التطبيقات الطبية الحيوية. يعتقد أن التفاعلات بين الجسيمات النانوية والبروتينات هي أساس التأثير البيولوجي للجسيمات النانوية. يتم تحفيز تفاعل الأكسدة للعديد من المواد عن طريق إنزيمات مؤكسدة تسمى البيروكسيداز. يرتفع نشاط البيروكسيداز اللعابي في أمراض اللثة لدراسة تأثير الجسيمات النانوية لثاني أكسيد التيتانيوم على نشاط بيروكسيداز اللعاب ، تم تسجيل 75 مشاركاً في هذه الدراسة. مجموعة التهاب دواعم السن 44 مشاركاً ومجموعة غير التهاب دواعم السن مع 31 مشاركاً. تراوحت أعمار المشاركين من 35 إلى 50 عامًا لكلا المجموعتين. تم استخدام المعلمات السريرية لمؤشر اللويحة ، مؤشر اللثة ، فحص عمق الجيب ومستوى الارتباط السريري في هذه الدراسة لتحديد وجود أو عدم وجود مرض اللثة. تم جمع اللعاب غير المحفز من جميع المشاركين وتحليل نشاط إنزيم البيروكسيداز تحت تأثير الجسيمات النانوية لثاني أكسيد التيتانيوم. أظهرت مجموعة التهاب دواعم السن زيادة نشاط إنزيم البيروكسيداز مقارنة بمجموعة غير التهاب دواعم السن ، ولم يظهر نشاط البيروكسيداز اللعابي أي ارتباط بالمعيار السريري. تم زيادة نشاط بيروكسيداز اللعاب بواسطة جزيئات ثاني أكسيد التيتانيوم النانوية. ظهرت هذه الدراسة انه بالامكان أن يتسبب السطح الصلب للجسيمات النانوية في إحداث تغييرات في جزيء البروتين المرتبط والذي يؤدي بدوره إلى تغييرات في تأثير الجسيمات النانوية على الأنسجة الحية أو الكائن الحي. يلعب **TiO2 NPs** دوراً في زيادة نشاط بيروكسيداز اللعاب داخل لعاب مرضى التهاب دواعم السن المزمن.