



Preparation and Characterization of 1, 3-Bis (methylsalicylate5-yl diazo) Benzene and Its Complexes with Some of First Period Transition Metals

Khaula M. Sultan

Dept. of Chemistry/ College of Education for Pure Science (Ibn- Al Haitham)/
University of Baghdad

Received in:22/March/2015 , Accepted in:30/June/2015

Abstract

In this study azo dye was prepared by the reaction of m-phenylenediazonium chloride with methyl salicylate, the resultant compound was used as a ligand for complex formation with Fe^{+2} , Cu^{+2} , Zn^{+2} , Ni^{+2} and Co^{+2} ions. The prepared ligand was characterized by ^1H NMR, UV-Vis., And FTIR spectroscopy, CHN analysis, in addition the complexes were characterized by TGA, UV-Vis., FTIR and conductivity methods. The results indicate that the ligand chelated through phenoxy and carboxyl groups as a O4 quadra dentate ligand, the Co complex complet its hexagon coordination by bonding with chlorine and the complex would be electrolytic in opposite with rest complexes.

Key words: Azo compounds, 3d metal complexes, transition metal complexes, Azo ligands

Introduction

Organic reagents have wide spread and miscellaneous applications due to its stability and high speed of reaction with different metal ions [1], in addition many of them have a high selectivity and sensitivity which draw a considerable attention. Azo compounds are one of the mostly important types of organic reagents because azo bond can act as coordination center usually [2,3] but not always[4], at the same time azo group is considered as a strong chromophoric center in these compounds [5]. This character makes the azo compounds play a major rule in qualitative and quantitative chemical analysis of metal ions [6], in addition to their use as indicators, in industrial field azo compound used from long time as drugs and dyes, at present they have a wide application in printing, electronic photography, color formers, liquid crystal display, laser technology, data storage, optical actuators, photorefractive media, fiber dye, food and cosmetics industry [7]. This study aims to prepare a ligand from Methylsalicylate with m-phenelendiamine by diazo coupling and then attach the prepared ligand with ionic metals Co and Ni.

Experimental

Materials


All chemicals used in this study were analytical grade and used without further purification. $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$, $\text{NiCl}_2 \cdot 6\text{H}_2\text{O}$, $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, ZnSO_4 and $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ m-phenelyndiamine hydrochloride, NaNO_2 , HCl , Methylsalicylate, Methanol, Ethanol, DMSO, H_2SO_4 and KOH

Instruments

C.H.N (Shimadzu) elemental analyser, ^1H NMR (Bruker 400MHz the solvent deuterized DMSO). FTIR spectra with KBr disc in the range (400-4000) cm^{-1} (Shimadzu FTIR-8400S). Electronic spectra were recorded on (Shimadzu UV-Vis. 160A) spectrophotometer. Conductivity were measured for 0.0001M of complexes in ethanol at 25°C using (Philips PW-Digital Conductometer). Thermal analysis was done by DSC-TGA (Linseis).

Synthesis of the Ligand

The Ligand was synthesized by dissolving (2 g) of m-phenylenediamine hydrochloride in a mixture of (2 ml) H_2SO_4 , 10 ml Ethanol, 10 ml of distilled water, the mixture was cooled to 5°C . 10ml of cooled 10% aqueous solution of NaNO_2 was added dropwise with stirring in order to obtain the diazonium salt solution. After 30 minute the diazonium solution was slowly added to a cooled solution of a mixture of (3 g) methylsalicylate and (1.2 g) of KOH to obtain the Ligand, the dark colored mixture containing the solid precipitate was filtered off, washed then re-crystallized with 70% ethanol. The reaction scheme is:



Preparation of metal complexes

The complexes was prepared by the addition of (10 ml)aqueous solution of the metal salt its concentration 1 M to 0.01 mmole of the Ligand dissolved in 20 ml 50% methanolic aqueous solution. The darck precipitate was settled over night then recrystallized from 70% ethanol.

Results and Discussion

^1H NMR


Figure(18) indicates ^1H NMR spectra of the Ligand contain a singlet band at 2.4 ppm belong to the methyl protons, a multiplet band between (6-9ppm) belongs to the Aromatic protons, finally a singlet band at (10.512 ppm) belongs to the phenolic protons.

Electric conductivity

The measured electric conductivity for the complexes in DMSO solvent, the results indicates that the Co complex is charged complex while the other complexes were neutral complexes.

Thermal analysis

The TGA thermograms of, figures(1,2,3,4 and 5) Co, Fe, Zn and Cu complexes showing two conversion regions while in Ni complex they were three . The first one is represented by losing of hydration water molecules in all complexes [8]. The second conversion region is the most characteristic and it represents the decomposition point of the complex, at this stage all the complexes lose their organic part leaving the metal (II) oxide, in Co complex case this its producing oxide was combined with KCl while in Ni complex this stage produces NiCO_3 which is further decomposed at the third conversion to NiO [9,10].



FTIR spectra:

Figure (6) shows the FTIR spectrum for the ligand that display a broad band at 3381 cm^{-1} belongs to the stretching vibration of phenolic OH group. This band is merged with weak aromatic and aliphatic C-H stretching bands at 3067 and 2953 cm^{-1} respectively. A strong band at 1674 cm^{-1} is due to stretching vibration of C=O bond. The low value that took place as a result of the resonance with aromatic ring which decreases its bond order and hence its force constant [11]. In addition, the stretching vibration of N=N bond appears at 1612 cm^{-1} . The C-O vibration frequency displays two values, the first one at 1215 cm^{-1} belongs to phenolic group, and the second one is part of carboxyl group. Finally, there is a very important absorption band at 847 cm^{-1} belongs to out of the plane bending vibration of C-H aromatic group which is characteristic to the para substituted aromatic compounds [12]. The existence of this group is a good indicator to explain that: the azo group takes the para position relative to phenolic OH group and does not take the ortho position. Figures (7, 8, 9, 10 and 11) show the FTIR spectra of the complexes. In complexes there is an existence of weak broad band above 3300 cm^{-1} assigned to stretching vibration of OH groups in hydration water molecules [13]. There is a great alteration in the C=O absorption band in all complexes. This suggests that the carbonyl group plays a major role in the coordination with the metal ion. The stretching vibration of azo group is red shifted to lower frequency. This happens due to the change of the resonance state in the aromatic ring that which caused by the alteration in its electronic density through the formation of the complex. The C-O bands is also affected by

coordination we believe the later band that belongs to phenolic C-O, bond has lost its H atom and coordinated with metal ion to form the complex. As a result of this coordination, we observed the appearance of a new band below 520 cm^{-1} assigned to M-O bond.

Electronic spectra

The electronic spectra of compounds in DMSO solvent are shown in Figures (12, 13, 14, 15, 16 and 17). The ligand shows strong absorption bands at 310, 364 and 457nm with molar absorbance 5×10^3 , 5.6×10^3 , and $4.8 \times 10^3\text{ mole.cm}^{-2}$ respectively. The UV-vis. Spectrum indicates that the 364nm band belongs to $\pi \rightarrow \pi^*$ transition of $-\text{N}=\text{N}-$ azo chromophoric group[14]. While the 310nm band belongs to $\pi \rightarrow \pi^*$ transition of phenolic group[5], finally the shoulder band at 457nm is due to $n \rightarrow \pi^*$ transition of the same group[5]. The excited state in the $\pi \rightarrow \pi^*$ transition of azo group plays a major role in the conformational transformations in the ligand, because it acts to ease the rotation N=N bond. This process is observed in the cis-trans transformations of the azo compounds. In the electronic spectra of the complexes, we have observed that the $\pi \rightarrow \pi^*$ transition of azo group is not greatly affected by coordination which supported the assumption that the azo group did not act as a coordination center with metals. On the other hand there is a considerable alteration of the bands that belong to phenolic group which considered as evidence that the phenolic group is one of the coordination centers that chelated with metal ion. In addition we observed the increase of the intensity of band of azo group, this may relate to the fact that the quantum probabilistic transition of this bond is stabilized after complex formation. Indeed the formation of complex through the contribution of the phenoloic and the carbonyl groups upon complexation through the chelating with metal ions require fold of the two salicylic parts that close to each other in the ligand which are accomplished by twisting of the $-\text{N}=\text{N}-$ bond. The transition state of this bond may help to satisfy this condition. The above discussion is proceeding for all complexes as well. Finally, the high values of the molar absorbance of the ligand chromophores act to cover the d-d transitions in the metal ions of the complexes [15].

References

- 1- Lodeiro, C. and Pina, F., (2009), Luminescent and chromogenic molecular probes based on polyamines and related compounds, *Coordination Chemistry Reviews*, 253, 1353–1383.
- 2- Anitha ,K. R.; Venugopala Reddy and Fasiulla,(2015), Studies on metal (II) complexes of bisazo dye 2, 21 [benzene-1,3-diyl di (E) diazene 2,1-diyl] dianiline, *Arch. Appl. Sci. Res.*, 7 (1),59-65.
- 3- Anitha, K. R.; Venugopala Reddy and Rao, K. S.Vittala, (2011), Synthesis and antimicrobial evaluation of metal (II) complexes of a novel bisazo dye 2, 21 [benzene-1, 3-diyl di (E) diazene 2, 1-diyl] bis (4-chloroaniline), *J. Chem. Pharm. Res.*, 3(3), 511-519.
- 4- Raj, H. D. and Patel Y. S.,(2015), Synthesis, characterization and antifungal activity of metal complexes of 8-hydroxyquinoline based azo dye, *Advances in Applied Science Research*, 6(2), 119-123.
- 5- Abd-Alredha, L; AL-Rubaie, R. and Mhessn, R. J. (2012) Synthesis and Characterization of Azo Dye Para Red and New Derivatives, *E-Journal of Chemistry*, 9(1), 465-470.

- 6- Patel, K. S.; Patel, J. C.; Dholariya, H. R.; Patel, V. K.; and Patel, K. D.; (2012), Synthesis of Cu(II), Ni(II), Co(II), and Mn(II) Complexes with Ciprofloxacin and Their Evaluation of Antimicrobial, Antioxidant and Anti-Tubercular Activity, *Open Journal of Metal*, 2, 49-59.
- 7- Sidir, Y. G.; Sidir, I.; Berber, H. and Tasal, E. (2011), An experimental study on relationship between hammett substituent constant and electronic absorption wavelength of some azo dyes, *Journal of Science and Technology*, 1, 7-11.
- 8- Stamatatos, T. C.; Katsoulakou, E.; Nastopoulos, V.; Raptopoulou, C. P.; Manessi-Zoupa, E. and Perlepes S. P., (2003), Cadmium Carboxylate Chemistry: Preparation, Crystal Structure, and Thermal and Spectroscopic Characterization of the One-dimensional Polymer $[Cd(O_2CMe)(O_2CPh)(H_2O)_2]_n$, *Z. Naturforsch.* 58b, 1045 – 1054.
- 9- Petrofsky, J. L.,(1964), Complexes OF Iron(II) Cobalt(II)Nickel(II) and Copper(II) With 2-a Pyridylmethyleneaminomethyl)Pyridine, Doctora Thesis, University Of Florida, USA.
- 10- Isaacs T. (1963), the Mineralogy and chemistry of the Nickelcarbonates, Doctora Thesis, Sheffield University, UK.
- 11- Gans, P. (1975), Vibrating molecules, 1st. edition, Chapman & Hall, London.
- 12- Silverstein, R. Webster, F. and Kiemle, D.(2005), Spectrometric Identification of Organic Compounds, 7th. Edition, John Wily & Sons, Inc., USA.
- 13- Nakamoto, K. (1982), Infra-Red Spectra of Inorganic and Coordination Compounds, 2ed. Edition,Mc-Grow Hill, Japan.
- 14- Mohammed, I. A. and Mustapha A. (2010), Synthesis of New Azo Compounds Based on N-(4- Hydroxyphenyl)maleimide and N-(4-Methylphenyl)maleimide, *Molecules*, 15, 7498-7508.
- 15- Lappert ,M. L. (1968), *Inorganic Electronic Spectroscopy*, 1st. edition, Elsevier, Netherlands.

Table (1): FTIR band values

Compound	νH_2O cm^{-1}	$\nu N = N$ cm^{-1}	$\nu C = O$ cm^{-1}	$\nu C - O$ cm^{-1}	$\nu M - O$ cm^{-1}	νOH phenolic cm^{-1}
L	-----	1612	1674	1213	-----	3381
Ni-comp.	3345	1595	1668	1231	484	-----
Co-comp.	3354	1601	1630	1234	446	-----
Cu-comp.	3300	1599	1682	1200	513	-----
Fe-comp.	3300	1599	1682	1209	490	-----
Zn-comp.	3381	1606	1666	1192	513	-----

Table (2): Elemental analysis of the Ligand

Element %	C	H	N
Practical	59.855	4.234	17.603
Theoretical	60.830	4.148	17.603

Table (3): UV-Vis. Band values

Compound	λ azo $\pi - \pi^*$ (nm)	$\epsilon \cdot 10^{-3}$ L.mol ⁻¹ .cm ⁻¹	λ phenolic $\pi - \pi^*$ (nm)	$\epsilon \cdot 10^{-3}$ L.mol ⁻¹ .cm ⁻¹	λ phenolic $n - \pi^*$ (nm)	$\epsilon \cdot 10^{-3}$ L.mol ⁻¹ .cm ⁻¹
L	364	5.6	310	5	457	4.8
Ni-comp.	364	6.1	-----	-----	461	5
Co-comp.	366	5.3	-----	-----	466	4.9
Cu-comp.	365	4.5	-----	-----	465	4.2
Fe-comp.	370	5.1	-----	-----	465	4.9
Zn-comp.	370	5	-----	-----	469	4.9

Table (4) :Physical properties of the Ligand and its complexes

Compound	M.wt/ g.mol ⁻¹	Colour	Yeild%	m.p. ^o C (dec.)	Λ_m / ohm ⁻¹ .cm ² .mol ⁻¹
Ligand	434	Reddish brown	62	235	-----
NiL.6H ₂ O	598.7	Dark brown	86	337.7	1.207
K ₂ [CoLCl ₂]. H ₂ O	599	Dark brown	88	224.8	32.1
FeL. H ₂ O	505.85	Dark brown	85	360.9	1.33
CuL. H ₂ O	513.5	Dark brown	84	321.7	1.32
ZnL. H ₂ O	515.4	Dark brown	88	423.6	1.02




Figure (1): Thermogram of the Ni complex.




Figure (2): Thermogram of the Co complex.




Figure (3): Thermogram of the Cu complex.




Figure (4): Thermogram of the Fe complex.




Figure (5): Thermogram of the Zn complex.




Figure (6): FTIR spectrum of the ligand.




Figure (7): FTIR spectrum of the Ni complex .




Figure (8): FTIR spectrum of the Co complex




Figure (9): FTIR spectrum of the Cu complex




Figure (10): FTIR spectrum of the Fe complex




Figure (11):FTIR spectrum of the Fe complex.




Figure (12): electronic spectrum of the ligand.




Figure (13): electronic spectrum of the Ni complex




Figure (14): UV-Vis. of the Co complex.




Figure (15): UV-Vis. of the Cu complex.




Figure (16) :UV-Vis. of the Fe complex.




Figure (17): UV-Vis. of the Zn complex.




Figure (18): ¹H NMR spectrum of the Ligand.




Figure (19): The general structure of Ni, Cu, Fe and Zn complexes.




Figure (20) :The general structure of Co complex.

تحضير و تشخيص 1,3-Bis (methylsalicylate5-yl diazo)benzene

مع معقداته لبعض عناصر الدورة الانتقالية الاولى

خولة محمد سلطان

قسم الكيمياء/ كلية التربية للعلوم الصرفة (ابن الهيثم)/ جامعة بغداد

استلم في: 22/اذار/2015 ، قبل في: 30/حزيران/2015

الخلاصة

تم تحضير صبغة ازو من تفاعل m-phenyldiazonium chloride مع methyl salicylate. في هذه الدراسة استعمل المركب المحضر كليكند لتكوين معقدات تناسقية مع ايونات Fe^{+2} , Cu^{+2} , Zn^{+2} , Ni^{+2} , Co^{+2} اذ تم تشخيص الليكند المحضر مع معقداته بمطيافيتي 1H NMR و FTIR و UV-vis. وتحليل العناصر CHN فضلا عن ان المعقدات المحضرة شخضت ايضا بقياس TGA و FTIR و UV-vis. و التوصيلية اذ بينت النتائج ان الليكند يرتبط مع الايونات من خلال مجاميع الفينوكسي و الكربونيل و بذلك يكون ليكند رباعي السن من نوع O4. كما انه في معقد الكوبلت فانه يتم اكمال التناسق السداسي من خلال الارتباط بالكلور وبذلك يكون هذا المعقد الكتروليتي بخلاف المعقدات الاخرى.

الكلمات المفتاحية: مركبات الازو, معقدات عناصر 3d, معقدات العناصر الانتقالية, ليكندات الازو.