

Synthesis, Characterization, and Antibacterial Evaluation of New N-Phenyl-naphthalimides linked to Benzothiazole moiety

A. S. Hamed

College of Veterinary Medicine/ University of Fallujah

Abstract

In this work several new naphthalimides linked to substituted benzothiazole moiety were prepared. Preparation of the new naphthalimides was performed via multistep synthesis which include reacted of naphthalic anhydride with 4-aminobenzoic acid, then the product subsequently introduced in reaction with thionyl chloride producing 4-(N-naphthalimidyl) benzoyl chloride which in turn introduced in reaction with substituted benzothiazoles affording the target new naphthalimides. The prepared compound were screened for their antimicrobial activity against gram positive (*Staphylococcus aureus*, *Streptococcus pyogenes*), gram negative (*Escherichia coli* and *Pseudomonas aureginosa*) and yeast (*Candida albicans*). The results indicated that they exhibit good antimicrobial activity against the tested organism.

تحضير وتشخيص وتقدير الفعالية البايولوجية لـ N- فنيل نفتال ايميدات جيدة مرتبطة بمكونة

البنزو ثايازول

احمد سليمان حمد

كلية الطب البيطري / جامعة الفلوجة

الخلاصة

تم في هذا البحث تحضير عدد من النفثال ايميدات الجديدة المرتبطة بمكونة البنزو ثايازول المعوضة. تم تحضير النفثال ايميدات الجديدة بطريقة التحضير متعدد الخطوات، تم في الخطوة الأولى مفاعلة انهيديد النفثالك مع 4- امينو حامض البنزويك ومن ثم تم مفاعلة الناتج مع كلوريد الثايونيل لتكوين 4- (N- نفتال ايميديل) كلوريد البنزويل والذي يدخل في تفاعل مع بنزو ثايازولات معوضة لتكوين النفثال ايميدات الجديدة. قيست الفعالية البايولوجية للنفثال ايميدات الجديدة ضد بكتريا موجبة لصبغة غرام (ستافيلوكوكس اريوس وستربتوتوكوكس بايوجينز) وبكتريا سالبة لصبغة غرام (اشريشيا كولاي ويسيدوموناس اوريجينوزا) والخميرة كانديدا البيكانس، حيث أثبتت النتائج امتلاكها فعالية بايولوجية ضد الأحياء المجهرية قيد الدراسة.

Introduction

Cyclic imides represent a very important compounds in drug discovery due to their wide spectrum of biological and pharmacological properties (1, 2, 3, 4). Naphthalimides first discovered by Brana and co-workers (5, 6), are DNA-targeted chemotherapeutic agents acting primarily by attacking DNA at some level (synthesis, replication, or processing) (7). Therefore plenty of naphthalimide based anticancer drugs have been synthesized (8, 9, 10), and promising results have been obtained (11). The benzothiazole scaffold is prevalent in a variety of pharmacologically active synthetic and natural compounds exhibiting antimicrobial (12, 13, 14, 15, 16, 17), anticancer (18, 19, 20), anthelmintic (21), and anti-diabetic activity (22). They are widely found in bioorganic and medicinal chemistry with application in drug discovery (23, 24). Taking into consideration the above described beneficial effects of cyclic imides and benzothiazole, we realized that it would be of interest to synthesize novel structural hybrids containing both heterocyclic ring systems.


Materials and Methods

Chemicals used in this work are supplied from BDH and Fluka companies and are used without further purification. Melting points were determined on Gallenkamp capillary melting point apparatus and were uncorrected. FTIR spectra were recorded on SHIMADZU FTIR-8400 Fourier Transform Infrared spectrophotometer using KBr discs. ^1H NMR and ^{13}C NMR spectra were recorded on Bruker 300MHz instrument using deuterated dimethyl sulfoxide (DMSO-d_6) as a solvent and trimethylsilane (TMS) as internal reference.

- **Synthesis of 4-(N-naphthalimidyl) benzoic acid [1]:** The titled compounds were prepared according to literature procedures (25). (0.01mol, 1.98g) of naphthalic anhydride and (0.01mol, 1.37g) of 4-aminobenzoic acid were placed in a wide dry Pyrex tube which was immersed in an oil bath and provided with a thermometer. The oil bath was heated until fusion of the amic acid then temperature was maintained at ten degrees above the melting point of the used amic acid for (30-45) minutes. After cooling to room temperature the obtained solid was boiled with 10% sodium bicarbonate for 20 mins and purified by recrystallization from ethanol. Yield 85%, melting point $> 300^\circ\text{C}$.
- **Synthesis of 4-(N-naphthalimidyl)benzoyl chloride [2]:** A mixture of 10g of 4-(N-naphthalimidyl) benzoic acid and (1.25g, 7.7mL) of thionyl chloride in 100mL round bottomed flask fitted with a double surface reflux condenser carrying a calcium chloride guard tube was heated on water bath with occasional shaking for 1 hour. The excess of thionyl chloride was distilled at 77°C (26) .
- **Synthesis of 4-(N-naphthalimidyl)-N-(substitutedbenzothiazol-2-yl) benz- amide [3-12]:** To the solution of (0.01 mol) of 4-(N-naphthalimidyl)benzoyl chloride in 100mL. of ethanol was added (0.01mol) of substituted benzothiazol. The mixture was refluxed on a water bath for 6 hours. After cooling, the precipitate was collected, washed with distilled water, and recrystallized from suitable solvent(27). Physical properties of prepared compounds are listed in Table (1).
- **Biological Study:** The cup plate method (28) using nutrient agar medium was employed in studying the antimicrobial activity of the prepared imides against four types of bacteria including *Staphylococcus aureous*, *Streptococcus pyogenes*, *Escherichia coli* and *Pseudomonas aureginosa* respectively and yeast *Candida albicans* fungi. Dimethyl form amide (DMF) was used as sample solution which fixed at (0.1 mL) for all compounds using a sterilized cork borer cups were scooped out of agar medium contained in a petri dish which was previously inoculated with the microorganisms. The test compound solution (0.1 mL) was added in the cups and the petri dishes were subsequently incubated at 37°C for 48 hrs. Zones of inhibition produced by each compound was measured in (mm) and the results are listed in Table (3).

Results and Discussion

- **Chemistry:** Since both naphthalimides and benzothiazoles are biologically active components having wide spectrum of medicinal and pharmacological applications, the present work is directed toward synthesis of new compounds containing these two active moieties with expected biological activity. The target of present work is synthesis of new naphthalimides linked to benzothiazole moiety. Performing of this target involved many steps, the first one involved synthesis of 4-(N-naphthalimidyl) benzoic acid [1] via direct reaction between 1,8-naphthalic anhydride and 4-aminobenzoic acid under fusion condition. Compound [1] was introduced in reaction with thionyl chloride in the second step producing 4-(N-naphthalimidyl) benzoyl chloride [2]. Reaction of compound [2] with different substituted benzothiazole in the third step gave the corresponding 4-(N-naphthalimidyl)-N-(substitutedbenzothiazol-2-yl) benzamide [3-12]. The synthetic route of the new compounds is outlined in Scheme (1).



Structures of the prepared compounds were confirmed by FTIR, ^1H NMR and ^{13}C NMR spectra data. FTIR spectrum of compound [1] showed appearance of $\nu(\text{C}=\text{O})$ of carboxylic acid at $(11687) \text{ cm}^{-1}$ and $\nu(\text{C}=\text{O})$ imide at $(1700) \text{ cm}^{-1}$ absorption bands indicating success of imidation reaction. Besides the spectrum showed characteristic absorption bands at $(3325-3461) \text{ cm}^{-1}$ due to $\nu(\text{OH})$ carboxylic acid and $\nu(\text{NH})$ amide and at $(1666) \text{ cm}^{-1}$ due to $\nu(\text{C}=\text{O})$ amide. FTIR spectra of imides [3-12] showed clear absorption bands at $(3270-3365) \text{ cm}^{-1}$ due to $\nu(\text{NH})$ amide, bands at $(1643-1682) \text{ cm}^{-1}$ due to $\nu(\text{C}=\text{O})$ amide and bands at $(1680-1718) \text{ cm}^{-1}$ due to $\nu(\text{C}=\text{O})$ imide. Other bands appeared at $(1588-1665) \text{ cm}^{-1}$, $(1542-1595) \text{ cm}^{-1}$ and at $(625-710) \text{ cm}^{-1}$ due to $\nu(\text{C}=\text{N})$ thiazol, $\nu(\text{C}=\text{C})$ aromatic and $\nu(\text{C}-\text{S})$ thiazol respectively. All details of FTIR spectral data of compounds [3-12] are listed in Table (2)(29). ^1H NMR spectrum of compound [1] showed clear signals at $(\delta = 7.7-8.5) \text{ ppm}$ and at $\delta = 12.3 \text{ ppm}$ belong to aromatic protons and (OH) proton while ^{13}C NMR spectrum of the same compound [1] showed signals at $(\delta = 124.2-136.3) \text{ ppm}$ and 162.8 ppm belong to aromatic carbons and $(\text{C}=\text{O})$ carbons respectively. ^1H NMR spectrum of compound [4] showed clear signals at $\delta = 2.32 \text{ ppm}$, $(\delta = 7.2-8.6) \text{ ppm}$ and $\delta = 11.8 \text{ ppm}$ due to CH_3 aromatic protons and (NH) proton, while ^{13}C NMR spectrum of the same compound [4] showed signals at $\delta = 22 \text{ ppm}$, $(\delta = 118.5-157) \text{ ppm}$, 162.8 ppm and 164.1 ppm belong to CH_3 , aromatic carbons, $(\text{C}=\text{N})$ and $(\text{C}=\text{O})$ carbons respectively. ^1H NMR spectrum of compound [7] showed signals at $\delta = 3.85 \text{ ppm}$, $(\delta = 7.31-8.72) \text{ ppm}$ and $\delta = 10.9 \text{ ppm}$ due to (OCH_3) , aromatic protons and (NH) proton respectively, while ^{13}C NMR spectrum of the same compound [7] showed signals at $\delta = 55.8 \text{ ppm}$, $(\delta = 106.5-153.9) \text{ ppm}$, 157.8 ppm and 165.2 ppm belong to (OCH_3) , aromatic carbons, $(\text{C}=\text{N})$ and $(\text{C}=\text{O})$ carbons respectively. ^1H NMR spectrum of compound [11] showed clear signals at $\delta = 2.2 \text{ ppm}$, $(\delta = 7.43-8.56) \text{ ppm}$ and $\delta = 11.2 \text{ ppm}$ belong to two methyl groups, aromatic protons and (NH) proton respectively, while ^{13}C NMR spectrum of the same compound [11] showed signals at $\delta = 15.4$ and 18.3 ppm belong to two methyl groups and signals at $(\delta = 113.5-140.1) \text{ ppm}$, 141.28 ppm and 163.82 ppm belong to aromatic carbons, $(\text{C}=\text{N})$ and $(\text{C}=\text{O})$ carbons respectively.

- **Biological Study:** Antimicrobial activity of the synthesized imides were tested against four types of bacteria and *Candida albicans* yeast using cup plate method. Most of the tested bacterial organisms showed sensitivity to synthesized imides

compared to standard antibiotics. Inhibition zones caused by each compound was measured in (mm) and the results are listed in Table (3). The results indicated that compounds (8, 9) are highly active against all types of tested bacteria. Compound [10] also highly active against *S. aureus* and *S. pyogenes* and moderate active against *E. coli*. Compound (10) is highly active against *Candida albicans* yeast while compound (8) showed moderate activity against *Candida albicans* fungi. The rest of imides were found to be moderately active against the tested organisms.

Table (1) Physical properties for prepared imides [3-12]











| Comp. No. | Compound structure | Color | Melting point °C | Yield % | Solvent of Recr. |
|-----------|---|-----------------|------------------|---------|------------------|
| 3 |  | White | 230-232 | 75 | Ethanol |
| 4 |  | Light brown | 255-257 | 83 | Ethanol |
| 5 |  | Off white | 295-297 | 70 | Ethanol |
| 6 |  | Yellow | >300 | 65 | Ethanol |
| 7 |  | Violet | >300 | 85 | Ethanol |
| 8 |  | Gray | 220-222 | 75 | Ethanol |
| 9 |  | Deep yellow | >300 | 68 | Ethanol |
| 10 |  | Greenish yellow | >300 | 78 | Ethanol |
| 11 |  | Brown | >300 | | Ethanol |
| 12 |  | Brown | 245-247 | 87 | Ethanol |

Table (2) FTIR spectral data of naphthalimides [3-12]

| Comp. No. | FTIR spectral data cm ⁻¹ | | | | | | |
|-----------|-------------------------------------|--------------|--------------|-----------------|-----------------|-----------------|---|
| | v(N-H) amide | v(C=O) amide | v(C=O) imide | v(C=N) thiazole | v(C=C) aromatic | v(C-S) thiazole | others |
| 3 | 3310 | 1678 | 1680 | 1610 | 1585 | 670 | - |
| 4 | 3382 | 1682 | 1715 | 1638 | 1560 | 675 | - |
| 5 | 3270 | 1675 | 1718 | 1600 | 1589 | 677 | v(C-Cl)aromatic 1095 |
| 6 | 3285 | 1666 | 1685 | 1625 | 1593 | 688 | v(C-NO ₂) 1500, 1410 |
| 7 | 3360 | 1665 | 1718 | 1665 | 1542 | 710 | v(C-O-C) 1220,1175 |
| 8 | 3295 | 1650 | 1695 | 1635 | 1589 | 708 | v(C-Cl)aromatic 1068 |
| 9 | 3310 | 1670 | 1689 | 1615 | 1593 | 678 | v(C-NO ₂) 1440, 1327 v(C-Cl) 1055 |
| 10 | 3325 | 1678 1643 | 1698 | 1595 | 1581 | 668 | - |
| 11 | 3365 | 1678 | 1710 | 1645 | 1578 | 673 | - |
| 12 | 3305 | 1680 | 1705 | 1588 | 1595 | 625 | - |

Table (3) Inhibition zones for some of the prepared compounds measured in (mm)

| Comp. No. | Gram-positive bacteria | | Gram-negative | | Yeast |
|------------|------------------------|--------------------|----------------|----------------------|-------------------------|
| | <i>S. aureus</i> | <i>S. pyogenes</i> | <i>E. coli</i> | <i>P. aeruginosa</i> | <i>Candida albicans</i> |
| 3 | 12.5 | 10 | 9.5 | 8.6 | 8.8 |
| 5 | 13 | 11.2 | 10.8 | 11.5 | 9.5 |
| 6 | 10.2 | 8.5 | 9.5 | 8.5 | 7.2 |
| 8 | 14.6 | 12.5 | 14 | 12.4 | 12 |
| 9 | 14.8 | 12.5 | 15.2 | 13.4 | 8.5 |
| 10 | 16 | 11.5 | 10.5 | 8.2 | 15.7 |
| ampicillin | 17 | 12.5 | 12 | 14 | - |
| DMSO | - | - | - | - | - |

References

- Abdel-Aziz, A. A. M. (2007). Novel and versatile methodology for synthesis of cyclic imides and evaluation of their cytotoxic, DNA binding, apoptotic inducing activities and molecular modeling study. *Eur. J. Med. Chem.*, 42: 614-618.
- Lima, E. O.; Queiroz, E. F.; Andricopulo, A. D.; Nunes, R. J.; Yunes, R. A. & Cechinel, V. (1999). Evolution of antifungal activity of N-aryl malimides and N-phenyl alkyl-3,4-dichloromalimides. *Bol. Soc. Chil. Quim.*, 44: 185-189.
- Borchhardt, D. M.; Castilho, M. S. & Andricopulo, A. D. (2007). Classical and fragment-based hologram structure-activity relationship for a series of analgesic cyclic imides. *Lett. Drug. Des. Dis.*, 5:57-61.
- Al-Azzawi, A. M. & Hamed, A. S. (2014). Synthesis and antimicrobial screening of new naphthalimides linked to oxadiazole, thiadiazole and triazole cycles. *Int. J. Res. Pharm. Chem.*, 4(2): 283-290.
- Brana, M. F. & Ramos, A. (2001). Naphthalimides as anti-cancer agents: synthesis and biological activity. *Cur. Med. Chem. Anti-Cancer Agents*, 1:237- 2556.
- Brana, M. F.; Cacho, M.; Gradillas, A.; de Pascual-Teresa, B. & Ramos, A. (2001). Intercalators and anticancer drugs. *Cur. Pharm. Des.*, 7: 1745-1780.
- Frei, E.; Teicher, B. A.; Holden, S. A.; Cathcart, K. N. S. & Wang, Y. (1988). Preclinical studies and clinical correlation of the effect of alkylation dose. *Cancer Res.*, 48: 6417-6423.
- Hariprakash, H. K.; Kosakowska-Cholody, T.; Meyer, C.; Cholody, W. M.; Stinson, S. F.; Tarasova, N. I. & Michejda, C. J. (2007). Optimization of naphthalimide-imidazoacridone with potent antitumor activity leading to clinical candidate (HKH40A, RTA 502). *J. Med. Chem.*, 50: 5557-5560.

9. Sami, S. M.; Dorr, R. T.; Alberts, D. S.; So'lyom, A. M. & Remers, W. A. (2000). Analogues of amonafide and azonafide with novel ring systems. *J. Med. Chem.*, 43: 3067-3073.
10. Zhu, H.; Huang, M.; Yang, F.; Chen, Y.; Miao, Z.; Qian, X.; Xu, Y.; Qin, Y.; Luo, H.; Shen, X.; Geng, M.; Cai, Y. J. & Ding, J. (2007). R16, a novel amonafide analogue, induces apoptosis and G2-M arrest via poisoning topoisomerase II. *Mol. Cancer Ther.*, 6: 484-495.
11. Aibin, W.; Yufang, X. & Qian, X. (2010). Novel naphthalimide-indomethacin hybrids as potential antitumor agents: effects of linkers on hypoxic/oxic cytotoxicity and apoptosis-inducing activity, *Monatshefte für Chemie- Chemical Monthly*, 141: 893-899.
12. Seenaiyah, D.; Reddy, P. R.; Reddy, G. M.; Padmaja, A.; Padmavathi, V. & Krishna, N. S. (2014). Synthesis, antimicrobial and cytotoxic activities of pyrimidinyl benzoxazole, benzothiazole and benzimidazole. *Eur. J. Med. Chem.*, 77:1-7.
13. Malik, J. K.; Manvi, F. V.; Nanjwade, B. K.; Singh, S. & Purohit, P. (2010). Review of the 2-amino substituted benzothiazoles: different methods of the synthesis. *Pharm. Lett.*, 2: 347-359.
14. Sharma, N. K. & Priyanka, K. K. (2010). Benzothiazole: the molecule of diverse biological activities. *Int. J. Cur. Pharm. Res.*, 2: 1-6.
15. Rostamizadeh, S. & Housaini, S. A. G. (2005). Microwave-assisted preparation of 2-substituted benzothiazoles, *Phosphorus Sulfur Silicon Relat. Elem.*, 180: 1321-1326.
16. Patil, S. S. & Bobade, V. D. (2010). Simple and efficient one-pot synthesis of 2-substituted benzoxazole and benzothiazoles. *Synth. Com.*, 40: 206-212.
17. Al-Qalaf, F.; Mekheimer, R. A. & Sadek, K. U. (2008). Cerium (IV) ammonium nitrate (CAN) catalyzed one-pot synthesis of 2-arylbenzothiazoles. *Molecules*, 13: 2908-2914.
18. Guo, H. Y.; Li, J. C. & Shang, Y. L. (2009). A simple and efficient synthesis of 2-substituted benzothiazoles catalyzed by H₂O₂/HCl. *Chin. Chem. Lett.*, 20: 1408-1410.
19. Azarifar, D.; Maleki, B.; Setayeshnazar, A. & Simple, A. (2009). Microwave-assisted, and solvent-free synthesis of 2-arylbenzothiazoles by acetic acid-promoted condensation of aldehydes with 2-aminothiophenol in air. *Phosphorus Sulfur Silicon Relat. Elem.*, 184: 2097-2102.
20. Pratap, U. R.; Mali, J. R.; Jawale, D. V. & Mane, R. A. (2009). Baker's yeast catalyzed synthesis of benzothiazoles in an organic medium. *Tetrahedron Lett.*, 50: 1352-1354.
21. Devmurari, V. P. & Ghodasara, T. J. (2010). Synthesis and antibacterial activity of some substituted 2-phenyl benzothiazole. *Arch. Appl. Sci. Res.*, 2: 198-203.
22. Boger, D. L. (1978). A convenient preparation of 2-substituted benzothiazoles. *J. Org. Chem.*, 43: 2296-2297.
23. Khokra, S. L.; Arora, K.; Mehta, H.; Aggarwal, A. & Yadav, M. (2011). Common methods to synthesize benzothiazole derivatives and their medicinal significance. *Int. J. Pharm. Sci. Res.*, 2: 1356-1377.
24. Chugunova, E.; Boga, C.; Sazykin, I.; Cino, S.; Micheletti, G.; Mazzanti, A.; Sazykina, M.; Burirov, A.; Khmelevtsova, L. & Kostina, N. (2015). Synthesis and antimicrobial activity of novel structural hybrids of benzofuroxan and benzothiazole derivatives. *Eur. J. Med. Chem.*, 93: 349-359.
25. Peng, Y.; Song, G. & Qian, X. (2001). Imitation of cyclic carboxylic anhydrides under microwave irradiation. *Synth. Comm.*, 31: 1927-1931.
26. Vogel, A. I. (1996). *Textbook of practical organic chemistry*, 5th ed., Longman.
27. Oruc, E. E.; Rollas, S.; Kandemirli, F.; Shvets, N. & Dimoglo, A. S. (2004). 1,3,4-Thiadiazole derivatives. Synthesis, Structure elucidation, and structure-antituberculo- sis activity relationship investigation. *J. Med. Chem.*, 47:6760-6767.
28. Chavan, A. & Pai, N. (2007). Synthesis and Biological Activity of N-Substituted-3-chloro-2-azetidiones. *Molecules*, 12: 2467-2477.
29. Silverstein, R. M. & Bassler, G. C. (1981). *Spectrometric identification of organic compounds*", 4th Ed., John and Sons.