

Changes in levels of Interleukin-6 (IL-6) and Interleukin-8 (IL-8) in the Serum of Preterm Delivery Pregnant Women Affected by Gingivitis

Doi: <https://doi.org/10.32007/jfacmedbagdad.2152>

Shaden H. Maddah* BDS
Ghada I. Taha* BSc, MSc, PhD



This work is licensed under a [Creative Commons Attribution-Noncommercial 4.0 International License](https://creativecommons.org/licenses/by-nc/4.0/)

Abstract

Background: Periodontal disease increases local and systemic inflammatory responses in pregnant women, which may lead to premature delivery.

Objectives: To detect maternal serum levels of proinflammatory cytokine (Interleukine-6, Interleukin-8) with preterm delivery in pregnant women suffering from gingivitis.

Cases and methods: In this case-control study, a total of 85 pregnant women, were included in the study, of whom 55 had gingivitis: 25 preterm deliveries and 30 full-term deliveries. Thirty healthy pregnant women with healthy gingiva were in the control group. They were selected from Al-Ramadi Teaching Hospital for Gynecology and Pediatrics, from November 2022 to May 2023. Blood samples were collected from all participants and the biomarkers cytokines Interleukine-6 and Interleukin-8 were detected by enzyme-linked immunosorbent assay (ELISA) kits.

Results: The current study showed an increased level of interleukin-6 and interleukin-8 in the pre-term delivery gingivitis group followed by full-term-delivery pregnant groups having gingivitis compared with the control group, with statistically highly significant differences ($p < 0.05$).

Conclusion: Pregnant women with gingivitis with the highest levels of Interlukine-6 and Interlukine-8 were more prone to have premature delivery than those with lower levels of Interleukins and those without gingivitis.

Keywords: full-term delivery; Gingivitis; Interleukine-6; Interleukin-8; Periodontal disease; preterm delivery.

J Fac Med Baghdad
2023; Vol.65, No. 4
Received: June, 2023
Accepted: Aug, 2023
Published: Jan., 2024

Introduction :

Periodontal diseases are bacterial infections of tooth-supporting tissues that inflame and destroy the periodontium. The complex bacterial structure of a protective and adhesive matrix is what maintains this bacterial infection affixed to the periodontal tissue (1). Bacterial infection severely degrades the periodontium and creates a continual and systemic assault with bacterial compounds, and host-derived inflammatory mediators may start and perpetuate systemic diseases (2). Because periodontal disease affects 90% of the population (either gingivitis or periodontitis), dental health and its relationship to systemic health are global health problems (3). New studies links periodontal disease to heart disease, diabetes, respiratory infections, and pregnancy complications. Pregnancy can also negatively impact gingival health,

as variations in hormone levels during pregnancy can induce gingivitis. Pregnant women may also have a higher risk of developing new periodontal diseases, as preexisting periodontal diseases become active during pregnancy (4,5)

Hill suggested that early-stage microorganisms linked to gingivitis and periodontitis may trigger a placental response, leading to premature delivery. Hormonal alterations are well-established to cause a host response to infection, but the growth and alterations of sub-gingival biofilm during pregnancy are poorly understood (6).

Forty percent of pregnant women have periodontitis (7,8), and their babies are seven times more likely to be premature or underweight. Pregnancy hormones cause inflammation, which promotes periodontal disease. Gingivitis affects 50-70% of pregnant women due to hormonal changes. Pregnancy-induced plasma progesterone and estrogen levels may alter subgingival bacteria, the maternal immune system, and pro-inflammatory mediator production (9,10). Cytokines are well-known mediators that govern local, systemic,

**Corresponding Author:* Dept. of Basic Science in Oral Microbiology, College of Dentistry, University of Baghdad. *shaden.husham@gmail.com
Ghada_ibraheem@codental.uobaghdad.edu.iq

and inflammatory responses and play a role in numerous biological processes, such as inflammation, by regulating the length and intensity of the inflammatory response (11). Cytokines control homeostasis and inflammation. Female sex hormones' ability to enhance cytokines production by human gingival fibroblasts at high concentrations comparable to pregnant women's plasma suggested that they may contribute to periodontal disease progression during pregnancy (12,13). Interleukin-8 is a cytokine generated in response to inflammation by several immunological inflammatory cells. Interleukin-8 functions begin with neutrophil activation and play a role in polymorphonuclear neutrophils (PMNs) migration to the inflammatory region (14). Chronic and excessive IL-8-mediated chemotactic and triggering impact on neutrophils in the inflammatory gingiva may generate local periodontal tissue injury (15). According to Shahshahan's research, maternal blood interleukin-6 and interleukin-8 concentrations may be employed as acceptable indicators for predicting preterm labor (16). In industrialized and developing nations, preterm birth (PTB) is a major public health issue and the most important biological factor of neonatal survival. They affect families and the healthcare system. Preventable risk factors for preterm birth must be continuously sought (17). Preterm birth is important because it predicts increased mortality and morbidity in infants born with this condition and reflects the mother's exposure to other risk factors like socioeconomic conditions, malnutrition, and diseases (18). In 1996, Offenbacher et al. found that periodontal disease is a risk factor for preterm low birth weight (9). Hill, also observed that periodontal bacteria may cause upper genital tract infections in pregnant women, resulting in premature delivery (6). Thus, this study aims to detect maternal serum levels of proinflammatory cytokine (Interleukin-6, Interleukin-8) with preterm delivery in pregnant women suffering from gingivitis.

Subjects, materials, and methods

The current study was performed on 85 pregnant women 55 of whom had gingivitis and was divided into two groups: 25 preterm deliveries and 30 full-term deliveries, in addition to 30 healthy pregnant women with healthy gingiva as a control group.

The cases were taken from Al-Ramadi Teaching Hospital for Gynecology and Pediatrics, Al-Ramadi, Iraq, during the period November 2022 to May 2023.

Inclusion Criteria for the gingivitis groups:

1. Pregnant women in their third trimester months (term pregnancies were those who delivered at ≥ 37 weeks of gestation, preterm pregnancies were those who delivered at < 37 weeks of gestation) (17).
2. Good general health
3. 18 – 42 years old
4. Iraqi nationality

5. Has at least 20 teeth in the oral cavity
6. Having gingivitis with Pocket depth ≤ 3 mm

Exclusion Criteria:

1. Periodontal therapy for the past 3 months.
2. History of smoking or alcohol drinking.
3. Use of antibiotics during the past three months.
4. Wearing orthodontic appliances or prosthodontics
5. Twin pregnancy.
6. Genital tract infection.
7. Urinary tract infection.
8. No systemic or autoimmune disease.
9. Not diabetic, no history of allergy or rhinosinusitis.

Oral Examination:

A clinical examination was performed by a dentist for every subject under the study.

The periodontal status of all teeth was assessed using a periodontal probe, the periodontal parameters include (8,9): Gingival index (GI), bleeding on probing (BOP), and probing pocket depth (PPD).

1. Gingival Index: The occurrence of gingival inflammation at 4-surfaces of each tooth was assessed using the criteria of the gingival index system (9).

Score 0: normal gingiva

Score 1: mild inflammation, slight change in color, slight edema, no bleeding on probing.

Score 2: moderate inflammation, redness, edema and glazing, bleeding on probing.

Score 3: severe inflammation, marked redness and edema, ulceration, tendency to spontaneous bleeding.

2. Bleeding on Probing: A periodontal probe is inserted into the base of the periodontal pocket/sulcus for six surfaces of each tooth (8,9) (mesiofacial, facial, distofacial, mesiolingual, lingual, and distolingual) and is moved lightly along the tooth (root) surface. After probing, the site was given a score:

a. If bleeding occurred within 30 seconds (score 1)

b. For the non-bleeding site (Score 0)

3. Probing Pocket Depth: Refers to the distance from the gingival margin to the most apical penetration of the periodontal probe. The sites of measurements were Mesiofacial, facial, distofacial, mesiolingual, lingual, and distolingual (8,9). No pressure was used to insert the periodontal probe.

Blood Sample Collection: Five millilitres of venous blood were collected from each subject under aseptic conditions before delivery, moved to a sterile gel tube, and serum was separated by centrifugation at 3000rpm, divided into small aliquots, and stored until analysis (18).

Enzyme-Linked Immunosorbent Assay (ELISA): Detection of pro-inflammatory biomarkers by using Human Interleukin-6 and Interleukin-8 enzyme immunoassay kit (ELK Biotechnology, company) in serum samples.

Principle of the Procedure: This kit uses sandwich enzyme immunoassay. This package includes a

microtiter plate pre-coated with either IL-6 or IL-8 antibody. Microtiter plate wells containing standards or samples get a biotin-conjugated IL-6 or IL-8 antibody. Incubate microplate wells with avidin-HRP. TMB substrate solution only colours wells containing IL-6, IL-8, biotin-conjugated antibodies, and enzyme-conjugated avidin. Sulfuric acid blocks the enzyme-substrate reaction, and the color change is measured at 450 nm±10 nm. The concentration of biomarkers in the samples is then determined by comparing the optical density (OD) of the samples to the standard curve.

Statistical analysis

The Statistical Package for Social Sciences (SPSS) version 26 and Microsoft Excel 2010 were employed for data processing. Differences between group means were studied. Fisher's exact test, ANOVA, and Pearson correlation were also used.

Results

Table 1 compares the three study groups in terms of their age, bleeding on probing, Interlukine-6, and Interlukine-8. The differences between the mean ages of the three study groups were not statistically significant (p>0.05). The level of BOP % in preterm-delivery pregnant groups having gingivitis was higher than in the full-term delivery group having gingivitis and the healthy pregnant control group, with significant differences (p<0.05). The mean±SE was (0.19±0.019), (0.17±0.016), and (0.09±0.028), respectively. The mean±SE value of IL-6 level in pre-term delivery was higher than in full-term-delivery pregnant groups having gingivitis compared with healthy pregnant control group, with significant differences (p<0.05). The mean±SE was (45.2±7.90), (22.2±9.64), and (7.7±0.75), respectively. Likewise, the mean±SE IL-8 levels in the pre-term delivery group were higher than the full-term-delivery groups with gingivitis and the control group, (128.5±11.72), (115.1±0.93) and (13.9±1.44) respectively, with significant differences (p<0.05).

Table 1: The mean value of age, BOP%, and the biomarkers in the three groups

Variable	Group			ANOVA test (p. value)
	Full term Mean±SE	Pre term Mean±SE	Controls Mean±SE	
Age (year)	25.07±1.097	26.12±1.400	27.73±1.250	F=0.553 0.296
Bleeding on probing (BOP)%	0.17±0.016	0.19±0.019	0.09±0.028	F=5.751 0.005
Interlukine-6 (ng/ml)	22.2±9.64	45.2±7.90	7.7±0.75	F=7.950 0.001
Interlukine-8 (ng/ml)	115.1±0.93	128.5±11.72	13.9±1.44	F=77.750 0.0001

Table 2 reveals the correlation between IL-6 and IL-8 in Full-term and Preterm delivery having gingivitis

groups. There was a negative non-significant correlation between IL-6 and IL-8 in both groups (r=-0.015), (r= -0.072), (p>0.05).

Table 2: Correlation between IL-6 and IL-8 in the gingivitis study groups

Biomarkers	Full term		Pre term	
	r	p	r	p
IL-6 and IL-8	-0.01	0.943	-0.072	Ns 0.706
	5	Ns		

r= Pearson correlation, Ns= non-significant (p>0.05)

Discussion

The present study showed the percentage of BOP % in preterm-delivery pregnant groups having gingivitis was significantly higher than in the full-term delivery group having gingivitis compared with the healthy pregnant group. This indicates that the pregnant women in this study had suffered the bad outcomes of gingivitis.

The significantly higher level of interleukin-6 in preterm-delivery than full-term delivery pregnant groups having gingivitis compared with the healthy pregnant woman group suggests that the local generation of these cytokines in the periodontal pocket due to periodontal disease may raise blood concentration and amniotic fluid concentration (19,20). Several prior studies back up these findings, showing that any infection that causes chorioamnionitis and leads to higher cytokine levels in amniotic fluid, particularly high levels of IL-6, may cause preterm rupture of membranes and delivery (20,21). The results are consistent with those of Latorre Uriza et al., who found that cytokines (interleukin-2, IL-6, IL-10, and TNF-α) increased as the periodontal disease progressed from healthy through gingivitis to chronic periodontitis, with the highest levels in pregnant patients (22). Greig et al. and Von Minckwitz *et al.* revealed that periodontal disease patients release proinflammatory cytokines and mediators into the gingival crevicular fluid (GCF) or the bloodstream, reaching the placenta-fetus interface (23). IL-1, IL-6, and TNF-α stimulate chorion prostaglandin synthesis, cervical ripening, and uterine contraction, increasing preterm labor risk (24,25). An earlier study by Van Dyke et al. agrees with the present investigation, showing that periodontal infection increases systemic inflammatory markers IL-3, IL-4, IL-5, IL-6, IL-10, and TNF-α (26). Even though elevated levels of these mediators in the serum, amniotic fluid, and GCF have been linked to a variety of adverse pregnancy outcomes. Deortbudak *et al.* and Madianos *et al.* found no evidence that periodontitis patients with elevated mediators in GCF, serum, or amniotic fluid had pregnancy difficulties (27,28). Our findings showed a noticeable increase in the IL-8

levels in pre-term delivery compared with full-term-delivery pregnant groups having gingivitis and the control group. This is a line with a previous study by Hebisch *et al.* who demonstrated that periodontal infections linked with preterm delivery during cervical dilatation increased maternal serum IL-8 and IL-6 levels in serum, amniotic fluid, maternal, and retroplacental blood (29). Although the increased percentage of periodontopathic bacteria in subgingival plaque might have produced the elevation of such cytokines, the causes for the rise were not obvious and should be investigated further (29). Another study, found these cytokines IL-6, IL-8, and TNF- α have a rise during periodontal disease and also chorioamnionitis (30). Furthermore, Early signals to enhance the production of pro-inflammatory molecules such as IL-6 and IL-8, according to Hasegawa *et al.*, lead to uterine contraction and cervical ripening, which leads to parturition (21). Monocytes, fibroblasts, lymphocytes, and endothelial cells release interleukin-8, which draws neutrophils to the infection site. Its pro-inflammatory and neutrophil chemotactic characteristics may cause periodontitis (31,32). Lagdive *et al.* discovered higher IL-8 levels in gingival tissue, which has been linked to progressive periodontitis (15). Actually, pregnancy increases gingival edoema and bleeding. Pregnancy-induced high oestrogen levels increase gingival inflammation and gram-negative anaerobe growth., particularly *Prevotella nigrescens* (33,34). Nevertheless, the oestrogen hormone inhibits neutrophilic chemotaxis as well as enzyme activity in periodontal tissues (33, 35). Periodontal tissues upregulate endothelium adhesion molecules, enhance chemotactic agent secretion, and exacerbate leukocyte chemotaxis in response to bacterial infection, secreting IL-8 and macrophage chemoattractant peptide-1 to reduce estrogen concentrations (36-39).

Conclusion

The findings of this study suggest a potential association between maternal oral health (gingivitis) and adverse pregnancy outcomes (premature delivery) mediated by elevated levels of proinflammatory cytokines (Interleukin-6 and Interleukin-8).

Conflicts of Interest: None

Author contributions:

Shaden Husham Maddah: a master's student, primarily responsible for conceptualizing the research topic, conducting the experiments, collecting and analyzing data, and writing the initial draft of the manuscript, also interpretation of results and the preparation of the final manuscript.

Assist. Prof. Dr. Ghada Ibrahim Taha: The supervisor provided substantial contributions to the design and

execution of the study. She offered continuous guidance throughout the research process, from methodology development to result analysis. Also critically reviewed and revised the manuscript, ensuring the integrity and accuracy of the work.

Both authors read and approved the final manuscript.

Authors' declaration:

Conflicts of Interest: The authors declare no conflict of interest.

We confirm that all the Figures and Tables in the manuscript belong to the current study. Besides, the Figures and images, which do not belong to the current study, have been given permission for re-publication attached to the manuscript. Authors sign on ethical consideration's approval-Ethical Clearance: The project was approved by the local ethical committee in (the college's in-house ethics committee of Dentistry, University of Baghdad (Project No.)) according to the code number (713822).

References

1. Abdulkareem AA, Al-Taweel FB, Al-Sharqi AJB, Gul SS, Sha A, Chapple ILC. Current concepts in the pathogenesis of periodontitis: from symbiosis to dysbiosis. *J Oral Microbiol.* 2023 Apr 2;15(1):2197779. <https://doi.org/10.1080/20002297.2023.2197779>
2. Al-Alousi HW. Frequency of Salivary Human Cytomegalovirus in Iraqi Patients with Chronic Periodontitis. *Journal of the Faculty of Medicine Baghdad.* 2013; 55(2), 162-165. <https://doi.org/10.32007/jfacmedbagdad.552647>
3. Al-obaidi M, Al-Ghurabi BH. Association between *Aggregatibacter actinomycetemcomitans* bacterial load and NLRP3 inflam-masome activation in periodontitis patients with Diabetes. *Journal of the Faculty of Medicine Baghdad,* 2023;64(4), 268-275. <https://doi.org/10.32007/jfacmedbagdad.6441972>
4. Hameed NM, Alwaheb AM. Trajectories of Salivary Hormones in Pregnant Women with Anxiety and their Effect on Gingival Health Condition. *Indian Journal of Forensic Medicine & Toxicology,* 2022;16(1), 1458-1465. <https://doi.org/10.37506/ijfnt.v16i1.17706>
5. Ali SS, Ali MS, AL-Gebori AM. The influence and assessment of E-cadherin sera levels with periodontitis and gingivitis. *Journal of Survey in Fisheries Sciences,* 2023; 10(3S), 3151-3161.
6. Hill GB. Preterm birth: associations with genital and possibly oral microflora. *Ann Periodontol.* 1998 Jul;3(1):222-32. <https://doi.org/10.1902/annals.1998.3.1.222>
7. Spahr A, Klein E, Khuseyinova N, Boeckh C, Muche R, Kunze M, et al. Periodontal infections and coronary heart disease: role of periodontal bacteria and importance of total pathogen burden in the Coronary

- Event and Periodontal Disease (CORODONT) study. *Arch Intern Med.* 2006 Mar 13;166(5):554-9. .
<https://doi.org/10.1001/archinte.166.5.554>
8. Lief S, Boggess KA, Murtha AP, Jared H, Madianos PN, Moss K, et al. The oral conditions and pregnancy study: periodontal status of a cohort of pregnant women. *J Periodontol.* 2004 Jan;75(1):116-26. <https://doi.org/10.1902/jop.2004.75.1.116>
9. Offenbacher S, Katz V, Fertik G, Collins J, Boyd D, Maynor G, et al. Periodontal infection as a possible risk factor for preterm low birth weight. *J Periodontol.* 1996 Oct;67(10 Suppl):1103-13.
<https://doi.org/10.1902/jop.1996.67.10s.1103>
10. AL- Muhamadawy AK. Periodontal health status and treatment needs among Iraqi dental students. *Journal of the Faculty of Medicine Baghdad,* 2010;51(4), 378-381.
11. Ertugrul AS, Sahin H, Dikilitas A, Alpaslan N, Bozoglan A. Comparison of CCL28, interleukin-8, interleukin-1 β and tumor necrosis factor-alpha in subjects with gingivitis, chronic periodontitis and generalized aggressive periodontitis. *J Periodontal Res.* 2013 Feb;48(1):44-51.
<https://doi.org/10.1111/j.1600-0765.2012.01500.x>
12. Daily ZA, Mohammed AN. Periodontal Health Status and Assessment of Osteocalcin levels in Saliva of Diabetic Patients and Systemically Healthy Persons (Comparative study). *Journal of Baghdad College of Dentistry,* 2017; 29(1), 89-95.
<https://doi.org/10.12816/0038632>
13. Yokoyama M, Hinode D, Masuda K, Yoshioka M, Grenier D. Effect of female sex hormones on *Campylobacter rectus* and human gingival fibroblasts. *Oral Microbiol Immunol.* 2005 Aug;20(4):239-43.
<https://doi.org/10.1111/j.1399-302X.2005.00222.x>
14. Paul AM, Victor DJ, Gana Prakash PS. Role of Interleukin-8 in periodontal disease. *International Journal of Clinical Dental Science,* 2012;3.
15. Lagdive SS, Marawar PP, Byakod G, Lagdive SB. Evaluation and comparison of interleukin-8 (IL-8) level in gingival crevicular fluid in health and severity of periodontal disease: a clinico-biochemical study. *Indian J Dent Res.* 2013 Mar-Apr;24(2):188-92.
<https://doi.org/10.4103/0970-9290.116675>
16. Shahshahan Z, Hashemi L. Maternal serum cytokines in the prediction of preterm labor and response to tocolytic therapy in preterm labor women. *Adv Biomed Res.* 2014 May 28;3:126.
<https://doi.org/10.4103/2277-9175.133243>
17. Dasanayake AP. Poor periodontal health of the pregnant woman as a risk factor for low birth weight. *Ann Periodontol.* 1998 Jul;3(1):206-12.
<https://doi.org/10.1902/annals.1998.3.1.206>
18. Al Mosawi A O. Need and Demand According to Basic Periodontal Index to Patient with Periodontal Disease Attending Department of Periodontology at the College of Dentistry/University of Baghdad. *Latin American Journal of Pharmacy,* 2021;40(SI), 271-275.
19. Sanz M, Kornman K. Working group 3 of joint EFP/AAP workshop. Periodontitis and adverse pregnancy outcomes: consensus report of the Joint EFP/AAP Workshop on Periodontitis and Systemic Diseases. *J Clin Periodontol.* 2013 Apr;40 Suppl 14:S164-9. <https://doi.org/10.1111/jcpe.12083>
20. Stevens VL, Hoover E, Wang Y, Zanetti KA. Pre-Analytical Factors that Affect Metabolite Stability in Human Urine, Plasma, and Serum: A Review. *Metabolites.* 2019 Jul 25;9(8):156.
<https://doi.org/10.3390/metabo9080156>
21. Hasegawa K, Furuichi Y, Shimotsu A, Nakamura M, Yoshinaga M, Kamitomo M, et al. Associations between systemic status, periodontal status, serum cytokine levels, and delivery outcomes in pregnant women with a diagnosis of threatened premature labor. *J Periodontol.* 2003 Dec;74(12):1764-70.
<https://doi.org/10.1902/jop.2003.74.12.1764>
22. Keelan JA, Blumenstein M, Helliwell RJ, Sato TA, Marvin KW, Mitchell MD. Cytokines, prostaglandins, and parturition--a review. *Placenta.* 2003 Apr;24 Suppl A:S33-46.
<https://doi.org/10.1053/plac.2002.0948>
23. von Minckwitz G, Grischke EM, Schwab S, Hettlinger S, Loibl S, Aulmann M, et al. Predictive value of serum interleukin-6 and -8 levels in preterm labor or rupture of the membranes. *Acta Obstet Gynecol Scand.* 2000 Aug;79(8):667-72. PMID: 10949232. <https://doi.org/10.1034/j.1600-0412.2000.079008667.x>
24. Lewis DF, Barrilleaux PS, Wang Y, Adair CD, Baier J, Kruger T. Detection of interleukin-6 in maternal plasma predicts neonatal and infectious complications in preterm premature rupture of membranes. *Am J Perinatol.* 2001 Nov;18(7):387-91.
<https://doi.org/10.1055/s-2001-18694>
25. Latorre Uriza C, Velosa-Porras J, Roa NS, Quiñones Lara SM, Silva J, Ruiz AJ, et al. Periodontal Disease, Inflammatory Cytokines, and PGE2 in Pregnant Patients at Risk of Preterm Delivery: A Pilot Study. *Infect Dis Obstet Gynecol.* 2018 Aug 1;2018:7027683.
<https://doi.org/10.1155/2018/7027683>
26. Greig PC, Murtha AP, Jimmerson CJ, Herbert WN, Roitman-Johnson B, Allen J. Maternal serum interleukin-6 during pregnancy and during term and preterm labor. *Obstet Gynecol.* 1997 Sep;90(3):465-9.
[https://doi.org/10.1016/S0029-7844\(97\)00294-9](https://doi.org/10.1016/S0029-7844(97)00294-9)
27. Dörtbudak O, Eberhardt R, Ulm M, Persson GR. Periodontitis, a marker of risk in pregnancy for preterm birth. *J Clin Periodontol.* 2005 Jan;32(1):45-52. <https://doi.org/10.1111/j.1600-051X.2004.00630.x>
28. Madianos PN, Bobetsis YA, Offenbacher S. Adverse pregnancy outcomes (APOs) and periodontal disease: pathogenic mechanisms. *J Clin Periodontol.*

2013 Apr;40 Suppl 14:S170-80.

<https://doi.org/10.1111/jcpe.12082>

29. Hebisch G, Grauaug AA, Neumaier-Wagner PM, Stallmach T, Huch A, Huch R. The relationship between cervical dilatation, interleukin-6 and interleukin-8 during term labor. *Acta Obstet Gynecol Scand.* 2001 Sep;80(9):840-8.

<https://doi.org/10.1034/j.1600-0412.2001.080009840.x>

30. Gücer F, Balkanli-Kaplan P, Yüksel M, Yüce MA, Türe M, Yardim T. Maternal serum tumor necrosis factor-alpha in patients with preterm labor. *J Reprod Med.* 2001 Mar;46(3):232-6. PMID: 11304864.

31. Van Dyke TE, van Winkelhoff AJ. Infection and inflammatory mechanisms. *J Clin Periodontol.* 2013 Apr;40 Suppl 14:S1-7.

<https://doi.org/10.1111/jcpe.12088>

32. Carta G, Persia G, Falciglia K, Iovenitti P. Periodontal disease and poor obstetrical outcome. *Clin Exp Obstet Gynecol.* 2004;31(1):47-9. PMID: 14998188.

33. Goepfert AR, Jeffcoat MK, Andrews WW, Faye-Petersen O, Cliver SP, Goldenberg RL, et al. Periodontal disease and upper genital tract inflammation in early spontaneous preterm birth. *Obstet Gynecol.* 2004 Oct;104(4):777-83.

<https://doi.org/10.1097/01.AOG.0000139836.47777.6d>

34. Jayaprakash K, Khalaf H, Bengtsson T. Gingipains from *Porphyromonas gingivalis* play a significant role in induction and regulation of CXCL8 in THP-1 cells. *BMC Microbiol.* 2014 Jul 18;14:193.

<https://doi.org/10.1186/1471-2180-14-193>

35. Pan W, Wang Q, Chen Q. The cytokine network involved in the host immune response to periodontitis. *Int J Oral Sci.* 2019 Nov 5;11(3):30. PMID: 31685798; PMCID:PMC6828663

<https://doi.org/10.1038/s41368-019-0064>

36. Ide M, Papapanou PN. Epidemiology of association between maternal periodontal disease and adverse pregnancy outcomes--systematic review. *J Clin Periodontol.* 2013 Apr;40 Suppl 14:S181-94. PMID: 23627328.

<https://doi.org/10.1111/jcpe.12063>

37. Mamalis A, Markopoulou C, Lagou A, Vrotsos I. Oestrogen regulates proliferation, osteoblastic differentiation, collagen synthesis and periostin gene expression in human periodontal ligament cells through oestrogen receptor beta. *Arch Oral Biol.* 2011 May;56(5):446-55. PMID: 21130420.

<https://doi.org/10.1016/j.archoralbio.2010.11.001>

38. Nebel D, Jönsson D, Norderyd O, Bratthall G, Nilsson BO. Differential regulation of chemokine expression by estrogen in human periodontal ligament cells. *J Periodontol Res.* 2010 Dec;45(6):796-802. PMID: 20701669.

<https://doi.org/10.1111/j.1600-0765.2010.01308.x>

39. Straub RH. The complex role of estrogens in inflammation. *Endocr Rev.* 2007 Aug;28(5):521-74. PMID: 17640948.

<https://doi.org/10.1210/er.2007-0001>

How to cite this Article

husham shaden, Taha G. Changes in levels of IL-6 and IL-8 in the serum of preterm delivery pregnant women affected by Gingivitis. *JFacMedBagdad [Internet]. [cited 2023 Dec. 12];65(4). Available from: <https://iqjmc.uobaghdad.edu.iq/index.php/19/JFacMedBaghdad36/article/view/2152>*

التغيرات في مستويات IL-6 و IL-8 في مصل النساء الحوامل اللواتي ولدن قبل الأوان المصابات بالتهاب اللثة

شادن هشام مداح /كلية طب الاسنان /العلوم الاساسية/جامعة بغداد

غادة ابراهيم طه /كلية طب الاسنان /العلوم الاساسية/جامعة بغداد

الخلاصة

الخلفية : أمراض اللثة هي عدوى مهمة عند النساء الحوامل، قد تعمل كعامل خطر للولادة المبكرة عن طريق زيادة الاستجابات الالتهابية الموضعية والجهازية.

المواد والطرق: تم تقسيم إجمالي 85 امرأة حامل إلى ثلاث مجموعات. 25 من الولادات قبل الأوان و 30 من الولادات الكاملة ومصابات بالتهاب اللثة، بالإضافة إلى 30 من النساء الحوامل اللواتي يتمتعن بصحة جيدة مع اللثة السليمة كمجموعة ضابطة. كان متوسط عمر المرضى (1.097 ± 25.07)، (1.400 ± 26.12) سنة، و (1.250 ± 27.73) سنة للمجموعة الضابطة. تم جمع عينة الدم من جميع المشاركين وتم استكشاف المؤشرات الحيوية

السيتوكينات IL-6 و Interleukin-8 بواسطة مجموعات مقايسة الممتز المناعي المرتبط بالإنزيم (ELISA)

النتائج: أظهرت الدراسة الحالية زيادة أعلى في مستويات IL-6 و IL-8 في الولادة المبكرة المصابة بمجموعة التهاب اللثة تليها مجموعة الولادات الكاملة والتي تعاني من التهاب اللثة مقارنة بمجموعة التحكم، مع وجود فروق ذات دلالة إحصائية عالية (p < 0.01)

الخاتمة: كانت النساء الحوامل المصابات بالتهاب اللثة مع أعلى مستويات إنترلوكين 6- وإنترلوكين 8- أكثر عرضة للولادة المبكرة مقارنة بالنساء اللاتي لديهن مستويات منخفضة من الإنترلوكين وأولئك اللواتي لا يعانين من التهاب اللثة.

الكلمات المفتاحية: التهاب اللثة، أمراض اللثة، الولادة المبكرة، الولادة الكاملة، إنترلوكين 6، إنترلوكين 8

Copyright (c) 2023 shaden husham, Ghada Taha