

## Some Chemical Properties of Soil from Southern Iraqi Marshes

\*Faris J. M. Al-Imarah and \*\*Haleemah J .Al-Aradi

\*Dept. of Chem. & Mar. Environ. Poll., Marine Science Centre,  
Basra Univ., Basra- Iraq.

\*\* Dept. of Biological Development of Shatt Al-Arab & N. Arabian Gulf –  
Marine Science Center – Basra University

### Abstract

A study for the evaluation of some properties of soil of certain sites in southern Iraqi marshes Al-Imarah marsh, and Al-Ghebaysh (Central Marsh), Umm El-Neaj and Umm El-Ward (Al-Hewaizah Marsh and Hareer marsh and El-Bargha within Al-Hammar Marsh has been conducted for investigation of their types and suitability for harvesting of certain agricultural crops. Soil samples were collected from each site for a depth of 10 - 90 cm, and transferred to the lab in Marine Science Centre. In the lab the following parameters were determined: pH, Electrical Conductivity (EC), Trace Metals (Cobalt, Co, Iron, Fe, Manganese, Mn, and Nickel, Ni, Total percent of Nitrogen, Dissolved Phosphorous, Sodium and Potassium). Moreover, soil texture was evaluated as % sand, silt and clay for the detection of the type of the soil Results revealed that pH for the studied soil range between lower value of 7.4 in Ghebaysh and 8.1 in Hareer marsh. Conductivity ranged between lower value 4.59 and higher value 13.16 mS/cm in Umm El-Ward and El-Bargah marsh respectively. Concentrations of studied trace metals were alternative among the sites, lower value was 170.526 µg/g dry weight Ni in the soil of Umm El- Neaj and higher value as 7887.727 µg/g dry weight Fe in the soil of Harrer marsh. The soil of highly enriched with nitrogen 0.621% in Umm El-Neaj and phosphorous 1.42 mg/g in Umm El-Ward, while lower values were 0.420% nitrogen in Al-Imarah marsh and 0.85 µg/g P in El-Ghebaysh. Potassium and sodium were ranged between 4.21- 6.35 ppm and 512.13- 661.31 ppm in Al-Imarah respectively. The study revealed that most of the southern marsh lands are suitable for planting of different crops.

**Key Words: Soil, Features, Southern Iraq, Marshes, Soil Texture.**

### Introduction

The Southern Iraqi marshes represent an important part of the

Southern Region Steppe of Iraq, it is called Sahel sedimentary due to deposition of great amounts of salts from Tigris and

Euphrates rivers (Wikipedia,2015). Marshes of southern Iraq occupies 35000km<sup>2</sup>, around 9000 km<sup>2</sup> represent a permanent marsh which filled covered with water over all the year. Other 9000 are non permanent marshes filled with water during rain, while the rest is shallow areas and has no water but cover with *Phragmites australis* and *Typha domingensis* (Mahmood ,2008). The soils of Iraqi marshes are unique type among wetlands and semi lands not only for west Asia but overall the world. They are characterized by unique features even they were deteriorate due to desiccation during the period 1991-2003 in which they turned to dry lands and most of people living their were flee to the main cities around. Animals and plants were cut to very low levels ,only 10% of the marshland were preserved ,60% were become desert and 30 % were used for agriculture (Van *et al* , 1993)

Chemical elements are exist in all soils, Iron (Fe) is the most abundant element (Savhell,1969) , it is present in the range 0.5 – 5 % , and total Cobalt (Co) in the soil is low and range between 100 and 300 µg/g in the basic rocks and 1 – 3 µg /g in

the gypsums rocks (Mitchell ,1972 ). Copper present in very low quantities, the earth crust contains copper in the range 0.1 – 1 % (Venogradov, 1959).

The concentrations of chemical elements are altering in the soil depending upon its texture, iron, manganese and nickel are decreased in sandy soil (Hussain, *et al.*, 1993; Kovda , 1988).

Lands of Iraqi marshes are characterized as being plain lands some of it is fertile which could be used for planting economical crops while others contain high levels of salts. Most lands of Iraqi marshes are covered with water all over the year which produce large quantities of rice and corn while *Phragmites australis* and *Typha domingensis* are growing heavily in non planting areas.

### **Aim of the study**

As marshlands are economically important, this study is aimed at investigation of some physico-chemical features for their soils to identify their suitability for agriculture to enhance the agriculture reality of most field crops specially the cereal crops.

### **Methods and apparatus**

Sites selected for this investigations are 6 stations spread among Southern Iraqi marshes as follows: 1. Al-Imarah marsh, 2. Al-Gebaysh marsh within Central marshes, 3. Umm El-Naaj, and 4. Umm El-Ward within Al-Hwaizah marsh, and 5. Hareer marsh and 6. El-Barghah within Al-Hammar marsh, as shown in Figure 1.

### Collection of samples

Soil samples from the six stations were collected by special tool within the 20 cm depth and put in plastic bags and transferred in cool box to the labs of Marine Science and kept in fridge prior to analysis.

### Laboratory Analysis

In the lab, soil samples thawed, grinded and pH, EC, total percent nitrogen by micro Kildal (Page et al

,1982) and dissolved phosphorus were measured by adopting standard methods. Chemical elements, Cobalt, iron, manganese, and nickel were determined by adopting the atomic absorption method using Atomic Absorption Spectrophotometer type Pye Unicam SP9 according to Sturgeon et al, (1982) while sodium and potassium were determined by flame photometry.

### Results and Discussion

The measured parameters in this study pH, EC, TN, TP and chemical elements Co, Fe, Mn, and Ni in the soils of six different locations within Southern Iraqi marshlands are presented in table 1., while the results of soil texture analysis are shown in figure 2.




Figure 1. Map of Southern Iraqi marshes showing the locations sampling sites

**Table 1. Physio chemical parameters for soil from different sites within Southern Iraqi Marshes**

Site / Parameters	1. Al-Imarah	2 El-Chebays	3 Umm El-Naaj	4 Umm El-Ward	5 Hareer	6 El-Barghah
pH	7.8	7.4↓	7.5	7.7	8.1↑	7.8
EC (d S/cm)	12.95	6.17	8.35	4.59↓	11.95	13.16↑
Co (µg/g)	351.779	364.752	307.804↓	307.804↓	395.748↑	371.532
Fe (µg/g)	7340.13	7541.895	7468.391	7200.318↓	7887.727↑	7526.546
Mn (µg/g)	721.523↓	884.43↑	730.368	736.074	743.311	736.074
Ni (µg/g)	213.115	341.052↑	170.526↓	198.947	193.532	284.21

<b>TN %</b>	<b>0.420↓</b>	<b>0.463</b>	<b>0.621↑</b>	<b>0.511</b>	<b>0.505</b>	<b>0.561</b>
<b>TP (µg/g)</b>	<b>1.03</b>	<b>0.85↓</b>	<b>0.98</b>	<b>1.42↑</b>	<b>1.12</b>	<b>0.94</b>
<b>K (mg/kg)</b>	<b>5.33</b>	<b>4.92</b>	<b>4.21↓</b>	<b>4.74</b>	<b>5.62</b>	<b>6.35↑</b>
<b>Na (mg/kg)</b>	<b>661.31↑</b>	<b>598.52</b>	<b>631.23</b>	<b>600.23</b>	<b>553.42</b>	<b>512.13↓</b>

**Table 2. Analysis of soil texture of selected sites in Southern Iraqi Marsh lands.**

<b>Site / Parameters</b>	<b>1. Al-Imarah</b>	<b>2 El-Chebaysh</b>	<b>3 Umm El-Naaj</b>	<b>4 Umm El-Ward</b>	<b>5 Hareer</b>	<b>6 El-Barghah</b>
<b>Sand %</b>	<b>12.364</b>	<b>6.3704</b>	<b>5.5052</b>	<b>0.9716</b>	<b>5.274</b>	<b>15.082</b>
<b>Silt %</b>	<b>51.312</b>	<b>62.096</b>	<b>55.664</b>	<b>66.208</b>	<b>63.19</b>	<b>43.424</b>
<b>clay %</b>	<b>36.323</b>	<b>31.533</b>	<b>38.831</b>	<b>32.820</b>	<b>31.534</b>	<b>41.494</b>
<b>Type of soil</b>	<b>Silt</b>	<b>Silt</b>	<b>Silt</b>	<b>Silt</b>	<b>Silt</b>	<b>Silt</b>

As it is maintained in table 1, it is appearing that the nature of soils of Southern Iraq marshes directed towards the basicity with a pH range from 7.4 in st. 2 (El-Chebaysh) to 8.1 in st 5 (Hareer), while there are large differences in the electric conductivities EC which range between 4.59 dS/cm in st. 4 (Umm El-Ward) and 13.16 dS/m in st. 6 (El-Barghah). Total nitrogen ranged from 0.413 % in st. 1 (Al-Imarah) to 0.621% in st 3 (Umm El-Naaj), total phosphorus

between 0.85 (µg/g) in st. 2 (El-Gebaysh) to 1.42 µg/g in st. 4 (Umm El-Ward).

The studied marsh soils were rich in chemical elements, Co reported values between 307.804 and 395.748 µg/g in stations 3 or 4 and 5 respectively, for Fe ,the highest values were reported as 7541.895 and 7887.727 µg/g in sts. 2 and 5 respectively, while for Mn the lowest value reported was 721.523 µg/g in st. 1 and the highest value as 884.43 µg/g in st. 2, both within Central

marsh. For Ni, values were alternative, the lowest 170.526  $\mu\text{g/g}$  in st. 3 and the highest 341.052  $\mu\text{g/g}$  in st. 2.

On the other hand for K low values were reported ,the greatest were 6.35 mg / g dry weight in st 6 and 5.62 mg / g dry weight in st 5, both within Al-Hammar marsh. For Na values reported ranged between 512.130 mg / g in st. 6 and 661.310 mg / g in st. 1. Those values for K and Na reported in this study don't agree with what reported earlier as 17ppm for K and 100 ppm for Na in soils from stations nearby st. 5 and 6 of this study (Hussain *et al.*, 1993). The increased of Na and K in the soil could be explosion on the basis of marshes desiccation in which salinity is increased in most of standard sites. Moreover, the standard sites were enriched with aquatic organism before desiccation leading to increased salinity and this results were agreeing with, (Ali et al ,2012)

Texture analysis shown in table 2 revealed that the texture of most studied soils from the southern Iraqi marshes were contrast in type and

site, all showed high percents of clay and low percents of sand, therefore all were classified as soils with a texture of silty soil , accordingly these soils are characterized to be cheap in conservation of nutrients and humidity to be dry very quick and it has good porosity and ventilation , and this results was agree with (Nughamesh and Ali, 2005).

### Conclusion

It is concluded that chemical elements are contra verse in values in the soils of Iraqi marshes, Co and Ni were very low compared to Fe and Mn. More over the texture of the studied soils were mostly silt, therefore it classified as silt texture, and this type of soil is the best to be suitable for agriculture of cereal crops. Finally, we recommend using these sites for agriculture to rise the agriculture field in southern Iraq.

**References:**

- Douabul, Ali, A.Z. ;Nadia , A. Al-Mudhafer ;Abdulzahra ,A. Alhello ; Hamid , T. Al- Saad and Sama ,S. Al-Maarofi (2012). Restoration versus Re-flooding :Mesopotamia Marshland ,Hydrology Current Research , 3:5.
- Hussain, H. H. , Jassim, M. A. and Nasser, T. Ak. (1993). Determination of some trace elements in soil samples from southern Iraq. Basra J. Agric. Sci., 6(1):111-120.
- Kovda, V. A. (1988). A study of nine element contents in soils of Amur river area. Sov. Soil Sci., 12: 1369-1377.
- Mahmood ,A.A.(2008). Concentration of pollutants in water ,sediments and aquatic plants in some wetlands in south of Iraq. Phd ,Uni. Basrah Iraq P:256 .
- Mitchell, R. L. (1972). Cobelt in soil and uptake by plants. Agro Chimica, 21: 121-135.
- Nughames, R. G, and Ali, S. A. (2005). Study of Physico chemical features for marshes water and their sediments and soils within the Governorate of Dhe Qar. Marine Mesopotamica, 20(1):67-80.
- Page, A.L. ;R.H. Miller and Keeney ,D.R.(1982). Methods of soil analysis .Part (2).2nd .ed Madison Wisconson, USA p:1159.
- Savchell, V. (1969). Trace elements in Agriculture. V.H.A.Co., New York, London.
- Sturgeon ,R. E. ;J. A. Desaulnicrs ; S. S. Berman and D.S.Russel (1982). Determination of trace metals in estuarine sediment by graphite furnace atomic absorption spectrophotometer Anal. Chem. Acta .134:288-291.
- Van Wijck ,C.;Comelis-Jan De Groot ,C.(1993) . The impact of desiccation of a freshwater marsh (Garcines Nord, Camargue ,France ) on sediment – water –vegetation interaction part 2:Hydobiologia 252 :95-103.
- Venogradov, G. P. (1959). The geochemistry of rare and dispressed chemical elements in soil. Consultants Bureau, New York, 209pp.

## بعض الخصائص الكيميائية لترب أهوار جنوب العراق

\* فارس جاسم محمد الامارة و \*\* حليلة جبار عبد الرزاق العرادي

\* قسم الكيمياء وتلوث البيئة البحرية, مركز علوم البحار

\*\* قسم التطور الإحيائي في شط العرب وشمال الخليج العربي مركز علوم البحار /

جامعة البصرة, جامعة البصرة, البصرة - العراق

### المستخلص

اجريت دراسة لتقييم بعض مواصفات ترب من ستة مواقع موزعة بين اهوار العراق الجنوبية وهي هور الامارة والجبايش ضمن الهور الوسطي وام النعاج و ام الورد ضمن هور الحويزة و هور حرير و البركة ضمن هور الحمار وذلك لتحديد نوعيتها ومدى ملائمتها لزراعة انواع مختلفة من المحاصيل الزراعية. تم جمع عينات بعمق 10- 90 سم من التربة ونقلت الى المختبر في مركز علوم البحار لقياس المواصفات التالية: الاس الهيدروجيني والتوصيلية الكهربائية و تقدير العناصر النزرة التالية : الكوبلت والحديد والمنغنيز والنيكل, وكذلك تقدير النسبة الكلية للنتروجين والفسفور الذائب والصوديوم والبوتاسيوم فضلاً عن تقييم نسجة التربة بتحديد النسب المئوية لكل من الرمل والغرين والطين لتحديد نوع التربة. بينت النتائج ان الاس الهيدروجيني للتربة المدروسة تراوح بين ادنى قيمة 7.4 في الجبايش و اعلى قيمة 8.1 في هور حرير , وتراوحت التوصيلية الكهربائية بين ادنى قيمة 4.59 الى 13.16 مللي سيمنس/ سم في ام الورد والبركة على التوالي. وكانت تراكيز المعادن النزرة المدروسة متفاوتة بين محطات الدراسة وكان ادناها 170.526 مايكروغم/غرام وزن جاف نيكل في تربة ام النعاج و اعلاها 7887.727 مايكروغم/غرام وزن جاف حديد في تربة هور حرير وكانت اعلى التربة اغناءً بالنتروجين هي 0.621 % في ام النعاج والفسفور 1.42 مايكروغرام/ غم في ام الورد وادناها بالنتروجين 0.420 % في هور الامارة والفسفور 0.85 مايكروغرام/غم في الجبايش . وتراوحت قيم البوتاسيوم والصوديوم بحدود 4.21 – 6.35 جزء بالمليون و 512.13 – 661.31 جزء بالمليون في هور الامارة على التوالي. يستخلص من هذه الدراسة ان معظم الارض في الاهوار صالحة لزراعة انواع مختلفة من المحاصيل الزراعية .