

العلاقة بين انواع دوال الهوية

رنا بهجت ياسين

رغد وميض فارس

اسيل علاء عوض علي

قسم الرياضيات – كلية التربية للبنات – جامعة تكريت

The relationship between types of identifications

Rana B.t Yaseen

Raghad W. Faris

Assel A. Awad

Dep. of mathematics-College of education for women University of Tikrit.

في هذا البحث تستخدم تعاريف المجموعات المفتوحة من انماط

(α – open , pre – open , b – open , β – open)

لتحديد تعاريف جديده لدوال الهوية في الفضاءات التوبولوجية، اسميناها

α – identification, pre – identification, b – identification , β – identification

وناقشنا العلاقة فيما بينهم . وايضا "بعض صفات تلك الدوال تُرست وبُرهننت .

الدالة المفتاحية :

α – identification, pre – identification, b – identification , β – identification

Abstract

In this paper , used the definitions of (α – open , pre – open , b – open , β – open) sets in order to limit the identifications in topological space namely (α – identification, pre – identification , b – identification , β – identification) functions and we discuss the relationship between them , as well as several properties of these functions are proved.

Keyword :

α – identification, pre – identification, b – identification

and β – identification

Introduction and Preliminaries:

The concept of continuous(α –continuous, pre –continuous , b –continuous, β –continuous) function, irresolute(α – irresolute , pre – irresolute , b – irresolute , β – irresolute)function and contra – continuous(contra – α – continuous, contra pre – continuous , contra – b – continuous , contra – β – continuous) have been introduced and investigated by Mashhour [12 ,13],Andrjevic [3] ,El-Monsef [5],(Maheshwair and Thakur) [10] , (Jafaris and Noiri) [7, 8] and Calda [4] respectively. By using" semi-, (α – , pre – , β – , b–) open sets " have been introduced and investigated by Levine [9],Njasted [18] , Mashhour [12,13] , Andrjevic [3], El-Monsef [5] respectively.

AL-kutabi [1] in 1996 , introduces and studies some weak identifications , the notion of semi-identification, Mazl [14] introduces the notion of b- identification. In this work , we study the concepts of types of identifications and discuss the relation between them .Also, we investigate it's relationship with other types of identifications.

" Throughout this paper \mathcal{H} , \mathcal{M} and \mathcal{X} ,will denote topological spaces for a subset \mathcal{A} of space $(\mathcal{H}, \mathcal{T})$, $\text{int}(\mathcal{A})$, $\text{cl}(\mathcal{A})$, denoted the interior and closure of a set \mathcal{A} , respectively ", and we indicate them by the following symbols : $\text{gof} = \mathcal{W}$, $f^{-1} = \mathcal{S}$, $g^{-1} = \mathcal{h}$, $f(f^{-1}) = \mathcal{F}$.

" A subset \mathcal{A} of a space \mathcal{H} is said to be:

1. α –open set [18](for short \mathcal{D} –) if $\mathcal{A} \subseteq \text{int}(\text{cl}(\text{int}(\mathcal{A})))$. So \mathcal{A}^c called α – closed (for short $\mathcal{D} =$).
2. pre –open set [12] (for short p–) if $\mathcal{A} \subseteq \text{int}(\text{cl}(\mathcal{A}))$. So \mathcal{A}^c called pre – closed (for short p =).
3. β –Open set [5] (for short \mathcal{B} –) if $\mathcal{A} \subseteq \text{cl}(\text{int}(\text{cl}(\mathcal{A})))$. So \mathcal{A}^c called β – closed (for short $\mathcal{B} =$).
4. b –open set [3] (for short b –) if $\mathcal{A} \subseteq (\text{cl}(\text{int}(\mathcal{A})) \cup \text{int}(\text{cl}(\mathcal{A})))$. So \mathcal{A}^c called b – closed (for short b =)." .

The family of all $(\mathcal{D}-, p-, \mathcal{B}-, b-)$ sets is denoted by $\mathcal{D}O(\mathcal{H}), pO(\mathcal{H}), \mathcal{B}O(\mathcal{H}), bO(\mathcal{H})$.

Remark : the diagram below shows the relationship between open sets .

$$\text{open} \rightarrow \mathcal{D}- \rightarrow p- \rightarrow b- \rightarrow \mathcal{B}-$$

figure (1)

" The converse of these implications are not true in general".

Example 1 :

Let $\mathcal{H} = \{d, k, p, O, C\}$ on $\mathfrak{S} = \{\mathcal{H}, \varphi, \{p, O\}, \{d, k\}, \{d, k, p, O\}\}$.

Then

- A subset $\{d\}$ of \mathcal{H} is $p-$ but it does not $\mathcal{D}-$.
- A subset $\{d, k, C\}$ of \mathcal{H} is $b-$ but it does not $p-$.
- A subset $\{p, C\}$ of \mathcal{H} is $\mathcal{B}-$ but it does not $b-$.

"The following definitions and results were introduced and studied".

Definition 2: "Let a function of a space \mathcal{H} into a space \mathcal{M} then:

- 1- f is called open (closed) function if the image of each open (closed) set in \mathcal{H} is open(closed) set in \mathcal{M} [6].
- 2- f is called $\mathcal{D}-$ ($\mathcal{D}=\$) function if the image of each $\alpha-$ open ($\mathcal{D}=\$) set in \mathcal{H} is $\mathcal{D}-$ ($\mathcal{D}=\$) set in \mathcal{M} [13].
- 3- f is called $p-$ ($p=\$) function if the image of each $p-$ ($p=\$) set in \mathcal{H} is $p-$ ($p=\$) set in \mathcal{M} [12].
- 4- f is called $b-$ ($b=\$) function if the image of each $b-$ ($b=\$) set in \mathcal{H} is $b-$ ($b=\$) set in \mathcal{M} [3].
- 5- f is called $\mathcal{B}-$ ($\mathcal{B}=\$) function if the image of each $\mathcal{B}-$ ($\mathcal{B}=\$) set in \mathcal{H} is $\mathcal{B}-$ ($\mathcal{B}=\$) set in \mathcal{M} [5]. "

Remark : the diagram below holds for a functions .

$$\text{open fun.} \rightarrow \mathcal{D}- \text{ fun.} \rightarrow p- \text{ fun.} \rightarrow b- \text{ fun.} \rightarrow \mathcal{B}- \text{ fun.}$$

figure (2)

"Now by [3,5,12,13]and the following examples illustrate that The converse of these implication are not true in general" .

Definition 3 : A function $f : \mathcal{H} \rightarrow \mathcal{M}$ is called:

- 1- A continuous function if \mathfrak{S} of any open set in \mathcal{M} is a open set in \mathcal{H} [6].
- 2- $\alpha-$ continuous function if \mathfrak{S} of any open set in \mathcal{M} is $\mathcal{D}-$ set in \mathcal{H} [13].
- 3- pre -continuous function if \mathfrak{S} of any open set in \mathcal{M} is $p-$ set in \mathcal{H} [12].
- 4- b -continuous function if \mathfrak{S} of any open set in \mathcal{M} is $b-$ in \mathcal{H} [2].
- 5- $\beta-$ continuous function if \mathfrak{S} of any open set in \mathcal{M} is $\mathcal{B}-$ set in \mathcal{H} [5].

Remark : Mubarki in 2013 presented the following diagram that illustrates the relationship between the types of continuous functions . [15]

$$\text{cont.} \rightarrow \alpha- \text{ cont.} \rightarrow \text{pre- cont.} \rightarrow b- \text{ cont.} \rightarrow \beta- \text{ cont.}$$

figure (3)

"The converse of these implications are not true in general and the following examples".

Example. 4:

Let $\mathcal{H} = \{d, k, p, O, C\}$ on $\mathfrak{S} = \{\mathcal{H}, \varphi, \{p, d\}, \{d, k\}, \{d, k, p, d\}\}$

1-Then , $f : \mathcal{H} \rightarrow \mathcal{H}$ defined by $f(d) = k, f(k) = d, f(p) = p, f(O) = k, f(C) = C$, is pre -continuous function but it is not $\alpha-$ cont.

2- Then , $f : \mathcal{H} \rightarrow \mathcal{H}$ defined by $f(d) = d, f(k) = k, f(p) = p, f(O) = O, f(C) = k$, is b -cont. but it is not pre - cont.

3- Then, $f : \mathcal{H} \rightarrow \mathcal{H}$ defined by $f(d) = p, f(k) = c, f(p) = d, f(o) = o, f(c) = k$, is β -cont. but it is not b -cont.

Definition 5 :

A mapping $f : \mathcal{H} \rightarrow \mathcal{M}$ is called irresolute function [10] (resp. α -irresolute [10], pre-irresolute [13], b -irresolute [3] β -irresolute [5]) if $\mathfrak{H}(u)$ is open ($\mathcal{D}-, p-, b-, \mathfrak{B}-$) in \mathcal{H} for each open ($\mathcal{D}-, p-, b-, \mathfrak{B}-$) in \mathcal{M} .

"Diagram (4)" :

irresol. \rightarrow α -irresol. \rightarrow pre-irresol. \rightarrow b -irresol. \rightarrow β -irresol.

generally speaking, the opposite of the implication is not necessarily true, as follows instance.

Example 6 :

Let $\mathcal{H} = \{d, k, p, o, c\}$ on $\mathfrak{T} = \{\mathcal{H}, \varphi, \{p, o\}, \{d, k\}, \{d, k, p, o\}\}$

1- Then, the $f : \mathcal{H} \rightarrow \mathcal{H}$ defined by $f(d) = d, f(k) = p, f(p) = k, f(o) = o, f(c) = c$, is pre-irresol. and not α -irresol.

2- Then, the $f : \mathcal{H} \rightarrow \mathcal{H}$ defined by $f(d) = d, f(k) = k, f(p) = c, f(o) = o, f(c) = p$, is b -irresol. and not pre-irresol.

3- Then the $f : \mathcal{H} \rightarrow \mathcal{H}$ defined by $f(d) = p, f(k) = k, f(p) = d, f(o) = c, f(c) = o$, is β -irresol. and not b -irresol.

Definition 7 :

A function $f : \mathcal{H} \rightarrow \mathcal{M}$ is called contra-continuous

(resp. contra α -continuous, contra pre-continuous [6,7], contra b -continuous [2] contra β -continuous [4]), if $\mathfrak{H}(u)$ is closed ($\mathcal{D} =, p =, b =, \mathfrak{B} =$) in \mathcal{H} , for each open set u of \mathcal{M} .

"Diagram (5)" :

contra-cont. \rightarrow contra α -cont. \rightarrow contra pre-cont. \rightarrow contra b -cot.
 \rightarrow contra β -cot.

The examples show that the reversal of the chart is incorrect.

Example 8 :

Let $\mathcal{H} = \{d, k, p, o, c\}$ on $\mathfrak{T} = \{\mathcal{H}, \varphi, \{p, o\}, \{d, b\}, \{d, k, p, o\}\}$

1-Then, $f : \mathcal{H} \rightarrow \mathcal{H}$ defined by $f(d) = c, f(k) = k, f(p) = d, f(o) = o, f(c) = p$.

Is contra pre-cont. but it is not contra α -cont.

2- Then, $f : \mathcal{H} \rightarrow \mathcal{H}$ defined by $f(d) = p, f(k) = o, f(p) = d, f(o) = k, f(c) = c$.

Is contra b -cont. but not contra pre-cont.

3-Then, $f : \mathcal{H} \rightarrow \mathcal{H}$ defined by $f(d) = d, f(k) = o, f(p) = p, f(o) = k, f(c) = c$.

Is contra β -cont. but not contra b -cont.

A Study of some new types of identifications:

In this section, we introduce new definitions of (α -identification, pre-identification, b -identification, β -identification) functions by using ($\mathcal{D}-, p-, b-, \mathfrak{B}-$) sets and study the relations between them.

Definition 9 : " A function $f : \mathcal{H} \rightarrow \mathcal{M}$ is called α -identification iff f is onto and one of the following condition satisfies "

--- U is $\mathcal{D}-$ in \mathcal{M} iff $\mathfrak{H}(u)$ is $\mathcal{D}-$ in \mathcal{H} .

--- U is $\mathcal{D} =$ in \mathcal{M} iff $\mathfrak{H}(u)$ is $\mathcal{D} =$ in \mathcal{H} .

For example : Let $\mathcal{H} = \{d, k, p, o\}$ and $\mathcal{M} = \{1, 2, 3\}$ be equipped with the topologies $\mathfrak{T}_{\mathcal{H}} = \{\mathcal{H}, \varphi, \{k, p\}, \{d, k\}, \{k\}\}$, $\mathfrak{T}_{\mathcal{M}} = \{\varphi, \mathcal{M}, \{1, 2\}, \{2, 3\}, \{2\}\}$

If $f : \mathcal{H} \rightarrow \mathcal{M}$ defined by $f(d) = 1, f(k) = 2, f(p) = 3, f(o) = 3$.

we get f is α – identification.

Proposition 10 :

Every α – irresolute and $\mathcal{D} - (\mathcal{D} =)$ onto functions is α – identification.

Proof : A $f : \mathcal{H} \rightarrow \mathcal{M}$, $f(U)$ is $\mathcal{D} -$ since f is onto and $\mathcal{D} -$,

so $(\mathcal{F}(U)) = U$ is $\mathcal{D} -$. and $U \subseteq \mathcal{M}$.

f is α – irresolute hanc $\mathfrak{S}(U)$ is $\mathcal{D} -$ in \mathcal{H} , so f is α – identification.

While if :

If $f : \mathcal{H} \rightarrow \mathcal{M}$ is onto , $\mathcal{D} =$ and α – irresolute ,

Hence $\mathfrak{S}(U)$ is $\mathcal{D} -$, implies $(\mathfrak{S}(U))^c = \mathfrak{S}(U^c)$ is $\mathcal{D} =$, which $(\mathcal{F}(U^c)) = (U^c)$ is $\mathcal{D} =$,

Since f is α – irresolute by def. so f is α – identification. p – in \mathcal{H} ".

Definition 11 :

" A function $f : \mathcal{H} \rightarrow \mathcal{M}$ is called pre – identification if f is onto and U is p – in \mathcal{M} iff $f^{-1}(u)$ is p – in \mathcal{H} ".

Remark : from figure (1) we get every α – identification is pre – identification but the opposite is not true .

As follows instance .

$\mathcal{H} = \{d, k, p, \mathcal{O}, \mathcal{C}\}$, $\mathcal{M} = \{d, k, p, \mathcal{O}\}$ be equipped with topologies

$\mathfrak{S}_x = \{\mathcal{H}, \varphi, \{p, \mathcal{O}\}, \{d, k\}, \{d, k, p, \mathcal{O}\}\}$ and $\mathfrak{S}_M = \{\varphi, Y, \{d, k\}, \{k, p\}, \{k\}\}$.

If $f : \mathcal{H} \rightarrow \mathcal{M}$ defined by $f(d) = d$, $f(k) = k$, $f(p) = p$,

$f(\mathcal{O}) = \mathcal{O}$, $f(\mathcal{C}) = p$, we get f is pre – identification.

Lemma 12 : A onto function $f : \mathcal{H} \rightarrow \mathcal{M}$ is called pre – identification , U is $p =$ in \mathcal{M} iff $\mathfrak{S}(U)$ is $p =$ in \mathcal{H} .

Proposition 13 :

Every pre – irresolute and $p - (p =)$ onto functions is pre – identification

Proof :

from " figure (1 ,4)" every $\mathcal{D} -$ function is p – function and α – irresolute is pre – irresolute , by Proposition 10 , we get every f α – irresolute is pre – irresolute.

Definition 14:

" A function $f : \mathcal{H} \rightarrow \mathcal{M}$ is called b – identification if f is onto and one of the following condition satisfies "

1) U is b – in \mathcal{M} iff $\mathfrak{S}(u)$ is b – in \mathcal{H} .

2) U is $b =$ in \mathcal{M} iff $\mathfrak{S}(u)$ is $b =$ in \mathcal{H} . [3]

"from figure (1) every p – is b – then for each pre – identification is b – identification."

We note from an example 1 :

A $f : \mathcal{H} \rightarrow \mathcal{H}$ be defined by $f(d) = d$, $f(k) = k$, $f(p) = \mathcal{C}$, $f(\mathcal{O}) = \mathcal{O}$, $f(\mathcal{C}) = p$, then f is b – identification but not pre – identification, since $\mathfrak{S}\{d, k, p\} = \{d, k, \mathcal{C}\} \notin \text{PO}(\mathcal{H})$.

Proposition 15 :

If $f : \mathcal{H} \rightarrow \mathcal{M}$ is onto , $b - (b =)$ and b – irresolute then f is b – identification. [3].

Proposition 16 :

The composition of two, α – identification(pre – identification, b – identification) functions is α – identification(pre – identification, b – identification).

Proof :

Suppose that $f : \mathcal{H} \rightarrow \mathcal{M}$, $g : \mathcal{M} \rightarrow \mathcal{N}$ are α – identifications

"Whenever The compo. of two onto functions is onto "

Now, if U be any \mathcal{D} -in \mathfrak{X} , by hypo. $\mathcal{G}, \mathfrak{f}$ are α -identifications then $\mathfrak{h}(U)$ is \mathcal{D} -in \mathcal{M} and we have $\mathfrak{H}(\mathfrak{h}(U)) = (\mathcal{W})^{-1}(U)$ is \mathcal{D} -in \mathcal{H} . implies U is \mathcal{D} -in \mathcal{H} , thus \mathcal{W} is α -identification.

Similarly, to prove \mathcal{W} is (pre-identification, b-identification).

Proposition 17 :

A $\mathfrak{f}: \mathcal{H} \rightarrow \mathcal{M}$ and $\mathcal{G}: \mathcal{M} \rightarrow \mathfrak{X}$ are functions and \mathfrak{f} is α -identification (pre-identification, b-identification)

then the following statement are valid :

- 1- If \mathcal{W} is α -cont. (pre-cont., b-cont.) then \mathcal{G} is α -cont. (pre-cont., b-cont.).
- 2- If \mathcal{W} is α -irresolute (pre-irresolute, b-irresolute.) then \mathcal{G} is α -irresolute. (pre-irresolute, b-irresolute.).
- 3- If \mathcal{W} is contra α -cont. (contra pre-cont., contra b-cont.) then \mathcal{G} is contra α -cont. (contra pre-cont., contra b-cont.).

Proof :

- 1) Let $\mathcal{W}: \mathcal{H} \rightarrow \mathfrak{X}$ is α -cont., Assume that k any an open set in \mathfrak{X} , Let $V = \mathfrak{h}(k)$ and $U = \mathfrak{H}(V)$, whenever $\mathcal{W}^{-1}(k) = \mathfrak{H}(\mathfrak{h}(k)) = U$ is \mathcal{D} -in \mathcal{H} , then $\mathcal{W}^{-1}(k)$ \mathcal{D} -in \mathcal{H} , but \mathfrak{f} is α -identif. then V is \mathcal{D} -in \mathcal{M} . So $\mathfrak{h}(k)$ \mathcal{D} -in \mathcal{M} , so \mathcal{G} is α -cont.
- 2) Assume that k any an \mathcal{D} -set in \mathfrak{X} , Let $V = \mathfrak{h}(k)$ and $U = \mathfrak{H}(V)$, we have $\mathcal{W}^{-1}(k) = \mathfrak{H}(\mathfrak{h}(k)) = U$, that is, U is \mathcal{D} -in \mathcal{H} , we get $\mathcal{W}^{-1}(k)$ \mathcal{D} -in \mathcal{H} , but \mathfrak{f} is α -identif. , then V is \mathcal{D} -in \mathcal{M} . whenever $\mathfrak{h}(k)$ \mathcal{D} -in \mathcal{M} . So \mathcal{G} is α -irresolute.
- 3) Assume that k any an \mathcal{D} -set in \mathfrak{Z} , Let $V = \mathfrak{h}(k)$ and $U = \mathfrak{H}(V)$, we have $\mathcal{W}^{-1}(k) = \mathfrak{H}(\mathfrak{h}(k)) = U$ is \mathcal{D} -in \mathcal{H} , then $\mathcal{W}^{-1}(k)$ \mathcal{D} -in \mathcal{H} , but \mathfrak{f} is α -identif. , then $V = \mathfrak{h}(k)$ is \mathcal{D} -in \mathcal{M} , thus \mathcal{G} is contra α -cont.

Definition 18 :

" A function $\mathfrak{f}: \mathcal{H} \rightarrow \mathcal{M}$ is called β -identification if \mathfrak{f} is onto and U is \mathfrak{B} -in \mathcal{M} iff $\mathfrak{H}(u)$ is \mathfrak{B} -in \mathcal{H} ".

" from figure (1) we get every b-identification is β -identification but the converse is not true". From instance 1: let $\mathfrak{f}: \mathcal{H} \rightarrow \mathcal{H}$ be defined by

$$\mathfrak{f}(d) = d, \mathfrak{f}(k) = k, \mathfrak{f}(p) = \mathcal{C}, \mathfrak{f}(0) = 0, \mathfrak{f}(\mathcal{C}) = p$$

then \mathfrak{f} is b-identification but not pre-identification, since $\mathfrak{H}\{d, k, p\} = \{d, k, \mathcal{C}\} \notin \mathfrak{p}0(\mathcal{H})$.

Proposition 19:

A onto function $\mathfrak{f}: \mathcal{H} \rightarrow \mathcal{M}$ is called β -identification if U is \mathfrak{B} -in \mathcal{M} iff $\mathfrak{H}(U)$ is \mathfrak{B} -in \mathcal{H} .

Proof : If U subset of \mathcal{M} , \mathfrak{B} then U^c is \mathfrak{B} -in \mathcal{M} , since \mathfrak{f} is β -identification, so $\mathfrak{H}(U)$ is \mathfrak{B} -in \mathcal{H} , (by def. \mathfrak{f} is onto, $(\mathfrak{H}(U))^c = \mathfrak{H}(U^c)$ is \mathfrak{B} -in \mathcal{H} . Similarly, if $\mathfrak{H}(U)$ is \mathfrak{B} -in \mathcal{H} , we get $\mathfrak{H}(U)^c = \mathfrak{H}(U^c)$ is \mathfrak{B} -in \mathcal{H} and \mathfrak{f} is β -identif., we get U is \mathfrak{B} -in \mathcal{M} .

Assume that U be \mathfrak{B} -in \mathcal{M} then U^c is \mathfrak{B} -in \mathcal{M} , whenever $(\mathfrak{H}(U))^c = \mathfrak{H}(U^c)$ is \mathfrak{B} -in \mathcal{H} , so $\mathfrak{H}(U)$ is \mathfrak{B} -in \mathcal{H} . Similarly,

if $\mathfrak{H}(U)$ is \mathfrak{B} -in \mathcal{H} , we get $(\mathfrak{H}(U))^c = \mathfrak{H}(U^c)$ is \mathfrak{B} -in \mathcal{H} ,

and then U^c is $\mathfrak{B} =$, so U is $\mathfrak{B} -$.

proposition 20 :

If $f : \mathcal{H} \rightarrow \mathcal{M}$ is onto , $\mathfrak{B} -$ ($\mathfrak{B} =$) and $\beta -$ irresolute then f is $\beta -$ identification.

Proof :

Assume that U is $\mathfrak{B} =$ in \mathcal{H} , $U \subseteq \mathcal{M}$, such that $\mathfrak{H}(U)$ is $\mathfrak{B} =$ in \mathcal{H} . whenever $(\mathcal{F}(U)) = U$, we get U is $\mathfrak{B} =$ in \mathcal{H} (Since $\mathfrak{H}(U)$ is $\mathfrak{B} =$ in \mathcal{H} , ,and f is $\mathfrak{B} =$ in \mathcal{H}). ,so U^c is $\mathfrak{B} -$ in \mathcal{H} , and since f is $\beta -$ irresolute then $\mathfrak{H}(U)$ is $\mathfrak{B} -$ in \mathcal{H} , whenever f is onto $(\mathfrak{H}(U))^c = \mathfrak{H}(U^c)$ implies $\mathfrak{H}(U)$ is $\mathfrak{B} -$ in \mathcal{H} , thus ,by Proposition 19 , then f is $\beta -$ identification.

Theorem 21 : The below stated expressions are hold .

- 1- every identification is $\alpha -$ identification.
- 2- every $\alpha -$ identification is pre - identification.
- 3- every pre - identification is $\mathfrak{b} -$ identification.
- 4- every $\mathfrak{b} -$ identification is $\beta -$ identification.

Proof : obvious.

Remark : " the above examples show that the inverse theorem is not necessarily true ."

Proposition 22 :

"The composition of two $\beta -$ identification functions is $\beta -$ identification".

Proof:

Let $f : \mathcal{H} \rightarrow \mathcal{M}$, $g : \mathcal{M} \rightarrow \mathcal{N}$ are $\beta -$ identifications

"Whenever The compo. of two onto functions is onto" .

,If U be any $\mathfrak{B} -$ in \mathcal{N} , by hypo. g, f are $\beta -$ identifications then $h(U)$ is $\mathfrak{B} -$ in \mathcal{M} and we have $\mathfrak{H}(h(U)) = (\mathcal{W})^{-1}(U)$ is $\mathfrak{B} -$ in \mathcal{H} , implies U is $\mathfrak{B} -$ in \mathcal{H} , thus \mathcal{W} is $\beta -$ identification.

Proposition 23 :

$f : \mathcal{H} \rightarrow \mathcal{M}$, $g : \mathcal{M} \rightarrow \mathcal{N}$ be functions and f is $\beta -$ identification then the following statement are valid :

- 1- If \mathcal{W} is $\beta -$ cont. then g is $\beta -$ cont.
- 2- If \mathcal{W} is $\beta -$ irresolute then g is $\beta -$ irresolute .
- 3- If \mathcal{W} is contra $\beta -$ cont. then g is contra $\beta -$ cont.

Proof :

1) Let $\mathcal{W} f : \mathcal{H} \rightarrow \mathcal{N}$ is $\beta -$ cont. , Assume that k any an open set in \mathcal{N} , Let $V = h(k)$ and $U = \mathfrak{H}(V)$, we have $W^{-1}(k) = \mathfrak{H}(h(k)) = U$ is $\mathfrak{B} -$ in \mathcal{H} , then $W^{-1}(k)$ $\mathfrak{B} -$ in \mathcal{H} ,but f is $\beta -$ identification , then V is $\mathfrak{B} -$ in \mathcal{M} .So $h(k)$ $\mathfrak{B} -$ in \mathcal{M} , thus g is $\beta -$ cont.

2) Assume that k any an $\mathfrak{B} -$ set in \mathcal{N} , Let $V = h(k)$ and $U = \mathfrak{H}(V)$, we have $W^{-1}(k) = \mathfrak{H}(h(k)) = U$,that is, U $\mathfrak{B} -$ in \mathcal{H} , we get $W^{-1}(k)$ $\mathfrak{B} -$ in \mathcal{H} ,but f is $\beta -$ identif. , then V is $\mathfrak{B} -$ in \mathcal{M} , thus g is $\beta -$ irresolute .

3) Assume that k any an $\mathfrak{B} -$ set in \mathcal{N} , Let $V = h(k)$ and $U = \mathfrak{H}(V)$, we have $W^{-1}(k) = \mathfrak{H}(h(k)) = U$ is $\mathfrak{B} -$ in \mathcal{H} . So $W^{-1}(k)$ $\mathfrak{B} -$ in \mathcal{H} ,but f is $\beta -$ identifi. , then $V = h$.

Remark : from the above discussion and known results we have the following implications .

identification \rightarrow α – identification \rightarrow pre – identification \rightarrow b – identification
 \rightarrow β – identification

figure (6)

Definition 24 : " A space $(\mathcal{H}, \mathfrak{S})$ is said to be $\alpha - \mathfrak{S}_1$ (pre- \mathfrak{S}_1 , b- \mathfrak{S}_1 , $\beta - \mathfrak{S}_1$) [8,11,16, 18] iff for each a pair of distinct points $x, y \in \mathcal{H}$, each belongs to an $\mathfrak{D} - (\mathfrak{p} - \mathfrak{b}, \mathfrak{B} - \mathfrak{B})$ sets which does not contain the other .

Theorem 25 : A function $f : \mathcal{H} \rightarrow \mathcal{M}$ is α – identification and \mathcal{M} is $\alpha - \mathfrak{S}_1$, then \mathcal{H} is $\alpha - \mathfrak{S}_1$.

Proof : let $x, y \in \mathcal{H}$, $x \neq y$, since \mathcal{M} is $\alpha - \mathfrak{S}_1$, there exist $\mathfrak{D} -$ sets M_1 and M_2 , of \mathcal{M} such that $f(x) \in M_1$ and $f(y) \in M_2$, $f(y) \notin M_1$ and $f(x) \notin M_2$.

Since function $f : \mathcal{H} \rightarrow \mathcal{M}$ is α – identification , we have

$$x \in \mathfrak{S}(M_1), y \in \mathfrak{S}(M_2) \text{ and } x \notin \mathfrak{S}(M_2), y \notin \mathfrak{S}(M_1)$$

hence then \mathcal{H} is $\alpha - \mathfrak{S}_1$.

Theorem 26 : A function $f : \mathcal{H} \rightarrow \mathcal{M}$ is pre – identification and \mathcal{M} is pre- \mathfrak{S}_1 , then \mathcal{H} is pre – \mathfrak{S}_1 .

Proof : let $x, y \in \mathcal{H}$, $x \neq y$, since \mathcal{M} is pre – \mathfrak{S}_1 , there exist $\mathfrak{p} -$ sets M_1 and M_2 , of \mathcal{M} such that $f(x) \in M_1$ and $f(y) \in M_2$, $f(y) \notin M_1$ and $f(x) \notin M_2$. Since function $f : \mathcal{H} \rightarrow \mathcal{M}$ is pre – identification , we have $x \in \mathfrak{S}(M_1), y \in \mathfrak{S}(M_2)$ and $x \notin \mathfrak{S}(M_2), y \notin \mathfrak{S}(M_1)$ hence then \mathcal{H} is pre – \mathfrak{S}_1 .

Theorem 27: A function $f : \mathcal{H} \rightarrow \mathcal{M}$ is b – identification and \mathcal{M} is b – \mathfrak{S}_1 , then \mathcal{H} is b – \mathfrak{S}_1 .

Proof : let $x, y \in \mathcal{H}$, $x \neq y$, since \mathcal{M} is b – \mathfrak{S}_1 , there exist b – sets M_1 and M_2 , of \mathcal{M} such that $f(x) \in M_1$ and $f(y) \in M_2$, $f(y) \notin M_1$ and $f(x) \notin M_2$. Since function $f : \mathcal{H} \rightarrow \mathcal{M}$ is b – identification ,

we have $x \in \mathfrak{S}(M_1), y \in \mathfrak{S}(M_2)$ and $x \notin \mathfrak{S}(M_2), y \notin \mathfrak{S}(M_1)$

hence then \mathcal{H} is b – \mathfrak{S}_1 .

Theorem 28 : A function $f : \mathcal{H} \rightarrow \mathcal{M}$ is β – identification and \mathcal{M} is $\beta - \mathfrak{S}_1$ then \mathcal{H} is $\beta - \mathfrak{S}_1$.

Proof : let $x, y \in \mathcal{H}$, $x \neq y$, since \mathcal{M} is $\beta - \mathfrak{S}_1$, there exist $\mathfrak{B} -$ sets M_1 and M_2 , of \mathcal{M} such that $f(x) \in M_1$ and $f(y) \in M_2$, $f(y) \notin M_1$ and $f(x) \notin M_2$. Since function $f : \mathcal{H} \rightarrow \mathcal{M}$ is $\beta -$ identification , we have $x \in \mathfrak{S}(M_1), y \in \mathfrak{S}(M_2)$ and $x \notin \mathfrak{S}(M_2), y \notin \mathfrak{S}(M_1)$

hence then \mathcal{H} is $\beta - \mathfrak{S}_1$.

References:

[1] Al-kutaibi, S.H., "on some types of identification" Dep. Of mathematics university of Tikreet . Tikreet univ. J.Sci. Vol 4.No.3. pp.49-59.(1998) .

[2] Al-omari, A. and Norrani, M.S., "some properties of contra - b –continuous and almost contra - b –continuous functions" Men.iac.sci.Kochi. univ. 22pp. 19-28 (2001).

[3] Andrijevic, D. "On b- open sets ", Mat. Bech., 48 ,pp.59-64 (1996).

[4] Caldas, M., Jafari, S. "some properties of contra - β -continuous functions " Men.iac.sci.Kochi. univ. 22,pp. 19-28 (2001).

[5] El-Monsef, M.E. and elat " β – open sets and β – Continuous mappings " Bull. Fac. Sci. Assiut. univ. Vol. 12, pp. 77-90 ,(1983).

[6] Jafari, S., Noiri, T. " contra - Continuous functions between topological spaces " Iranian-Int. J.Sci., Volume 2, pp. 153-167,(2001).

[7] Jafari, S., Noiri, T. " on contra -pre Continuous functions " Bull. Malaysian.Math.Sci.Soc. Volume 25, pp. 115-128,(2002).

- [8] Kar,A., and Bhattacharyya,P., " Some weak separation axioms", Bull. Calcutta Math. Soc.Vol. 82,PP. 415-422, (1990).
- [9]Levine, N. "Semi-open sets and semi - continuity in topological spaces", Amer. Math. Monthly , vol. 70, Pp.36-41,(1963).
- [10] Maheshwair,S.N.,and Thakur ,S.S., " α -irresolute Mapping " Tamkang J. Math. Vol. 11, pp.209 -214,(1980).
- [11] Mahmoud , R.A .,and El-Monsef M.E., " β -irresolute and β -topological invariant" Proc. Pakistan Acad. Sci., Vol. 27 ,pp. 285–296 (1990).
- [12] Mashhour,A.S., Abd El-Monsef ,M.E. and El-Deeb,S.N., "On precontinuous and weak precontinuous mappings "Pros.Math. Science Egypt, Vol.53,pp.47-53,(1982).
- [13] Mashhour,A.S., Hasonein,I.A., and El-Deeb,S.N., " α -Continuous and α -open Mapping)Acta. Math. Hunger Vol. 41, pp.213-218,(1983).
- [14] Mez'l, A.K. , "On b-identification" Journal of Kufa for Mathematics and Computer vol. 1,NO.3,pp.76-80,(2011).
- [15] Mubarki ,A.M.,Al-rshudi ,M.,Al-juhani, M.A., "some properties of contra - β^* – open sets and β^* –continuity in topological spaces " Journal of Tabiah unive. For sci. Vol.8 ,pp. 142-148 (2014).
- [16] Mustafa J.M., "Some separation axioms by b- open sets", Mu'tah Lil- Buhuth wad-Dirasat, (20) (3), 57-63 (2005).
- [17] Njastad,O. "On some classes of nearly open sets"Pacific.J.Math.,15, pp. 961-970(1965).
- [18] Raman,T.C.,Vidyottama K., K. Sharma, " α - Generalized & α - Separation Axioms for Topological Spaces" IOSR Journal of Math. Vol. 10, Issue 3 Ver. VI., PP 32-36 (2014).