

Theoretical Study of Charge Transfer in Styryl Thiazilo Quinoxaline Dyes STQ-1, STQ-2, and STQ-3 in Organic Media System

*Hadi.J.Mujbel. AL-agealy **
*Mohsin.A.Hassooni**

*Khaled.H.Harbbi**
*Rafah.I.Noori**

Received 23, November, 2011

Accepted 7, May, 2012

Abstract:

Charge transfer in styryl dyes STQ-1, STQ-2, and STQ-3 with organic media system has been studied theoretically depending on the Franck-Condon rule and continuum dielectric model. The reorientation energies λ (eV) were evaluated theoretically depending on dipole momentum, dielectric constant ϵ , and refractive index n . The rate constant of charge transfer has been calculated depending on the reorientation energy λ (eV), effective free energy ΔG^* , potential height barrier ΔG^\ddagger , and coupling coefficient V_{el} . A Matlab program has been written to calculate the rate constant of charge transfer and other parameters. The results of calculations show that STQ-2 dye is more reactive for charge transfer compared with STQ-1 and STQ-3 dyes.

Key words: Theoretical study, Charge transfer, STQ-1, STQ-2, and STQ-3 dye.

Introduction:

Charge transfer (CT) reactions are abundant in chemistry and biology [1]. Studies on charge transfer in organic molecular systems have witnessed rapid growth in recent years [2]. Understanding electron transfer at the molecular level will aid in developing molecular assemblies with unique properties and novel applications, such as molecular electronics devices [3]. The complete theory of charge transfer based on early work of Rudolf Marcus. A great amount of effort has been made in the studies of the electron transfer reaction with a variety of tools such as time-resolved spectroscopy, computer simulation methods, and analytical theory [4]. Theoretical treatments of charge transfer within a solution have focused on charge

transfer between weakly coupled charge transfer sites [5]. For electron transfer in solution, we most commonly consider electron transfer to proceed along a solvent rearrangement coordinate in which solvent reorganizes its configuration so that dipoles overcharge help to stabilize the extra negative charge at the acceptor site [6]. The study of charge transfer in organic media systems has become a very active area in research in recent years. In our study, the styryl dyes with charge donor-acceptor transfer on either side of the styryl bond are particularly attractive for optical and electronic properties. Figure (1) shows the structure of three dyes STQ-1, STQ-2, and STQ-3 used in this study [7].

*Baghdad University/College of Education Ibn Al-Haitham/Department of physics.




Fig. (1). Molecular structures of STQ-1, STQ-2 and STQ-3 dyes[7]

Theory

For a simple linear reactions , the reaction rate constant k_{LRC} , can be written with Franck-Condon rule[8].

$$k_{LRC} = \frac{\rho w_s}{2\pi} \exp\left(-\frac{\Delta G^\ddagger}{k_B T}\right) \dots (1)$$

Where w_s is the frequency of motion in the reactant potential well , ρ is the electronic transition probability at the transition region, ΔG^\ddagger is the activation free energy , and $k_B T$ is the thermal energy.

The probability ρ is a function of electronic matrix element V_{el} in the case of charge transfer. V_{el} is approximately proportional to an overlap integral between the wave functions of the reactant and product. When V_{el} is small, the free energy becomes non adiabatic case [8], in this case the charge transfer rate constant can be written as [8-11].

$$k_{CT} = \left(\frac{\pi}{h^2 k_B T \lambda}\right)^{\frac{1}{2}} |V_{el}|^2 \exp\left(-\frac{\Delta G^\ddagger}{k_B T}\right) \dots (2)$$

Where λ is the solvent reorientation energies due to solvent and anion rearrangement in the secondary coordination sphere.

The reorientation energies can be represented by expression [12].

$$\lambda = \frac{\mu_e^2}{4\pi\epsilon_s D^3} [f(\epsilon_s) - f(n^2)] \dots (3)$$

Where

$$f(\epsilon_s) = \frac{\epsilon_s - 1}{2\epsilon_s + 1} \dots (4)$$

$$\text{And } f(n^2) = \frac{n^2 - 1}{2n^2 + 1} \dots (5)$$

Here ϵ_s is a dielectric constant for varies solvent , n is refractive index, D is the Onsager cavity radius for the dye molecule and μ_e is the dipole moment of charge transfer in excited state effectively.

The effective free energy for reaction charge transfer that broken into the work it takes to bring the donor and acceptor together and the difference between the reduction potentials and determined by [13].

$$\Delta G^* = e[E_{ox} - E_{red}] - \lambda - \Delta E_{00} \dots (6)$$

Where e is the electron charge, E_{ox} , and E_{red} are the oxidation and reduction potentials respectively, and E_{00} is the energy of the ground state.

The driving force ΔG_0 for the charge recombination event can be calculated as [14].

$$h\nu = \Delta G_0 + \lambda \dots(7)$$

where h is the planks constant, and ν is the frequency of the absorption light to excited molecule.

The height barrier of charge transfer is the energy must have the electron transfer is given by [15].

$$\Delta G^\# = \frac{(\Delta G_0 + \lambda)^2}{4\lambda} = \frac{(h\nu)^2}{4\lambda} \dots (8)$$

Results:

The charge transport in some styryl thiozolo quinoxaline STQ-1, STQ-2, and STQ-3 dyes in organic media have been studied here. The charge transport rate constant is determined with many parameters; the reorientation energies $\lambda(eV)$, effective free energy $\Delta G^*(eV)$,

activation free energy $\Delta G^\#(eV)$, and the coupling coefficient matrix element $V_{el}(eV)$,

The reorientation energy λ of STQ-1, STQ-2, and STQ-3 were calculated using a more general expression equation(3) in several organic solvents, Cyclohexane, Diexane, Butanol, Ethanol, and Methanol, respectively. Equation (3) was applied where the values

of activity radius $D \cong 2A^\circ$ for STQ dyes, and the dipole moment of charge transfer μ_e taken from table (1) with ϵ and n are the static dielectric constant and refractive index for solvent from table (2). Results are summarized in the table(3).

Table (1) Molecular and photophysical properties of STQ-1, STQ-2, and STQ-3 styryl thiozolo dyes in organic media

Dye	Molecular weight	Melting point	λ_{ab} (nm)	λ_{em} (nm)	μ_e (Debye)	E_{ox}	E_{red}
STQ-1	360.4	230	490	630	7.43	0.998	-0.907
STQ-2	390.5	156	485	625	10.51	0.993	-0.927
STQ-3	384.5	205	530	623	5.22	0.933	-1.027

Table(2): Organic solvent properties. [16]

solvent	Refractive index (n) [16]	Dielectric constant ϵ [16]
Cyclohexane	1.424	2.01
Dioxane	1.420	2.21
Butanol	1.397	17.51
Ethanol	1.359	23.40
Methanol	1.326	32.63

Table (3) The result of calculation of the reorganization energy $\lambda(eV)$ for charge transfer for styryl thiozolo STQ-1, STQ-2, and STQ-3 dyes in organic media

solvent	STQ-1 Dye	STQ-2 Dye	STQ-3 Dye
Cyclohexane	0.3925	0.7853	0.1777
Dioxane	0.4911	0.9826	0.2224
Butanol	1.5254	3.0521	0.6908
Ethanol	1.6046	3.2106	0.7266
Methanol	1.6737	3.3488	0.7579

So the other parameters in the charge transport is the effective free energy that is provided by is the absorption of light and defined as the part of the work it takes to bring part of system

together, and the difference between the reduction potential of the system. The theoretical calculation values of the effective free energy $\Delta G^*(eV)$ can be estimated relative E_{oo} using

equation (6), where E_{ox} , and E_{red} are the oxidation and reduction potentials respectively are taken from

table (1), and λ from table (3). These values of estimation are shown in table (4).

Table (4) The result of charge transfer effective free energy ΔG° for styryl thiazol STQ-1, STQ-2, and STQ-3 dyes in organic media.

solvent	ΔG° (eV)STQ-1 Dye	ΔG° (eV)STQ-2 Dye	ΔG° (eV) STQ-3 Dye
Cyclohexane	1.5125	1.1346	1.7823
Dioxane	1.4139	0.9373	1.7376
Butanol	0.3796	-1.1321	1.2692
Ethanol	0.3004	-1.2906	1.2333
Methanol	0.2313	-1.4288	1.2021

The activation free energy $\Delta G^{\#}$ is the scale of the height of the barrier depends on absorption of spectrum energy and reorientation energy. The

activation free energy has been calculated using equation (8), results of $\Delta G^{\#}$ using various solvent with STQ dyes are listed in table (5).

Table (5): the activation free energy $\Delta G^{\#}$ for STQ-1, STQ-2, and STQ-3 in organic media

Wave length(nm)	$\Delta G^{\#}$ for STQ-1					$\Delta G^{\#}$ for STQ-2					$\Delta G^{\#}$ for STQ-3				
	500	550	600	650	700	500	550	600	650	700	500	550	600	650	700
Cyclohexane	3.901	3.224	2.709	2.308	1.990	1.950	1.611	1.354	1.153	0.994	8.07	7.122	5.984	5.099	4.396
Dioxane	3.118	2.577	2.165	1.845	1.590	1.558	1.288	1.082	0.922	0.795	6.885	5.690	4.781	4.074	3.512
Butanol	1.003	0.829	0.697	0.594	0.512	0.501	0.414	0.348	0.296	0.255	2.216	1.832	1.539	1.311	1.130
Ethanol	0.954	0.788	0.662	0.564	0.486	0.476	0.394	0.331	0.282	0.243	2.107	1.714	1.463	1.247	1.075
Methanol	0.914	0.756	0.635	0.541	0.466	0.457	0.377	0.317	0.270	0.233	2.02	1.669	1.403	1.195	1.030

After estimate the values of the reorientation energy λ , free energy (driving energy) ΔG° , effective free energy ΔG^* , activation free energy $\Delta G^{\#}$, and assume the electronic coupling coefficient $V_{el} \sim 0.5 eV$ that a key factors controlling of the charge

transfer then parameters and expression equation (2), we can calculated the rate constant of charge transfer, the results are listed in tables (6),(7), and (8) respectively for STQ-1, STQ-2, and STQ-3 with same solvents.

Table(6)The rate of charge transfer constant STQ-1 in organic media

Wave length	Rate of charge transfer(s^{-1})						
	500 nm	550nm	600nm	650nm	700nm	λ_{ab} 450nm	λ_{em} 630nm
Cyclohexane	404643×10^{-13}	1.3783×10^{-10}	1.6368×10^{-7}	9.33×10^{-7}	2.9840×10^{-5}	1.233×10^{-13}	2.002×10^{-7}
Dioxane	1.2570×10^{-7}	1.2277×10^{-5}	15.587×10^{-4}	1.414×10^{-2}	2.256×10^{-8}	4.4956×10^{-8}	4.135×10^{-3}
Butanol	4.9419×10^7	2.1601×10^8	7.3844×10^8	2.0900×10^9	5.0975×10^9	3.549×10^7	1.4066×10^9
Ethanol	1.0737×10^8	4.3620×10^8	1.4033×10^8	3.7730×10^9	8.8062×10^9	7.8348×10^7	2.5895×10^9
Methanol	1.9866×10^8	7.6199×10^8	2.3360×10^9	6.0294×10^9	1.3588×10^{10}	1.4691×10^8	4.2029×10^9

Table(7)The rate of charge transfer constantSTQ-2 in organic media

Wave length	Rate of charge transfer(s^{-1})						
	500 nm	550nm	600nm	650nm	700nm	λ_{ab} 485nm	λ_{em} 625nm
Cyclohexane	15.7689	2.7673×10^2	3.0125×10^3	2.2727×10^5	1.2842×10^5	5.9532	8.613×10^4
Dioxane	7.8950×10^3	7.7942×10^4	5.2535×10^5	2.6414×10^6	1.0541×10^7	3.6245×10^3	1.2163×10^6
Butanol	1.1696×10^{11}	2.4446×10^{11}	4.5187×10^{11}	7.6005×10^{11}	1.1887×10^{12}	9.1034×10^{10}	5.9212×10^{11}
Ethanol	1.7021×10^{11}	3.4304×10^{11}	6.1514×10^{11}	1.0084×10^{12}	1.5403×10^{12}	1.3413×10^{11}	7.9537×10^{11}
Methanol	2.2905×10^{11}	4.4847×10^{11}	7.8505×10^{11}	1.2609×10^{12}	1.8927×10^{12}	1.8227×10^{11}	1.0043×10^{12}

Table(8)The rate of charge transfer constantSTQ-3 in organic media

Wave length	Rate of charge transfer(s^{-1})						
	500 nm	550nm	600nm	650nm	700nm	λ_{ab} 530nm	λ_{em} 623nm
Cyclohexane	4.500×10^{-46}	1.698×10^{-40}	6.4800×10^{-36}	4.8898×10^{-32}	1.028×10^{-28}	7.7248×10^{-56}	4.7148×10^{-34}
Dioxane	6.724×10^{-34}	1.664×10^{-29}	7.6352×10^{-26}	9.5909×10^{-22}	4.341×10^{-20}	2.1568×10^{-35}	3.1178×10^{-24}
Butanol	0.22548	5.8489	88.278	8.7816×10^2	6.2890×10^3	6.6004×10^{-4}	2.6601×10^2
Ethanol	1.2871	2.8466	3.7577×10^2	3.3373×10^3	2.1687×10^4	5.0313×10^{-3}	1.0723×10^3
Methanol	5.1394	1.0003×10^2	1.1870×10^3	9.6340×10^3	5.7951×10^4	0.0252	3.2441×10^3

Discussion:

The charge transfer in dye-organic media complex were investigated by theoretical calculations. The Golden rule expression and the continuum model for reorientation energy applied in our theoretical studies of charge transfer reactions in dye –organic media provide a constant value for the reaction. One of the start point of these studies were calculated of reorientation energy of the STQ-1, STQ-2, and STQ-3 in several organic solvents, Cyclohexane, Diaxane, Butanol, Ethanol, and Methanol, respectively. The results shift increasing with the high polarity solvent and large dipole moment of dye in excited state . Table (3) shows the results of reorientation energies for STQ dyes. Large dielectric constant of solvent results large values of λ for STQ dyes compare with less polarity of solvent , that's are shown in table (3). On the other hand the large values of dipole moment μ_e results to large reorientation energy, that's very view, for STQ-2 have $\mu_e = 10.5 \text{ Debye}$ compare with

STQ-1 have $\mu_e = 7.43 \text{ Debye}$ and $\mu_e = 5.22 \text{ Debye}$ for STQ-3.

Table (4)give the results of the effective free energy. It is clear that the dye have small effective free energy where have large reorientation energy and vice versa. The negative results in table (5) for STQ-2 indicate the free energy ΔG_0 is a part of energy that broken to rearrangement of the system to start transfer of charge .

Table (5) show the effect of the barrier height $\Delta G^\#$ on the rate constant of charge transfer, we note that the barrier in STQ-2 dye is small compare with the other dyes , STQ-1 more large height barrier, and STQ-3 high large barrier that give large values rate constant in STQ-2 compare with the other dyes, that's means k_{ET} increasly, where height barrier $\Delta G^\#$ decreasing. Generally from the results of rate constant of charge transfer in STQ dyes in organic media ,we found the major change in rate constant occurs in large reorientation energy system and small barrier height and negatively effective free energy. On the other

hand large rate constant indicate a distribution of charge in the state of dye (ground and excited state) give rise to dipole and higher order moment.

Conclusions:

In this work, the charge transfer reactions of a styryl STQ dyes has been studied theoretically in organic media . The reorientation energy are calculated according to continuum model and energy are calculated according to continuum model and dipole moment.

It should be noted that the rate constant at charge transfer for STQ-2 is always large than the rate constant for STQ-1 and STQ-2 in the same organic media, this indicate because large μ_e , and we observe that solvent much polar guide much charge transfer. The most electronically active dye leads to height rate charge transfer.

The rate of charge transfer in STQ dyes in organic media system is a function of height barrier that's mean, when the potential barrier large results small rate constant of charge transfer and vice versa.

In summary, we can concluded from present results that STQ-2 dye is more reaction for charge transfer compare with STQ-1 and STQ-2 dye, means that STQ-2 have large a distribution of charge in the ground and excited state that give dipole moment results, lead to high charge transfer rate constant.

Reference:

1. Dimitry,A.Ch.;lev.I.K.;and Armen,Y.M.; 2001"Photo Synthetic Electron Transfer Controlled by Protein Relaxation"Bio Phys.J. 50 (1033-1034).
2. Eranda,M.; and Francis,D.S; 2007 " Design and Studies of Photo Induced

Electron Transfer in Pacman" Ann.GRASP. 167.

3. Bumm,L.A.; Arnold,J.J.; Dunbar,T.D.; Allara,D.L.; and Wiss,P.S.; 1999 " Electron Transfer Through Organic Molecular "J.Phys Chem.B, 103:8122-8127.
4. Jung,Y.J.; and Gao,J.; 2002 " Spectral Analysis of Electron Transfer Kinetic " J.Chem-Phys, 117(8): 3822-3823.
5. John ,N.G.; and Chandler,D.; 1992 " Quantum Theory for Free Energies of Electron Transfer " J.Chem. Phys., 97(7) 4958-4963.
6. Andrei,T.; 2008" Marcus Theory for Electron Transfer a Short Introduction " J. Club. Mazin,MPIP. Department of Chemistry.
7. Koti,A.S.R.; Bhattachrjee,B.; Haram,N.S.; Ranjan,D.; Periasamy,N.; Sanawana,N.D.; and Rangnekar,D.W. 2000; " Photo Physics of Same Styryl Thiazolo Quinoxaline Dyes in Organic Media " J.Phptchem. and PHOTO boil.A.Chemistry, 137: 115-123.
8. Keitaro,Y.; Keisuke, T.; and Yutak,N.; 1995 " Effect of the Solvent Dynamics and Vibrational Motion Electron Transfer "Ball.Chem.Soc.J, 68: 696-712.
9. Dennis,A.; and Joseph, T.; " Thermal activated Inverted Electron Transfer Kinetic " J.Am. Chem. Soc. 122:10950- 10965.
10. Andre,J.M., " Quantum Theory of Polymers " 2006, Socrates Erasmus Programe, University of Warsaw.
- 11.Deanna,M.D.;Alessandro,Peter,H.D .;Joseph,T.H.; Peter.C.J. and Keene,F.R.; 2006 " The Effective Electron Transfer Distance in Dinuclear Ruthenium Complexes "Eur.J.Inorg. Chem.,; 772-783.
12. Plenn,B. 2002; " Electron Transfer in donor-Bridge-Acceptor System and Derized Malerals ", Ph.D Thesis Debye Institute Vande University of Utrecht.
13. AL-agealy,H.J.; AL-obaidi,R.I.; And Al-saedi,R.F.; 2008 " Effect of

Solvent Types on Photo Induced Electron Transfer in Methylene Blue Dye " Ibn AL-Haitham J. for Pure and Appl.Sci, 21(3): 39-49.

14. AL-agealy,H.J.; 2010 " Electron Transfer between Excited Cresyl Violet CV Dye and Semiconductor at Acetonitrile Solvent " J.Thi-Qar Sci, 2(3): 213-221.

15. AL-agealy,H.J.; 2009 " Photo Induced Electron Transfer Between Ruthenium and Methyl Viologen " Ibn AL-Haitham J. for Pure and Appl.Sci, 22(3):93-113.

16. Chandrakhar,K.; Naik,L.R.; Suresh,K.H.; and Math,N.N.; 2006 " Steady state Absorption and Fluorescence Study " Indian. J. Pure Appl. Phys, 44:292-299.

دراسة نظرية للشحنة المنتقلة في صبغة الكواينوكسلاين STQ-1, STQ-2, دراسة نظرية للشحنة المنتقلة في صبغة الكواينوكسلاين STQ-3 في أنظمة الاوساط العضوية

خالد هلال حربي*
رفاه نوري اسماعيل*

هادي جبار مجبل*
محسن عنيد حسوني*

*جامعة بغداد/كلية التربية للعلوم الصرفة ابن الهيثم / قسم الفيزياء

الخلاصة:

دُرِس انتقال الشحنة في صبغات STQ-1, STQ-2, STQ-3 في أنظمة الاوساط العضوية وحسبت نظريا اعتمادا على قاعدة فلرانك كوندن وانموذج ثنائي القطب المستمر. وحسبت طاقات اعادة الالتحام λ (eV) نظريا اعتمادا على زخم ثنائي للقطب μ وثابت العزل ϵ ومعامل الانكسار n . كما تم حساب ثابت معامل الانتقال للشحنة اعتمادا على طاقة اعادة الترتيب λ (eV) والطاقة الحرة الفعالة ΔG^* وارتفاع حاجز الجهد ΔG^\ddagger ومعامل الازدواج. كتب برنامج بلغة ماتلاب لحساب ثابت معدل الانتقال للشحنة وبقية المعاملات اظهرت نتائج الحساب ان صبغة STQ-2 اكثر فعالية للانتقال الالكتروني قارن ب STQ-1 و STQ-3.