

DIAGNOSTIC STUDY ON MICROFILARIAE AND SOME BLOOD PROTOZOA IN QUAIL BIRDS (*COTURNIX COTURNIX JAPONICA*) IN NINEVEH GOVERNORATE

Rawaa G. Mohammad

Department of biology, Faculty of health and science, University of Koya, Iraq.

(Received 6 May 2012, Accepted 23 March 2012)

Keywords; microfilaria, quail, Knott technique.

ABSTRACT

The study covered a diagnosis for detection of microfilaria in 90 blood samples obtained from quail birds. The results recorded that total infestation rate was (15.6%).

Three species of microfilaria have been diagnosed in quail. Form A of detected larvae measured 106 – 125 μm in length and 2.5 – 3.0 μm in width, while form B measured 97–145.4 μm in length and 4.0 – 6.0 μm in width, finally form C dimensions measured 120–150 μm and 4.5 – 5.0 μm in width respectively. Knott technique detected most positive cases with higher sensitivity rate 85.7%, than other techniques used in the study. The rate of infection recorded in adult birds 14.5% was little higher than it in young birds 10.7% with no significant statistical differences between males and females. The study showed appearance of 7.7% of mixed infections of microfilaria with *Aegyptianella spp.* and 4.4% with *plasmodium spp.* Finally the study pointed that microfilaria appeared with high percentage in circulating blood than peripheral blood of 40 quail birds, as 15% and 10% in circulating and peripheral blood respectively.

INTRODUCTION

Japanese quail (*Coturnix coturnix japonica*) belong to Family: Phasianidae of the Order: Galiformes of the Class: Aves, (1), and has been introduced to Iraq as domestic bird because it has desirable meat which has tasteful, energy higher than in chickens meat and for eggs productions (2). The birds have considered one of the common birds farm in Egypt and Saudi Arabia for its characteristics mentioned above and has been used as laboratory animals(3), although it poses quality resistance to disease than those of chickens but may hindered by various problems such as haemoparasites which include one or more genera typically live in blood of the host during at least some development(4).

Filarial nematodes larvae whose adult stages live in internal organs and cavities, circulates in blood from where they are ingested by vector during blood meal (5) and may causes ecological and behavioral changes in avian host and induced morbidity and mortality in susceptible animals (6). The present study was conducted to determine the incidence and diagnosis of microfilaria in

blood of quail bird in nineveh governorate and teste the best method and site for diagnosis it in blood . The study considered the first deal with this aspect in quail at nineveh governorate.

MATERIAL AND METHODES

A total of (130) quail species (*Coturnix coturnix japonica*) were collected from different areas in nineveh governorate. Blood samples were collected from 90 quail birds in different ages and both sexes at the time of slaughter from jugular vein and the blood was kept in vials with anticoagulant . All samples examined by using three laboratory techniques; wet blood film technique (7), Giemsa stain(thin blood smear) technique(8), and knott's technique .

Microfilaria and other blood parasites measured with ocular micrometer and identified depending on data given by (9,10,11) while intensity calculated according to (12).

Thin blood smear prepared from (40)quail birds collected from preipheral blood (wing vein) according to (11) befor slaughter birds and from circulating blood (lung) as in (13) after slaughtered birds to compare between two sites for isolation of microfilaria . chi-square test were used to compare among the resultes (14).

RESULTS

Examination of (90) blood samples from quail birds (*Coturnix coturnix japonica*) revealed presence of microfilaria with total infection rate(15.5%). (Table ,2).Three forms of microfilaria have diagnosed in blood samples based on morphological characteristics with low intensity for all three forms as shown in table (1).Figures (1,2,and 3).

The current resultes appeared that knott's technique is the most sensitive technique for detection of microfilaria in blood among other techniques, it has detected (12) out of (14) positive sample detected by all techniques used in the study, with sensitivity reached to (85.7%) followed by wet blood film technique which considered also as suitable technique for microfilaria detection which is gave sensitivity (71.4%) and Giemsa stain with sensitivity (57.1%).(Table ,2). Figures(4 and 5).

The study showed that percentage of infections in adult birds was higher than it in young which represented with (14.5%) and (10.7%) in adult and young respectively, (table,3) .This difference was not significant at ($p>0.05$) . Also there was no significant differences between females and males . Females recorded (12.7%) of infection while males represnted (14.3%).(Table,4).

Table (5) showed presence of mixed infections of microfilaria with *Aegyptianella spp.* in 7 (7.7%), *Plasmodium spp.* in 4 (4.4%) of positive samples and mixed infection of microfilaria with both above parasite genera in 3 (3.3%) of positive blood samples .

The study notes high single infections of *Aegyptianella spp.* parasites. It is found in 16 blood samples of quail with percentage (17.7%) and intensity moderate which appears as redish

organism in erythrocytes with varied sizes from (0.4-1) μm . (Fig, 6), and low single infection with *plasmodium spp.* with percentage (4.4%) and low intensity. (Fig,6). (Table,6).

Comparative study between peripheral blood (wing vein) and circulating blood(lung) of 40 examined quail bird indicated that microfilaria was higher in circulating blood (15%) than in peripheral blood (10%) with significant difference between them at ($p < 0.05$). (Table ,7).

Table (1) Measurements of three forms of microfilaria recovered from (90) blood of quail birds.

Form of microfilaria	Length(μm) mean (range)	Width(μm)	Characteristic feature
Form A	112.4 (106-125)	2.5-3	Unsheathed
Form B	118.7 (97-145.4)	4.0-6.0	Unsheathed
Form C	129.4 (120-150)	4.5-5	Sheathed

Table (2) Number of positive blood samples with different techniques and sensitivity of each.

No. of samples	Positive samples No. (%)	Techniques					
		Knott's		Wet blood film		Giemsa stain	
		Positive	Sensitivity(%)	Positive	Sensitivity(%)	Positive	Sensitivity(%)
90	14 (15.5)	12	85.7	10	71.4	7	57.1

Table (3) Percentages of infection with microfilaria in (90) blood samples of quail bird according to age.

Age	No. of examined samples	No. of positive sample	Percentage of infection(%)	Signification
Adult(more than 10 weeks)	62	9	14.5	Not significant at $p > 0.05$
Young (less than 10 weeks)	28	3	10.7	

Table (4) Percentages of infection with microfilaria in (90) blood samples of quail birds according to sex.

Sex	No.examined	No.Positive sample	Percentage of infection(%)	Signification
Females	55	7	12.7	Not significant at $p>0.05$
Males	35	5	14.3	

Table(5) Mixed infections of microfilaria with the other blood parasites in (90) blood samples of quail birds.

Type of parasite mixed with microfilaria	No. of infected samples	Percentage of infection(%)
Microfilaria and <i>Aegyptianella</i> spp.	7	7.7
Microfilaria and <i>plasmodium</i> spp.	4	4.4
Microfilaria , <i>Aegyptianella</i> spp. and <i>plasmodium</i> spp.	3	3.3

Table (6) percentages of single infections with blood parasites in (90) blood samples in quail bird and intensity of each one.

Type of blood parasites	No. of Positive samples	Percentage of infection(%)	Intensity
Microfilaria	14	15.5	Low
<i>Aegyptianella</i> spp.	16	17.7	Moderate
<i>plasmodium</i> spp.	4	4.4	Low

Table(7) Prevalence of microfilaria according to site of collected blood sample in (40)quail birds.

Site of collection	No. of positive samples	Percentage of infection(%)
Peripheral blood(wing vein)	4	10
Circulating blood (lung)	6	15



Fig. (1) form A/ unsheathed microfliaria.




Fig. (2) form B/ unsheathed microfliaria.(posterior end).



Fig.(3) form C/ sheathed microfilaria.




Fig.(4) unidentified microfilaria in blood of quail bird(Knott technique,10x).



Fig.(5) unidentified microfilaria in blood of quail bird(Giemsa stain technique,40x).




Fig.(6) Single infection of *Aegyptianella* spp.in blood film of quail bird(Giemsa stain technique,100x).




Fig.(7) Single infection of *plasmodium* spp.in blood film of quail bird(Giemsa stain technique,100x).

DISCUSSION

Quail birds deserve care since they have many benefit characteristics as it is resistance to different diseases, not need big distance to farm, and considered as important source of animal proteins which is necessary to human.

In the current study (14) quail birds from (90) examined for presence of microfilaria harboured nematode larva (microfilaria) in blood with total infection rate (15.6%). This study is the first deals with this aspect in quail in Iraq, there is no study that deals with this parasites in our country but studies performed in different parts in world indicated presence the parasite in birds, such study in Texas (2) which observed that (25%) of quail examined in Brewster country, Texas was infected with microfilaria, and other study which considered avian filariasis occur in wide range in avian species and may be contributing factor in causing diseases or death (15). Distribution of infection with microfilaria depends upon environment demands of vector and different exposure of birds to vector, the exposure may be depends on the time of daily activities of individuals species and selection place for nesting, rest (4).

From the laboratory techniques used under this study to diagnose microfilaria in blood of quail, Knott's technique was the best for identification which recorded sensitivity (85.7%), these findings disagree with (7) but agree with (16) which was considered this technique as the best and give accurate results in number of positive samples diagnosed and time saved.

The morphology and size of microfilaria detected in our study approached with the results given by several studies such as (11), current study did not identify species because the adult worms; which parasitized the body cavity and air sac that may belong to several genera like ornithofilaria, sarconema and splendidofilaria; can not be observed.

The percentage of infection is little higher in adult birds (14.5%) than young birds (10.7%), this result agrees with (17) while disagrees with (18); this may reflect along prepatent period or a greater probability of exposure to infected intermediate host with age (19) on the other hand there is no statistical differences in rates of infection between male and female of quail examined, these results are similar to those obtained by (20) and may be attributed to that male and female affected equally for contributing factor to disease.

The study evidence appearing of mixed infections of microfilaria with *Aegyptianella* spp which is intraerythrocytic parasite of domestic birds which now placed in family Anaplasmataceae (21) as percentage (7.7%). (15) found similar association in chickens kept near cages of birds, also my study observed high single infection with *Aegyptianella* spp in blood of quail as percentage (17.7%) which indicates importance of this parasite in quail that it may become high pathogenic and often fatal (22) while low mixed infections of *plasmodium* spp with microfilaria (3.3%) and low single infections with *plasmodium* spp (4.4%) in quail indicate less

susceptible of quail to infection with plasmodium parasite. Presence of these three types of parasites (microfilaria, *Aegyptianella spp.* and *plasmodium spp.* in 3 blood smears of quail in my study varies from the results observed in other studies in world because of several factors may affect the presence of blood parasites such as type of bleeding, presence suitable environment to vector and immuno status of birds.

Results of the study agreed with the other accompanied studies such as (13,17), studied on Willo ptarmigan and passerine birds respectively, that indicated the relationship between the site of blood collection and the prescence of blood parasites. Circulating blood (heart and lung) is the more correct site for diagnosing study of microfilaria than peripheral blood (wing vein) that microfilaria congregate in deep circulation especially in lung where flow rate of blood is slow (23).

دراسة تشخيصية عن اليرقات الخيطية الدقيقة وبعض الاوالي الدموية في طائر السمان (*Coturnix coturnix japonica*) في محافظة نينوى

رواء غانم محمد

قسم علوم الحياة، هيئة العلوم والصحة، جامعة كويه، العراق.

الخلاصة

اشتمل البحث على دراسة تشخيصية للكشف عن اليرقات الخيطية الدقيقة في 90 عينة دم من طائر السمان. شاركت النتائج الى ان النسبة الكلية للخمج بلغت 15.6%، شخّصت ثلاثة انواع منها في طائر السمان تراوحت طول النوع A بين 106-125 مايكرومتر والعرض 2.5-3.0 مايكرومتر والنوع B بين 97-145.4 مايكرومتر والعرض بين 4.0-6.0 مايكرومتر والنوع C بين 120-150 مايكرومتر والعرض بين 4.5-5.0 مايكرومتر. كشفت تقنية Knott عن معظم الحالات الموجبة اذ سجلت اعلى نسبة كفاءة من بين التقنيات الاخرى المستعملة بلغت 85.7% سجلت نسبة الخمج في الطيور البالغة ارتفاعا قليلا عنها في الصغيرة العمر اذ بلغت 14.5% و 10.7% في الطيور البالغة والصغيرة على التوالي مع عدم وجود فرق معنوي بين الذكور والاناث. وظهرت الدراسة وجود خمج مختلط لطيفلي اليرقات الخيطية الدقيقة مع طفيلي *Aegyptianella spp.* بنسبة بلغت 7.7% ومع طفيلي *Plasmodium spp.* بنسبة بلغت 4.4% ووجد من خلال فحص عينة دم مأخوذة من الدم المحيطي الوريد الجناحي ومأخوذة من الدم الدائر (الرئة) من 40 طائر من السمان ان نسبة تواجد طفيلي اليرقات الخيطية الدقيقة في الدم الدائر اعلى من الدم المحيطي اذ بلغت 15% و 10% في الدائر والمحيطي على التوالي.

REFERENCES

- 1- Al-Lous B. Iraqi birds. Baghdad, Iraq, 1961.
- 2- Dehka K and Borah J. Hematological and biochemical changes in japanse quails *coturnix coturnix japonica* and chickens due to *Ascaridia galli* infection. J. poultry sci. 2008;7:704-710.
- 3- Al- Shaheen, Sajida A M. Sellation of body weight, growth rate, and potency of feeding conversion in qual bird (*Coturnix coturnix japonica*). Msc. Thesis, college of agriculture, Basrah uni. 2006

- 4- Hauptmanova K,Benediikt U and Literak I.Blood parasites in passerine birds in Solvakian east .carpathians.Acta protozool .2006;45:105-109.
- 5- Kucera J,Blood parasites of birds in central Europe.4.*Trypanosoma oplasma*”*Microfliaria* and other Haematozoa. Folia pparasitologica(praha).1982,29:107-113.
- 6- Bradly R,Fedynich AM and Lerich SP.Survey on Haemoparasities in scaled quail from elephant mountain wildlife management area .Brewster country ,Texas. Texas J Agriculture and natural Reso.2004,17:18-22.
- 7- Goldsmid JM.Studies on the laboratory diagnosis of human filariasis :Preliminary communication .J.clin.path.1970,23:632-635.
- 8- Razijalali MH,Alborzi AR,Avizeh R and Mosalianejad B.Astudy on *Dirofilaria immitis* in healthy urban dogs from Ahvaz ,Iran.Iran.J.Vet.Res.2010;4:357-362.
- 9- Kassai T.Veterinary Helminthology .1ed.Great Britian,Bath press,someret.1999. p:119-133.
- 10- Hauptmanova K,Barus V,Literak I and Benedikt V.Haemoproteids and microfilaria in hawfinches in the Czech Republic.Helminthologia.2004;4:125-133.
- 11-Rukhsana T.Infection of haematozoan parasites found in birds of NWFP (pakistan) pakistan j .Biolo.Sci.2005;8:1-5.
- 12-Tarofs A and Stutchbury B.Low infection prevalence of blood parasites in hoded warblers.J.Field. Ornithol.1997;28:75-78.
- 13-Holmstad P.Anwar A,Lezhova Tand Skorping A.Standerd sampling technique underestimate prevalence of avain hematozoa in Willow ptarmigan (*Lagopus lagopus*) J of Wildlife dis.2003;39:354-358.
- 14-Petrie A and Watson P.Statistics of veterinary and animal science.Black Well Science.London .2003.
- 15- Peirce MA and Bevan BJ.Blood parasites of imported psittacine birds.Vet.Rec.1977; 2:282-283.
- 16- Gadahi JA,Arijo AG,Abubakar M,Javaid SB and Arshed MJ.Prevalence of blood parasites in stray dog in Hyderabad Area:Comparative sensitivity of different diagnosis technique for detection of microfliaria.Vet World. 2008;9:229-232.
- 17- Kirkpatrik CE and Suthers HB.Epizootiology of blood parasites infection in passerine birds from Cetral New Jersey.Canadian J OF Zoology. 1988;66: 2374-2382.
- 18- Ravinder NM ,Hugh IJ and Thomas BS.Molecular evidence for hostspecificity of parasitic nematode microfliaria in some African rainforest birds.Molecular Ecology .2005;14:3977-3988.
- 19- Rabinowitz A,Sharon P,Vicki M.Microfliariae of *Tetrapetalonema llewellyni* in raccoons of cades cova.Great smoky mountains national park,Tennessee,Usa.J.Wildlife.Dis.1985;21:449-450.

20- Stannard AA and Cello RM. Onchocerca cervicalis infection in horses from western united states. American J. Vet. Res. 1975;36:69-74.

21- Peirce MA. A new species of Aegyptianella from south-east Asia. Vet. Rec. 1999 ;

145 :288 .

22- Gastle MD and Christensen BM. Isolation and Identification of *Aegyptianella pullorum* (Rickettsiales, Anaplasmataceae) in wild turkeys from north America. Avian Dis. 1984;29:437-445.

23- Hawking F. Circadian and other rhythms of parasites. Advances in parasitology. 1975;123-182.