

SYNTHESIS AND CHARACTERIZATION OF SOME NEW (3-ACETYL-2H-CHROMEN-2-ONE) DERIVATIVES

تحضير وتشخيص بعض المشتقات الجديدة ل (3-اسيتيل-2H-كرومين-2-كيتون)

Suzanne Jubair Abass*, Jallal Hassan Muhammed* and Lamyaa Salih*

***= College of Pharmacy / University of Kerbala**

ABSTRACT

Compound S₁ which is the starting compound was prepared by the reaction between salicylaldehyde and ethyl acetoacetate in the presence of piperidine. In order to prepare compound S₂, compound S₁ was refluxed with 2-aminothiazole that dissolved in ethanol in presence of glacial acetic acid. S₃ was prepared through addition of bromine that dissolved in chloroform to compound S₁. S₄ was prepared by heating compound S₃ with thiourea in the presence of sodium acetate. S₅ was prepared by refluxing of compound S₄ with ethyl acetoacetate, while S₄ on refluxing with succinic anhydride give S₆. S₆ when refluxed with *o*-phenylenediamine give compound S₇. All the synthesized compounds were characterized through their physical properties and diagnosed by using FTIR and ¹H NMR spectrophotometric techniques.

الخلاصة

تم تحضير المركب S₁ الذي هو المركب الابتدائي والأساسي لتحضير بقية المركبات من خلال التفاعل بين السلسلديهايد واثيل اسيتواسيتيت بوجود البيريدين ومن اجل تحضير المركب S₂ تم تصعيد المركب S₁ مع المركب 2-امينوثيازول المذاب في الايثانول بوجود حامض الخليك الثلجي، اما المركب S₃ فقد تم تحضيره بإضافة البروم المذاب في الكلوروفورم الى المركب S₁. تم تحضير المركب S₄ من خلال تسخين المركب S₃ مع الثايورييا بوجود خلات الصوديوم. المركب S₅ عندما تم تصعيده مع الاثيل اسيتواسيتيت تم الحصول على المركب S₆. اما المركب S₇ فتم الحصول عليه بتصعيد المركب S₆ مع الاورثو ثنائي الامين باستخدام الايثانول كمذيب. جميع المركبات المحضرة تم تميزها من خلال صفاتها الفيزيائية والكيميائية وتشخيصها باستعمال التقنيات الطيفية المتمثلة بتقنية الاشعة تحت الحمراء وتقنية الرنين النووي المغناطيسي للبروتون.

INTRODUCTION

Coumarins are large family of heterocyclic compounds that occupy an important position in natural and synthetic organic chemistry. Synthesis of coumarins and their derivatives has attracted a large attention from organic and medicinal chemists for many years as a large number of natural products contain this heterocyclic nucleus^[1]. There are many methods for synthesis of coumarins^[2,3,4], Knoevenagel reaction is the one that have been adopted in this research. On the other hand the nitrogen and sulfur heterocyclic system families are very interesting compounds due to their physicochemical properties especially in the matter of design new drugs^[5]. This study aim to combine the coumarinic system with the nitrogen and sulfur heterocyclic system in the hope of getting more potent pharmacological compounds.

EXPERIMENTAL

Materials:

All chemical materials were purchased from BDH, Scharlau and Himedia and were used without further purification.

Instruments:

Melting points were measured on a Gallan Kamp MFB-600 Melting point apparatus and were uncorrected. FTIR spectra were recorded as potassium bromide (KBr) disk on FTIR-8400S Fourier Transform Infrared Spectrophotometer "SHIMADZU". ¹H NMR spectra were recorded on Burker DMX- 500 NMR (300-600 MHz) Spectrophotometer with using DMSO as a solvent in Jordan University.

Preparation of compound S₁ ^[7]

(3-acetyl-2H-chromen-2-one)

Salicylaldehyde (3.05g ,0.025mol) and ethyl acetoacetate (3.25g ,0.025 mol) was mixed, stirred and cooled .To this mixture 10 mL of piperidine was added with shaking .The mixture was kept in refrigerator for 2 hrs , after that a yellow solid was obtained, then was recrystallized from ethanol. The physical properties were listed in table 1.

Preparation of compound (S₂) ^[8]

(3-[(1Z)-N-(4,5-dihydro-1,3-thiazol-2-yl)ethanimidoyl]-2H-chromen-2-one)

Compound S₁ (0.22g, 0.001mol) was dissolved in 40mL of warm ethanol contains 5 drops of acetic acid, and 2-aminothiazole (0.1g, 0.001mol) was added. The resulting mixture was refluxed for 10 hrs. After cooling, the formed solid was filtered off, dried and recrystallized from ethanol. The physical properties were listed in table 1.

Preparation of compound (S₃) ^[8]

(2-oxo-2H-chromene-3-carbonyl bromide)

To a solution of compound S₁ (2.19g ,0.01mol) in 100 mL of ethanol ,bromine (1.6g ,0.01mol) in 25mL chloroform was added with shaking. The mixture was warmed to decompose, then it was heated for 15 minutes. The mixture was cooled and filtered to get a solid precipitate which on washing with diethyl ether gave the desired product which was recrystallized from acetic acid to give colorless needles. The physical properties were listed in table 1.

Preparation of compound S₄

(3-(2-amino-4,5-dihydro-1,3-thiazol-4-yl)-2H-chromen-2-one)

A suspension of compound S₃ (1.49g, 0.005mol) and thiourea (0.38g, 0.005mol) in 50 mL ethanol was heated to give a clear solution that gave deposited crystals on further heating. The crystals were filtered, washed with ethanol and then boiled with water containing sodium acetate yielding the target compound. The product was recrystallized from ethanol. The physical properties were listed in table 1.

Preparation of compound S₅ ^[9]

(7-methyl-3-(2-oxo-2H-chromen-3-yl)-2,3-dihydro-5H-[1,3] thiazolo[3,2-a]pyrimidin-5-one)

A mixture of compound S₄ (0.24g, 0.001mol) and ethylacetoacetate (0.65g, 0.005mol) was refluxed for 3hrs. After cooling, the formed solid was filtered off, dried and recrystallized from ethanol. The physical properties were listed in table 1.

Preparation of compound S₆ ^[10]

(4-oxo-4-[[4-(2-oxo-2H-chromen-3-yl)-4,5-dihydro-1,3-thiazol-2-yl]amino]butanoic acid)

A mixture of compound S₄ (0.24g, 0.001mol) and succinic anhydride (0.15g, 0.001mol) in glacial acetic acid was refluxed for 3hrs and then was cooled. The formed solid was filtered off, dried and recrystallized from dioxane. The physical properties were listed in table 1.

Preparation of compound S₇ ^[11]

(3-(1H-benzimidazol-2-yl)-N-[4-(2-oxo-2H-chromen-3-yl)-4,5-dihydro-1,3-thiazol-2-yl]propanamide)


Compound S₅ (1.7g, 0.005mol) and *o*-phenylenediamine (0.54g, 0.005mol) were dissolved in absolute ethanol and was refluxed for 12hrs. The resulting mixture was cooled and poured onto 30mL of ice cold water containing 1mL of conc. HCl. The precipitate which was allowed to settle down, was filtered off, dried and recrystallized from ethanol. The physical properties are listed in table 1.

Table 1: The physical properties for the synthesized compounds

Comp. No.	Color	m.p ⁰ C	Yield%	Mol. Formula
S ₁	yellow	123-125	85	C ₁₁ H ₈ O ₃
S ₂	Greenish yellow	152-154	93	C ₁₈ H ₁₂ N ₂ SO ₂
S ₃	pale yellow	163-165	55	C ₁₁ H ₈ O ₃ Br
S ₄	Yellow	220-222	56	C ₁₂ H ₈ N ₂ SO ₂
S ₅	Brown	190-192	78	C ₁₅ H ₁₀ N ₂ O ₄
S ₆	Pale yellow	210-212	53	C ₁₆ H ₁₂ N ₂ SO ₅
S ₇	dark yellow	240-243	69	C ₂₂ H ₁₆ N ₄ SO ₃

RESULTS AND DISCUSSION

All the synthesized compounds were characterized through their physical properties and diagnosed by using FTIR and ¹H NMR spectrophotometric techniques.



Synthesis and characterization S₁

The synthesis of this compound was achieved by the reaction between salicylaldehyde and ethyl acetoacetate in the presence of piperidine. This compound is confirmed by FT-IR spectrum which shows appearance of a strong band at 1741 cm^{-1} for the stretching frequency of carbonyl of the lactone ring [12] and another band appears at 1680 cm^{-1} for the ketone which is conjugated with (C=C) group.

Synthesis and characterization of S₂

This compound was prepared as a Schiff's base which is the result from the reaction between the ketonic group of the coumarin molecule (compound S₁) with 2-amino thiazoline. Compound S₂ is confirmed by the FT-IR spectrum which shows band at 1608 cm^{-1} for (C=N) group and disappearance of the stretching band of the ketonic carbonyl group, while the stretching frequency of the (C-S) group of the thiazoline ring appeared at 1113 cm^{-1} .

Synthesis and characterization of S₃

This compound is synthesized by the bromination of compound S₁. FT-IR spectrum shows the appearance of a stretching frequency of (C-Br) group at 875 cm⁻¹ beside two carbonyl group stretching frequency at 1734 cm⁻¹ and at 1687 cm⁻¹.

Synthesis and characterization of S₄

Compound S₄ was prepared through the reaction between compound S₃ and thiourea. This compound was confirmed by its spectral data. FT-IR spectrum shows the appearance of double band at 3315 cm⁻¹ and 3383 cm⁻¹ belong to (NH₂) group and a stretching frequency at 1095 cm⁻¹ for (C-S) group beside the carbonyl group stretching frequency at 1747 cm⁻¹. It also show a single band at 1643 cm⁻¹ for (C=N) group. ¹H NMR spectrum shows peaks at: 2.4 ppm for (H, CH of thiazoline ring), 3.2 ppm for (2H, CH₂ of thiazole ring), 7.4-7.9 ppm for (4H, CH of aromatic ring + CH olefinic of coumarin ring), 9.8 ppm for (2H, NH₂).

Synthesis and characterization of S₅

The reaction between compound S₄ and ethyl acetoacetate produce compound S₅ which is confirmed by FT-IR spectrum that shows the appearance of a strong band at 1722 cm⁻¹ for the carbonyl group of the coumarin molecule and another band at 1637 cm⁻¹ for the carbonyl group of the pyrimidinone ring. It also shows the a stretching frequency at 2995 cm⁻¹ for the aliphatic (C-H) bond. ¹H NMR spectrum shows peaks at: 2.35 ppm for (3H, CH₃), 3.34 ppm for (H, CH of thiazoline ring), 3.99 ppm for (2H, CH₂ of thiazol ring), 6.45 ppm for (H, CH of pyrimidine ring) and 6.99-7.95 ppm for (5H, CH of coumarin rings).

Synthesis and characterization of S₆

This compound was prepared from the reaction of compound S₄ with succinic anhydride. FT-IR of this compound shows a broad band at 3257 cm⁻¹ for (O-H) group and another band at 3180 cm⁻¹ for (N-H) group beside three bands at 1722 cm⁻¹, 1714 cm⁻¹ and 1649 cm⁻¹, the first for stretching carbonyl group of the coumarin molecule, the second for the stretching carbonyl group of the carboxylic group and the third for the stretching carbonyl group of the amide group.

Synthesis and characterization of S₇

Compound S₆ on reaction with *o*-phenelendiamine produces compound S₇. FT-IR spectrum shows the appearance of two stretching frequencies at 3240 cm⁻¹ and at 3233 cm⁻¹ for two (N-H) groups and two bands at 1739 cm⁻¹ and 1645 cm⁻¹ the first for the carbonyl group of the coumarin molecule and the second for the carbonyl of the amide group. ¹H NMR spectrum shows peaks at: 2.35 ppm for (2H, CH₂ of aliphatic CH₂-C=O), 2.77 ppm for (2H, CH₂ aliphatic), 2.44 ppm for (H, CH of thiazoline ring), 3.83 ppm for (2H, CH₂ of thiazoline ring), 7.1-7.9 ppm for (8H, CH of aromatic ring + CH olefinic of coumarin ring), 11.5 ppm for (H, NH of imidazole ring) and 12.5 ppm for (H, NH of amide moiety).

Table 2: The FT-IR spectra data for the synthesized compounds

Comp No.	Characteristic bands of FT-IR (cm^{-1} , KBr disc)			
	ν (N-H) cm^{-1}	ν (C=O) cm^{-1}	ν (C-H) cm^{-1}	ν (others) cm^{-1}
S ₁	-----	1741,1680	ar.=3032 al.=2931	ν (C-O)= 1207 ν (C=C) =1610,1558
S ₂	-----	1734	ar.=3086 al. =2978	ν (C-S) =1113 ν (C=N)=1608,1612
S ₃	-----	1734,1687	ar.=3026 al. =2958	ν (C-O) =1074 ν (C-Br) =875
S ₄	3383,3315	1747	ar.=3002	ν (C-S) =1095 ν (C=N) =1643
S ₅	-----	1722,1637	ar. =3086 al. =2995	ν (C=N) =1604 ν (C-S) =1160 ν (C-N) =1187
S ₆	3144	1722,1714,1649	ar. =3051 al. =2982	ν (C-S) =1141 ν (C-O)=1185 ν (O-H)=3180
S ₇	3240,3233	1739,1645	ar =3059 al=2929	ν (C=N)=1600 ν (C-S) =1140 ν (C-S) =1199




Figure 1: FTIR Spectrum for compound S₁




Figure 1: FTIR Spectrum for compound S₄




Figure 3: ¹HNMR Spectrum for compound S₅

REFERENCES

- 1) Kinza Aslam, M. Kaleem Khosa, Nanzish Jahan and Sofia Nosheen, "Synthesis and applications of coumarines", *Pak. J. Pharm. Sci.*, 23(4),310-316 **2010**.
- 2) Bilijana R. Dekic, Nico S. Radulovic, Vidoslav S. Dekic, Rastko D.Vukicevic and Radosav M. Palic, "Synthesis and antimicrobial activity of new 4-heteroarylaminocoumarin derivatives containing nitrogen and sulfur as heteroatoms", *Molecules* ISSN 1420-3049, **2010**.
- 3) Borges F., Roleira F., Milhazes N., Santana L. and Uriarte E., "Simple Coumarins and Analogues in Medicinal Chemistry: Occurrence, Synthesis and Biological Activity", *Current Med. Chem.*; 12, 887-916, **2005**.
- 4) Trkovnik M., Ivezi Z., "Synthesis of some new coumarinquinolone carboxylic acid ", *J. Heterocyclic Chem.*, 37, 137, **2000**.
- 5) Nachiket S. Dighe, Shashikant R. Pattan, Santosh S. Dengale, Deepak S. Musmade, Madhuri Shelar, Vishal Tambe and Mangesh B. Hole , "Synthetic and pharmacological profiles of coumarins: A review ", *Sch.Res.Lib. Archives of Applied Science Research*, 2 (2): 65-71, **2010**.
- 6) Madhavan G.R., Balraju V., Malleshham B., Chakrabarti R., Lohray V.B., "Novel coumarin derivatives of heterocyclic compounds as lipid lowering agents", *Bioorg. Med. Chem. Lett.*, 13, 2547, **2003**.
- 7) Nadeem Siddiqi, Faiz Arshad and Suroor Khan, "Synthesis of some new coumarin incorporated thiazolyl symicarbazonas as anticonvulsants ", *Pharm. Soci.* 66(2), **2009**.
- 8) Upendra Kumar K., Yashovardhan, Sudhir Kumar B. and Ashok Kumar, "Synthesis of new 10-substituted phenothiazines as anti-inflammatory and analgesic agents", *Int.J.of Phrm. Bio.Sci.* ; 1, Issue -3, **2010**.
- 9) Ashraf A. Khalil , Sami G. Abdel Hamide ,Abdulrahman M. Al Obaid and Hussein I. Al Subbagh , "Substituted quinazolines ,part2 , Synthesis and in-vitro anticancer evaluation of new 2-substituted mercapto-3-H-quinazoline analogs", *Arch. Pharm. Med. Chem.* .; 2,95-103, **2003**
- 10) Irina O. Zhuravel, Sergiy M. Kovalinko, Sergiy V. Vlasov and Valentin P. Chernykh, "Solution-phase synthesis of combinational library of 3-[4-(coumarin-3-yl)-1,3-thiazol-2-yl carbamoyl] propanoic acid amides", *Molecules* ; 10, 444-456, **2005**.
- 11) Vinod Kumar P., Mridula U. and Mrinaljni U., "Benzimi-dazolyl quinolinyl mercaptotriazoles as potential antimicrobial and antiviral agents", *Acta. Pharm.*; 55: 47-56, **2005**.
- 12) Robert M. Silverstein ,Francis X. Webster , Daved J. Kiemle , "Spectrometric Introduction of Organic Compounds" ,7th Edition ,*John Wiley & Sons* , USA , **2005**