

Nitroso-R-salt as a sensitive spectrophotometric reagent for the determination of paracetamol in pharmaceutical preparations

*Raghad Sinan**

*Sadeem S. Abed**

Date of acceptance 19 /11 / 2008

Abstract

Nitroso-R-salt is proposed as a sensitive spectrophotometric reagent for the determination of paracetamol in aqueous solution. The method is based on the reaction of paracetamol with iron(III) and subsequent reaction with nitroso-R-salt to yield a green colored complex with maximum absorption at 720 nm. Optimization of the experimental conditions was described. The calibration graph was linear in the concentration range of 0.1 – 2.0 $\mu\text{g mL}^{-1}$ paracetamol with a molar absorptivity of $6.9 \times 10^4 \text{ L mol}^{-1} \text{ cm}^{-1}$. The method was successfully applied to the determination of paracetamol in pharmaceutical preparations without any interference from common excipients. The method has been statistically evaluated with British Pharmacopoeia method and no statistical difference between methods was found at the 95% confidence level.

Key words: Paracetamol; Spectrophotometry; Nitroso-R-salt; Pharmaceutical preparations.

Introduction

Nitroso-R-salt is the disodium salt of α -nitroso- β -naphthol disulfonic acid. This reagent was used for qualitative identification of ferrous ion and for copper and nickel ions at low pH [1].

Paracetamol (acetaminophen) has the chemical name N-(4-hydroxyphenyl) acetamide. It is important and extensively used as antipyretic-analgesic drug [2].

Various methods have been reported for the determination of paracetamol in pharmaceutical preparations. These include spectrophotometry [3-7], polarography [8], micellar electrokinetic chromatography [9], flow injection-spectrofluorimetry [10], flow injection fourier-transform infrared spectrometry [11], flow injection-spectrophotometry [12, 13], first derivative spectrofluorimetry [14],

square-wave voltammetry [15], flow injection-biamperometry [16], high-performance thin-layer chromatography [17], and reversed-phase capillary electrochromatography [18].

Most of spectrophotometric methods need either pre-hydrolysis step [19] or temperature and pH controls [20]. Therefore, development of a simple and sensitive spectrophotometric method seems to be desirable. The present paper describes a simple and sensitive spectrophotometric method for the evaluation of paracetamol in pharmaceutical preparations. The method is based on the reaction of paracetamol with iron(III) and subsequent reaction with nitroso-R-salt to yield a green colored complex. A comparative study of this method with

*Department of Chemistry, College of Science, University of Baghdad, Baghdad - Iraq

the official method reported in pharmacopoeia was made. The present method offers the advantages of simplicity, no need for extraction or heating, in addition to higher sensitivity in comparison with some of the existing spectrophotometric methods.

Materials and methods:

Apparatus

A Shimadzu UV-visible 260 digital double-beam recording spectrophotometer (Shimadzu, Kyoto, Japan) was used for all spectral and absorbance measurements with matched 1-cm quartz cells.

Reagents

All chemicals used were of analytical reagent grade. Paracetamol standard material was provided from the State Company for Drug Industries and Medical Appliances (SDI), Samarra – Iraq.

- 1- Paracetamol stock standard solution $1000 \mu\text{g mL}^{-1}$ was prepared by dissolving 0.1000 g of pure paracetamol in 5 ml of ethanol and diluting to the marked with distilled water in 100 mL volumetric flask. Working standard solutions were prepared by suitable dilution of the stock standard solution.
- 2- Ferric sulfate solution 1 mM was prepared by dissolving 0.0562 g of ferric sulfate in distilled water and diluting to the marked in 100 mL volumetric flask.
- 3- Nitroso-R-salt solution 10 mM was prepared by dissolving 0.3773 g of nitroso-R-salt in distilled water and diluting to the marked in 100 mL volumetric flask.

Pharmaceutical preparations of paracetamol

Pharmaceutical preparations were obtained from commercial sources.

- 1- Paracetol tablets (SDI, Iraq): 500 mg paracetamol for each tablet.
- 2- Panadol Extra tablets (Smithkline Beecham, Ireland): 500 mg paracetamol, 65 mg caffeine for each tablet.
- 3- Panatol tablets (Global Pharma, UAE): 500 mg paracetamol for each tablet.
- 4- Emidol tablets (Global Pharma, UAE): 500 mg paracetamol for each tablet.
- 5- Kanagesic tablets (Kanawati Medical Products, Syria): 450 mg paracetamol and 35 mg orphenadrine citrate for each tablet.
- 6- Hayamol injections (Ibn Hayyan Pharmaceutical HOMS, Syria): 375 mg paracetamol for each injection (5 mL).

Calibration graph

Aliquot of standard paracetamol solution (2.5 – 50.0 μg) was transferred into 25 mL calibrated flasks. To each flask, 1.5 mL of ferric sulfate solution (1 mM), shake well and followed by 3 mL of nitroso-R-salt solution (10 mM). The contents were diluted to the mark with distilled water and mixed; after 45 min, the absorbance value at λ_{max} 720 nm was measured against a reagent blank and a calibration graph was constructed.

Procedure for the assay of pharmaceutical preparations

- 1- Tablets solution ($1000 \mu\text{g mL}^{-1}$)

The average tablet weight was calculated from the contents of 20 tablets that had been finely powdered and weighed. A portion of this powder, equivalent to 100 mg of paracetamol, was accurately weighed, shaken with 5 mL of ethanol, diluted with distilled water in to a 100 mL volumetric flask and was filtered.

- 2- Injections solution ($1000 \mu\text{g mL}^{-1}$)

The contents of five injections were mixed. An aliquot corresponding to

100 mg of paracetamol (1.3 mL) was shaken with 5 mL of ethanol and diluted to 100 mL with distilled water in a volumetric flask.

Further appropriate solutions of pharmaceutical preparations were made by using simple dilution with distilled water. Two different concentrations of each solution of pharmaceutical preparations were analyzed in five replicate using the above procedure.

Results and discussion

The proposed spectrophotometric method for the determination of paracetamol is based on the oxidation reaction of paracetamol with iron(III) and subsequent chelation of iron(II) with nitroso-R-salt to form a green colored product.

Spectral characteristics

A green product is formed when paracetamol was allowed to react with iron(III) salts and subsequent reaction of iron(II) produced with nitroso-R-salt. The soluble colored complex has a maximum absorption at 720 nm as shown in Fig. (1).




Fig. (1): Absorption spectrum of complex of nitroso-R-salt with iron(II) which produced from oxidation reaction of paracetamol ($1.5 \mu\text{g mL}^{-1}$) with iron(III)

Optimum conditions for complex formation

In order to establish the optimum conditions necessary for a rapid and quantitative formation of the colored product with maximum stability and sensitivity, the investigators measured the absorbance of a series of solutions by varying one and fixing the other parameters at 720 nm. It was found that a 1 mM solution of ferric sulfate in the range 0.5 – 4.0 mL and a 10 mM solution of nitroso-R-salt in the range 0.5 – 5.0 mL were necessary to achieve a maximum color intensity of the product. Fig. (2) shows that 1.5 mL of ferric sulfate solution (1 mM) and 3 mL of nitroso-R-salt solution (10 mM) were enough to obtain the maximum absorbance. Therefore, 1.5 mL of ferric sulfate and 3 mL of nitroso-R-salt were recommended for all measurements.





Fig. (2): Optimum conditions for determination of paracetamol

The reaction between paracetamol and iron(III) in the presence of nitroso-R-salt was found to be instantaneous. However, the reaction is complete within 45 min at room temperature (25°C).

The effect of temperature on the colored product was studied at 5, 25 and 45°C and the results obtained indicated that the color was stable for at least 120 min at 25°C and was used in the recommended procedure.

Reaction mechanism

Paracetamol reduces iron(III) salts in aqueous medium to form iron(II) salts, which subsequently chelate with nitroso-R-salt to form a green colored product as given in the following equations [1, 21]:



Analytical parameters

Employing the conditions described under analytical procedure, a linear calibration graph [Fig. (3)] for paracetamol was obtained. It shows that Beer's law was obeyed in the range given in Table (1). The analytical and regression parameters [22] of proposed spectrophotometric method are compiled in Table (1), and demonstrate the highly sensitivity of the method.




Fig. (3): Calibration graph for paracetamol

Table (1): Analytical and regression parameters of proposed method Accuracy and precision

Parameter	Value
λ_{max} , nm	720
Beer's law limits ($\mu\text{g mL}^{-1}$)	0.1 – 2
Regression equation $y = a + b x$; $y = \text{absorbance}$, $x = \text{concentration } (\mu\text{g mL}^{-1})$	$y = 0.0165 + 0.4566 x$
Correlation coefficient, r	0.9980
Correlation of determination, r^2	0.9960
Intercept, a	0.0165
Slope, b ($\text{mL } \mu\text{g}^{-1}$)	0.4566
Standard deviation of the residuals, $S_{y/x}$	0.0230
Standard deviation of the intercept, S_a	0.0135
Standard deviation of the slope, S_b	0.0130
Molar absorptivity, ϵ ($\text{L mol}^{-1} \text{cm}^{-1}$)	6.902×10^4
Sandell's sensitivity, S (ng cm^{-2}) per 0.001 absorbance unit	2.1902
Limit of detection, LOD ($\mu\text{g mL}^{-1}$) $\text{LOD} = 3 S_{y/x} / b$	0.1513
Limit of quantification, LOQ ($\mu\text{g mL}^{-1}$) $\text{LOQ} = 10 S_{y/x} / b$	0.5044

To determine the accuracy and precision of the method, pure paracetamol solution was determined at two different concentrations. The results shown in Table (2), indicate that a satisfactory precision and accuracy could be obtained with the proposed method.

Interference

To test the efficiency and selectivity of the proposed analytical method to pharmaceutical preparations, a systematic study under the optimum experimental conditions was made for the effect of additives and excipients such as lactose, talc, starch, magnesium stearate and polyvinylpyrrolidone (PVP) that are usually present in dosage forms. The criterion of interference was an error of not more than $\pm 1\%$ in the absorbance. In this study, a wide range of concentrations was used in which the determination of the $2 \mu\text{g mL}^{-1}$ level of a drug was performed. Experimental results showed that there was no interference from additives or

excipients for the examined method up to 10-fold excess as shown in Table (3).

Table (2): Accuracy and precision of the proposed method

Concentration of paracetamol ($\mu\text{g mL}^{-1}$)		Erel., %	Rec., %	RSD, %
Taken	Found*			
0.3000	0.2992	- 0.267	99.733	1.273
1.0000	1.0153	+ 1.530	101.530	0.807

Erel. is relative error. Rec. is recovery.
RSD is relative standard deviation.
* Mean value of five determinations.

Table (3): Determination of $2 \mu\text{g mL}^{-1}$ of paracetamol in the presence of excipients

Excipient, $20 \mu\text{g mL}^{-1}$	Conc. of paracetamol, $\mu\text{g mL}^{-1}$	Erel., %	Rec., %	RSD, %
	Found*			
Lactose	2.0037	+ 0.185	100.185	0.612
Starch	1.9920	- 0.400	99.600	0.251
Talc	2.0191	+ 0.955	100.955	0.337
Mg stearate	1.9963	- 0.185	99.815	0.427
PVP	2.0005	+ 0.025	100.025	0.221

* Mean value of five determinations.

Analytical applications

The proposed method was applied successfully to the determination of paracetamol in pharmaceutical preparations [Table (4)]. The results were compared statistically using Student's t-test for accuracy, and a variance ratio F-test for precision [22] with the official method [23] at the 95% confidence level [Table (5)]. The results showed that the t- and F-values were less than the theoretical value, indicating there was no significant difference between both methods in *accuracy*, but the proposed method was *more precise* than the official

method. Consequently, we advise to adopt the proposed method in routine analysis and for quality control purposes for paracetamol and related pharmaceutical preparations.

Conclusion

The proposed method is found to be economical, simple, fairly and highly sensitive than other spectrophotometric methods [Table (6)]. It has the advantage of being accurate, does not require the removal of excipients, temperature control, pH control, solvent extraction step and pre-hydrolysis step. The statistical parameters and recovery study data clearly indicate the reproducibility and accuracy of the proposed method. It can be applied successfully to different pharmaceutical preparations.

Table (4): Determination of paracetamol in pharmaceutical preparations using the proposed method

Pharmaceutical Preparation	Concentration of paracetamol ($\mu\text{g mL}^{-1}$)		Erel., %	Rec., %	RSD, %
	Taken	Found*			
Paracetol tablets	0.5000	0.4925	- 1.500	98.500	0.563
	1.5000	1.4930	- 0.467	99.533	0.269
Panadol Extra tablets	0.5000	0.4957	- 0.860	99.140	0.204
	1.5000	1.5021	+ 0.140	100.140	0.126
Panadol tablets	0.5000	0.4882	- 2.360	97.640	1.570
	1.5000	1.5193	+ 1.287	101.287	0.226
Emidol tablets	0.5000	0.4877	- 2.460	97.540	1.380
	1.5000	1.5321	+ 2.140	102.140	0.350
Kanaganecic tablets	0.5000	0.5005	+ 0.100	100.100	0.505
	1.5000	1.5160	+ 1.067	101.067	0.351
Hayamol injections	0.5000	0.4979	- 0.420	99.580	0.426
	1.5000	1.5021	+ 0.140	100.140	0.370

* Mean value of five determinations.

Table (5): The comparison of the proposed method with standard official method using t- and F-statistical tests

Pharmaceutical preparation	Proposed method		Official method		s	Value	
	Rec.% * (x _i) ₁	($\bar{x}_1 - \bar{X}$) ₁ ²	Rec.% * (x _i) ₂	($\bar{x}_2 - \bar{X}$) ₂ ²		t (theor.)	F (theor.)
<i>Paracetamol pure</i>	99.757	0.0004	100.000	0.3446	1.482	0.409 (2.18)	2.266 (4.284)
<i>Paracetol tablets</i>	99.017	0.5184	98.881	0.2830			
<i>Panadol Extra tablets</i>	99.640	0.0094	100.726	1.7240			
<i>Panadol tablets</i>	99.464	0.0745	99.025	0.1505			
<i>Emidol tablets</i>	99.840	0.0106	99.242	0.0292			
<i>Kanagesic tablets</i>	100.584	0.7174	98.700	0.5084			
<i>Hayamol injections</i>	99.860	0.0151	99.314	0.0098			
	$\bar{X}_1 =$ 99.737	$\Sigma =$ 1.3458	$\bar{X}_2 =$ 99.413	$\Sigma =$ 3.0495	(n ₁ + n ₂ - 2) = 12	(n ₁ - 1) = 6 (n ₂ - 1) = 6	

* Mean value of five determinations.

Table (6): Comparison of the proposed method with other spectrophotometric methods

Reagents	λ_{max} , nm	Linear range, $\mu\text{g mL}^{-1}$	$\epsilon, \text{L mol}^{-1}\text{cm}^{-1}$	Remarks	Ref.
HNO ₃ + H ₂ SO ₄	355	(0.2 - 1) × 10 ³	4.747 × 10 ³	Treated with acetone and KOH solution	24
Cerium(IV) sulfate in 5M H ₂ SO ₄	410	30 - 160	8.112 × 10 ²	Required heating at 80°C for 90 min	25
Ammonium molybdate	670	0.1 - 6	2.600 × 10 ⁴	Required strongly acidic medium	26
Sodium 1,2-naphthoquinone-4-sulfonate and cetyltrimethyl ammonium bromide	570	1 - 20	1.118 × 10 ⁴	Required pre-hydrolysis of paracetamol	27
o-Cresol + NaIO ₄ + ammonium buffer (pH 10)	612	2 - 44	7.024 × 10 ³	Time consuming (50 min)	28
NaNO ₂ + HCl	430	180 - 300	1.038 × 10 ⁴	Treated with NaOH solution in flow injection system	29
Fe(III) + 1, 10-phenanthroline + acetate buffer (pH 4.5)	510	0.5 - 10	3.779 × 10 ⁴	Required heating at 60°C	20
Fe(III) + nitroso-R-salt	720	0.1 - 2	6.902 × 10 ⁴	Most sensitive, rapid and a facile work	Proposed method

References

1. Fritz, F. and Vinzenz, A., 1972. 6th ed., Spot Tests in Inorganic Analysis. Elsevier Publishing Company, Amsterdam, pp. 669.
2. The Merck Index on CD-ROM. 2000. 12th ed., Whitehouse Station, NJ, USA.
3. Romero, J. S. E., Rodriguez, L. A., Alvarezcoque, M. C. G. and Ramisramos, G., 1994. Spectrophotometric determination of phenols by coupling with diazotized 2,4,6-trimethylaniline in a micellar medium. *Analyst*. 119 (6): 1381-1386.
4. Erk, N. and Onur, F., 1997. Simultaneous determination of analgine and paracetamol in tablets by spectrophotometric methods. *Anal. Lett.* 30 (6): 1201-1210.
5. Mohamed, F. A., Abdallah, M. A. and Shammatt, S. M., 1997. Selective spectrophotometric determination of p-aminophenol and acetaminophen. *Talanta*. 44 (1): 61-68.
6. Nogowska, M., Muszalska, I. and Zajac, M., 1999. Simultaneous spectrophotometric determination of acetylsalicylic acid, paracetamol and caffeine in pharmaceutical preparations. *Chem. Anal.* 44 (6): 1041-1048.
7. Sena, M. M. and Poppi, R. J., 2004. N-way PLS applied to simultaneous spectrophotometric determination of acetylsalicylic acid, paracetamol and caffeine. *J. Pharm. Biomed. Anal.* 34 (1): 27-34.
8. Walsh, M. I., Elbrashy, A. M. and Sultan, M. A., 1994. Polarographic behaviour and determination of

- paracetamol and salicylamide after treatment with nitrous acid. *Microchim. Acta.* 113 (1-2): 113-124.
9. Boonkerd, S., Lauwers, M., Detaevernier, M. R. and Michotte, Y., 1995. Separation and simultaneous determination of the components in an analgesic tablet formulation by micellar electrokinetic chromatography. *J. Chromatogr.* 695 (1): 97-102.
10. Pulgarin, J. A. M. and Bermejo, L. F. G., 1996. Flow-injection stopped-flow spectrofluorimetric kinetic determination of paracetamol based on its oxidation reaction by hexacyanoferrate(III). *Anal. Chim. Acta.* 333 (1-2): 59-69.
11. Bouhsain, Z., Garrigues, S., Moralesrubio, A. and Delaguardia, M., 1996. Flow injection fourier-transform infrared spectrometric determination of paracetamol in pharmaceuticals. *Analyst.* 121 (5): 635-639.
12. Bouhsain, Z., Garrigues, S. and Delaguardia, M., 1996. Flow injection spectrophotometric determination of paracetamol in pharmaceuticals by means of on-line microwave-assisted hydrolysis and reaction with 8-hydroxyquinoline (8-quinolinol). *Anal. Chim. Acta.* 330 (1): 59-69.
13. Criado, A., Cardenas, S., Gallego, M. and Valcarcel, M., 2000. Continuous flow spectrophotometric determination of paracetamol in pharmaceuticals following continuous microwave assisted alkaline hydrolysis. *Talanta.* 53 (2): 417-423.
14. Murillo, J. A. and Garcia, L. F., 1996. Application of first derivative fluorescence spectrometry to the simultaneous determination of paracetamol and salicylamide in pharmaceuticals. *Anal. Lett.* 29 (3): 423-438.
15. Zen, J. M. and Ting, Y. S., 1997. Simultaneous determination of caffeine and acetaminophen in drug formulations by square-wave voltammetry using a chemically modified electrode. *Anal. Chim. Acta.* 342 (2-3): 175-180.
16. Danet, A. F., David, V. and David, I., 1998. Acetaminophen determination by flow injection analysis with biamperometric detection. *Rev. Roum. Chim.* 43 (9): 811-816.
17. Argekar, A. P. and Sawant J. G., 1999. Simultaneous determination of paracetamol and mefenamic acid in tablets by HPTLC. *JPC-J. Planar Chromatogr.-Mod. TLC.* 12 (5): 361-364.
18. Pucci, V., Mandrioli, R., Raggi, M. A. and Fanali, S., 2004. Reversed-phase capillary electrochromatography for the simultaneous determination of acetylsalicylic acid, paracetamol, and caffeine in analgesic tablets. *Electrophoresis.* 25 (4-5): 615-621.
19. Criado, A., Cardenas, S., Gallego, M. and Valcarcel, M., 2000. Fast urinary screening for paracetamol using on-line microwave assisted hydrolysis and spectrophotometric detection. *Analyst.* 125 (6): 1179-1183.
20. Abbas, A., Nahid, S. and Ali, R. Z., 2006. Spectrophotometric determination of salicylamide and paracetamol in biological samples and pharmaceutical formulations by a differential kinetic method. *Acta Chim. Slov.* 53: 357-362.
21. Belal, S. F., Elsayed, M. A., Elwalily, A. and Abdine, H., 1979. Spectrophotometric determination of acetaminophen and salicylamide through nitrosation and subsequent chelation. *Analyst.* 104 (1243): 919-927.

22. Miller, J. N. and Miller, J. C., 2000. 4th ed., Statistics and Chemometrics for Analytical Chemistry. Pearson Education Limited, London, pp. 271.
23. British Pharmacopoeia on CD-ROM. 2001. 3rd ed., The Stationery Office Ltd., London.
24. Elkheir, A. A., Belal, S., Elsadek, M. and Elshanwani, A., 1986. Spectrophotometric determination of acetaminophen, oxyphenbutazone and salicylamide by nitration and subsequent complexation reactions. *Analyst*. 111 (3): 319-321.
25. Sultan, S. M., Alzamil, I. Z., Arahman, A. M. A., Altamrah, S. A. and Asha, Y., 1986. Use of cerium(IV) sulphate in the spectrophotometric determination of paracetamol in pharmaceutical preparations. *Analyst*. 111 (8): 919-921.
26. Morelli, B., 1989. Spectrophotometric determination of paracetamol in pure form and in tablets. *J. Pharm. Biomed. Anal.* 7 (5): 577-584.
27. Nagaraja, P., Murthy, K. C. S. and Rangappa, K. S., 1998. Spectrophotometric method for the determination of paracetamol and phenacetin. *J. Pharm. Biomed. Anal.* 17 (3): 501-506.
28. Al-Abachi, M. Q. and Al-Ward, H. S., 2001. Spectrophotometric micro determination of paracetamol in pharmaceutical preparations via oxidative coupling with o-cresol and sodium periodate. *National J. Chem.* 4: 538-548.
29. Knochen, M., Giglio, J. and Reis, B. F., 2003. Flow-injection spectrophotometric determination of paracetamol in tablets and oral solutions. *J. Pharm. Biomed. Anal.* 33 (2): 191-197.

التقدير الطيفي للباراسيتامول في المستحضرات الصيدلانية باستعمال ملح النترزو-R ككاشف طيفي حساس

سديم صبحي عبد*

رغد سنان عبد الستار*

*قسم الكيمياء، كلية العلوم، جامعة بغداد، بغداد - العراق

الكلمات المفتاحية: باراسيتامول، التقدير الطيفي، ملح النترزو-R، المستحضرات الصيدلانية.

الخلاصة

يتضمن البحث تطوير طريقة طيفية حساسة لتقدير الباراسيتامول في المحلول المائي باستعمال كاشف طيفي حساس (nitroso-R-salt). تعتمد الطريقة على تفاعل الباراسيتامول مع أيونات الحديدك، ثم تفاعل أيونات الحديدوز الناتجة مع الكاشف (nitroso-R-salt) لتكوين معقد اخضر يمتلك أقصى امتصاص عند طول موجي 720 نانومتر. تم تثبيت الظروف التجريبية الفضلى للتفاعل. كان مدى الخطية من 0.1 إلى 2.0 مايكروغرام مل⁻¹ باراسيتامول وقيمة الامتصاصية المولارية مساوية إلى 6.9×10^4 لتر مول⁻¹ سم⁻¹. طبقت الطريقة بنجاح لتقدير الباراسيتامول في المستحضرات الصيدلانية، و تم مقارنة النتائج مع الطريقة القياسية المعتمدة من قبل الدستور البريطاني لتقدير الأدوية، و وجد أن الطريقتين القياسية و المقترحة لا تختلفان معنوياً في الدقة و المصدقية.