Study the Spectral Properties of Coumarine -47 Dissolved in Chloroform

Rafah Abdul Hadi Omer * Mohammed T.Hussian ** Lamia KandKalif *

> Received 20, December, 2012 Accepted 11, March, 2014

Abstract:

The spectral propetties (absorption and fluorescence) of Coumarine-47 laser dye have been studied. This type of laser dye belong the Coumarine family and it has dissolved in chloroform at different concentrations (1x10-5, 5x10-5, and 1x10-4 M) at room temperature. The achieved results have been pointed out to increase in the absorption and fluorescence as the concentration increased which are agreements with Beer – Lambert law. These have been also showed an expansion in the spectral range of absorption and fluorescence with a noticed shift in the direction of longer wavelength (Red-shift) with increasing concentration.

The quantum efficiency of the dissolved C_{47} in chloroform has been computed by using the brevious concentrations and their results are as follows (69%, 63%, and 45%) respectively. The radiative and fluorescence lifetime have been also computed as given (6.47ns, 4.67ns, and 2.06ns) and (4,07ns, 3.22ns, and 1.47ns) respectively.

Key words: Coumarin dye, Dye solutions, Spectroscopic properties.

Introduction:

The first demonstration of a dye laser, where the active medium was a solution of dye dissolved in solvent, was reported in 1966 by Sorokin and Lankard[1]. Since then, stimulated emission from many classes of organic compounds has been reported. Some of popular and the most heavily investigated are the rhodamines and coumarins. Lasers dyes are attractive in many respects as lasing media. Their stimulated emissions are tunable throughout their broad fluorescence spectrum. Laser dyes can be used in solution with appropriate solvents or doped into a solid such as the common polymethylmethacrylate polymer, (PMMA). This level of functionality allows their concentration in the host material, and hence their absorption and gain properties, to be easily controlled. Finally, the cost of the active medium, organic dyes, is negligibly small compared to that of inorganic solidstate lasers.

Solid-state dye lasers. first demonstrated by Soffer and McFarland in 1967[2] have many advantages over their liquid counterparts. They are nontoxic, nonflammable, and can be engineered to be rugged and compact by eliminating the dye flow hardware that is necessary in a liquid system. Liquid dye lasers rely on dye flow through the laser cavity to avoid thermal problems and the buildup of photodegraded dye molecules which decrease the lasing efficiency. Dye flow equipment is often bulky and complex. Because of photodegradation, or the need to change the output wavelength range, it is necessary for the dye solution to be changed periodically. As a consequence, significant time must be spent on laser maintenance, and the user may be exposed to large quantities of toxic solvents and dyes. Solid-state

^{*}University of Baghdad .college of science for women – physics department

^{**}University of Baghdad .college of science – physics department

dye lasers do not have a liquid gain media, so liquid circulating equipment is avoided and thermal problems are more easily compensated because there is no convection. Several solid host materials have been studied with polymer matrices being the favorite.

Unfortunately, today's popular molecules organic laser dye experiences irreversible photodegradation during optical pumping. Although laser-dyes have been under use and investigation for decades, many details of the processes responsible for dye stability and maintenance of their lasing properties from photodegradation under high pump repetition rates are still vague. explanations Some suggest the formation of dimers [3] (two mutually bonded dye molecules) or thermal breakdown due to local heat accumulation[4]. In either case, the degradation effect renders dye molecules incapable of contributing to stimulated emission. These degraded molecules can also reabsorb stimulated emission from nondegraded molecules, worsening the effect. The result is that lasing efficiency continuously and permanently decreases over a relatively short time, rendering the dye dopedpolymer useless as a lasing media and limiting the laser's operational lifetime. In applications in which gain medium replacement or movement is not

desirable. the fundamental photostability of the dye molecules is of paramount importance. Attempts have made delay been to the photodegradation process using additives in the polymer matrix [5, 6] or by rotating the pumped media [7, 8, 9], thereby continuously changing the location of excitation, but to my knowledge, no laser dye doped polymer has been found that can perform with the reliability and simplicity needed for long term use in a commercial system.

Theory

Once a molecule has absorbed a photon it can emit a photon upon returning to the electronic ground state from the electronic excited state. As with absorbance. the strength of the transition is governed by the transition moment. The related peaks in turn will be the dominant peaks in the resulting spectrum. Two processes that emit light from a dye molecule are fluorescence, where the emitted photon is from the decay of excited state S₁to the ground state S₀, and phosphorescence, which is a transition from the triplet state T_1 to the electronic ground state S₀. Most ground states are singlet due to the total spin being zero, cancelled out by pairs of electrons with opposite spin. The direct transition from S_0 to T_1 is forbidden since the value of the spin quantum number S would have to change. Similarly a transition from T₁to S_0 is slow due to the same reason, resulting in a slow process called phosphorescence. Triplet-triplet absorption bands tend to heavily overlap the fluorescence region. The triplet state has a long lifetime leading to a buildup of population inversion. However since the transition from S_0 to

However since the transition from S_0 to T_1 is forbidden, it would require very high dye concentrations for lasing. Fluorescence can be used at relatively low dye concentrations for laser applications. If the triplet state is at a lower energy than S_1 , it then competes with the buildup of the excited S1 state, lowering the quantum yield of the fluorescence [4].

Decay to the lowest level of S_1 is very quick provided that the temperature is low enough to prevent thermal excitation to a higher vibronic level within S_1 . The number of excited atoms N where k is the Boltzmann's constant is *Nexcited*= $N_0 \exp [\Delta E = kT]$. At room temperature $kT = 2.15 \times 10^{-2} eV$ so the population density of the excited state will be low.

The longevity of different excited states is important to the usefulness of a dye as a laser medium. The typical time scales for these processes are: 1ps from S_2 to S_1 (non-raditative transition), 10ns from (non-raditative S_1 to T_1 transition), 1µs from T_1 to S₀ (phosphorescence) and 1ns from S_1 to S_0 (fluorescence) (Figure 1) [5].



Fig.(1) Dye molecule energy state [5].

2-1. Fluorescence

Excitation of a molecule from absorption of a photon or thermal energy promotes the molecule to an excited energy state, setting up the possibility of emission of a photon. The excited molecules may lose energy through a non-radiative process to reach the lowest vibrational level of the excited state. Fluorescence is then the radiative loss of energy as a photon is emitted from a molecule in the excited state returning to the ground state. Sincea non-radiative loss of energy occurs between the absorption and subsequent fluorescence, the emitted photon is of lower frequency than the absorbed photon producing a red shift absorbance between the and fluorescence spectra.

2-2. QuantumEfficiency

The fluorescence efficiency Φ_{fm} , indicates the fraction of pump photons that are converted to fluorescence photons per dye molecule, where the maximum value is unity. Fluorescence quantum yield expresses the proportion of excited molecules that deactivate by emitting a photon[6]. The quantum vield of fluorescence is a measure of the rate of nonradiative transitions that compete with the emission of light. The knowledge of fluorescence quantum efficiency of organic dyes and its concentration dependence are essential for selecting efficient laser media [7]. fluorescence efficiency The for different concentrations of C₄₇ in chloroform solvent was determined and their values are listed in table (1).

The spectrum of the molecular fluorescence $F(\dot{v})$ gives the relative fluorescence intensity at wave-number (\dot{v}), this is related to the quantum efficiency by the following equation [8,9]

$$q_{fm} = \int_{0}^{\infty} F(\upsilon') d\upsilon' \quad \dots \quad (1)$$

In order to evaluate absolute quantum efficiency, we have to consider both the radiative and non-radiative processes taking place in the medium, therefore

$$q_{fm} = \frac{K_{fm}}{K_{fm} + \Sigma K_d} = \frac{K_{fm}}{K_{fm} + K_{IC} + K_{ISC}} - --(2)$$

Where K_{fm} fluorescence rate constant, K_{d} nonradiative rate constant, K_{IC} internal conversion rate constant and K_{ISC} internal system crossing.

Since $K_{fm}=1/\ \tau_{fm}$ and $\tau_{f}\!\!=1/\ K_{fm}+\Sigma\ K_{d}$

Therefore

$$q_{fm} = \frac{\tau_f}{\tau_{fm}} = \int_0^\infty F(\upsilon') - \dots (3)$$

Where τ_f is the fluorescence lifetime and τ_{fm} is the radiative lifetime of the excited state to circumscribe the possible reasons for q_{fm} variations. The radiative lifetime being a function of the absorption strength, it is invariant for any molecule unless its absorption spectrum changes. As this, to first order, is not the case in concentration quenching, the mechanism for q_{fm} variations must be changes in τ_f , caused, for example, by changes in the nonradiative lifetime of the excited state [10].

Results and Discussion:

To study the absorption spectrum for the chloroform, it can be shown from figure (2) that it has no absorption at the spectral range of C 47 dye within wavelength range (300 - 500nm). Absorption spectra of C₄₇ dissolved in chloroform with different concentration $(1x10^{-4}, 5x10^{-5} \text{ and } 1x10^{-5} \text{ M})$ are shown in figures (2) ,(3) and(4) respectively. From these figures one deduced C_{47} can that solution absorption spectrum has a wide spectral range at wavelength range (300 -500nm). It has also be shown that maximum absorption appears at higher concentration at a wavelength (393nm) and red shifted by \approx 13nm, while at lower concentration $(1 \times 10^{-5} M)$ in other word at wavelength 380nm this is due to the fact that an increase in concentration produce an increase in number of molecules in volumetric unit which effect the energy state. This increase in concentration produce an increase in perturbation field of the molecules and while at the concentration decreases the relative decrease which is intensity in agreement with Beer - Lambrat law. Table (1) presents the concentration variation with the maximum absorption wavelength (λ_{abs}) and relative intensity.



Fig.(2) Absorption spectra of chloroform

Table (1)The effect of dyeconcentrationvariation on theabsorptionspectra and the relativeintensity

Concentration (mol/liter)	I _{abs} (a.u)	λ_{max} (abs) (nm)	$\Delta\lambda_{abs}$ (nm)	$\Delta \lambda_{abs}(FWHM)(nm)$
1×10^{-5}	20.5	380	50	20
5×10^{-5}	61.5	385	70	25
1×10^{-4}	66.1	393	90	30

The spectral properties of chloroform with C_{47} were studied by recording the fluorescence emission spectra for different concentrations $(1x10^{-4}, 5x10^{-5} \text{ and } 1x10^{-5} \text{ M})$ as shown in figures (2),(3) and (4). The peak wavelength of fluorescence for the chloroform with C_{47} dye shows red shift from that for pure dye. This change may be attribute to the structure of dye, the fluorescence from the excited state dve molecule is reabsorbed by the ground state molecule which shifts the fluorescence peak to lower energies. The fluorescence bandwidth of the dye in chloroform solvent is boarder than that for pure dye, this is due to the fact that the spectral characteristics of the dye depends insolvent on the intermolecular between the dve and chloroform molecules.

The relative intensity was determined as a function of the dye concentrations and the results are tabulated in table (2).It can be shown that the highest relative intensity for the peak wavelength at the highest dye concentration.

Table (2)The effect of dyeconcentrationvariation on thefluorescencespectra and the relativeintensity

Concentration (mol/liter)	I _{flo} (a.u)	λ_{max} (flo) (nm)	$\Delta\lambda_{\rm flo}$ (nm)	$\Delta\lambda_{\rm flo}$ (FWHM) (nm)
1×10^{-5}	18	402	30	25
5×10^{-5}	44.5	405	60	40
1×10^{-4}	64.6	410	70	45



Fig. (3) Absorption and fluorescence spectra of C47 dissolved in chloroform at concentration (1x10-4 M)



Fig. (4) Absorption and fluorescence spectra of C47 dissolved in chloroform at concentration (5x10-5 M) $\,$



Fig. (5) Absorption and fluorescence spectra of C47 dissolved in chloroform at concentration (1x10-5 M)

The fluorescence quantum yield of the dye was determined on the basis of the absorption and fluorescence spectra of the dye dissolved in chloroform. This was calculated by using the following equation:



From the results of calculation by using computer model (Matlab 6.5) and the results tabulated in table (3).

Table	(3)		Rep	rese	nts	quantum
efficienc	eies	of	the	C ₄₇	in	chloroform
solvent						

Concentration	Quantum
(mol/liter)	efficiency% q _{fm}
1x10 ⁻⁵	69%
5x10 ⁻⁵	63%
1×10^{-4}	45%

Conclusion:

From the observed results of studying the fluorescence and absorption spectra of C_{47} dye dissolved in chloroform one can conclude that the increase in dye concentration showed an increase in relative intensity for both spectra, which is in agreement with Beer – Lambrt law. The Stock's shift for fluorescence spectrum of the C_{47} dye dissolved in chloroform is toward the longer wavelength (low energy) as

the concentration increased. It must be noted that the quantum yield of the dye solution decrease the as dve concentration increase. Finally as the concentration dye increase the interference region between the fluorescence and absorption spectra have been increased which produced a decrease in Stock's shift.

References:

[1] M. Maeda, 1984, Laser Dyes, "Properties of Organic Compounds for Dye Lasers", Academic Press, Inc., Orlando, Florida, pp. 1.

[2] K. Kuriki, T. Kobayashi, N. Imai, T. Tamura, S. Nishihara, Y. Nishizawa, A. Tagaya, Y. Koike, Y. Okamoto, 2000, Appl. Phys. Lett. 77, 331.

[3] K.H. Drexhage, 1990," Structure and Properties of Laser Dyes in Topics in Applied Physics, Dye Lasers, Springer-Verlag, Berlin, Germany, pp. 21-22.

[4] F. Amat-Guerri, A. Costela, J.M. Figuera, F. Florido, R. Sastre, 1993, Chem. Phys. Lett. 209, 352.

[5] C. Dudley,2004," Absorption, Fluorescence and Applied Spontaneous Emission of Blue - Emitted Dyes", Thesis, Washington State University. Internet, Practice 4.2,2009, [6] Fluorescence Quantum Yield Fluorescence Quantum Yield of Dyes in Highlighter Pen, No. 9. [7]C.V.Bindhu, S.S. Harilal, V.P.N. Nampoori and C.P.G. Vallabhan, 1999, "Solvent Effect on Absolute Fluorescence Quantum Yield of Rhodamine 6G Determined using Transient Thermal Lens Technique", Modern Physics Letters B, Vol. 13, No. 16, 563.

[8] G. Somasundaram, A. Ramalingam, J. Photochem. Photobiol, 1999, A125, 97.

[9] G. Somasundaram, A. Ramalingam, J. Lumin, 2000," Gain Studies of Coumarin 1 Dye - Doped Polymer Laser". 90, 5.

[10] G. Somasundaram, A. Ramalingam,2000, " Gain Studies of Coumarin 490 Dye – Doped Polymer Laser". Chem. Phys. Lett 324, 28.

لمباع قند خليف*

رفاه عبدالهادي عمر * محمد تقى حسين **

*جامعة بغداد / كلية العلوم للبنات / قسم الفيزياء ** جامعة بغداد / كلبة العلوم / قسم الفيز باء

الخلاصة :

تمت دراسة الخصائص الطيفية (الامتصاص والفلورة) لصبغة الكومرين – 47 الليزرية. وتنتمى هذه الصبغة الليزرية لعائلة الكومرين والمذابة في مذيب الكلوروفورم وبتراكيز (⁴-1x10⁻⁵, 5x10⁻⁵)) مولاري وبدرجة حرارة الغرفة. أظهرت النتانج زيادة في شدة الامتصاص والفلورة مع زيادة التركيز الذي يتفق مع بير – ُلامبرت وكذلك توسيع المدى الطيفي للامتصاصُ والفلورة وحدوث ازاحة لقمَّة طيف الامتصاصُ والفلورة باتجاه الاطوال الموجية الاطول (ازاحة – حمراء) مع زيادة التركيز.

أذتم حساب الكفاءة الكمية لصبغة الكومرين – 7 في مذيب الكلوروفورم وبحسب التراكيز السابقة فكانت النتائج كما يلي: (63% , 63% , 63% ,) على التوالي وكذلك زمن العمر الاشعاعي (6.47, 4.67, 2,06) نانوثانية وزمن عمر الفلورة (1.47, 3.22, 1.47) بنانوثانية وعلى التوالي .