

## Synthesis, Structural and Biological studies of 4-[(1-phenyl-2,3-dimethyl-3-pyrozoline-5-one)azo]-N,N-dimethyl aniline with some transition metal ions

Rehab A.M.Al-Hasani \*

Received 6, May, 2008

Acceptance 12, May, 2009

### Abstract:

The present work includes the preparation and characterization of {Co(II), Ni(II), Pd(II), Fe(III), Ru(III), Rh(III), Os(III), Ir(III), Pt(IV) and VO(IV)} complexes of a new ligand 4-[(1-phenyl-2,3-dimethyl-3-pyrozoline-5-one)azo]-N,N-dimethylaniline (PAD). The product (PAD) was isolated, studied and characterized by physical measurements, i.e., (FT-IR), (UV) Spectroscopy and elemental analysis (C.H.N). The prepared complexes were identified and their structural geometric were suggested in solid state by using flame atomic absorption, elemental analysis (C.H.N), (FT-IR) and (UV-Vis) Spectroscopy, as well as magnetic susceptibility and conductivity measurements. The study of the nature of the complexes formed in (ethanolic solution) following the mole ratio method, gave results which were compared successfully with those obtained from solid state studies. The apparent stability constant of the complexes have been studied with the time and their color were stable for more than (6) hours, as well as the molar absorptivities have been calculated. The antibacterial activity for the ligand (PAD) and their metal complexes were studied against two selected micro-organisms [(*Klesbiella pneumonia*) as gram negative] and [(*Streptococcus faecalis*) as gram positive]. Further more the antifungal activity against two micro-organisms (*Candida albicans* and *Aspergillus flavus*) were studied for the ligand (PAD) and their metal complexes.

**Key words:** Azo compounds, Coordination compound of Azo, Synthesis and crystal structure of Azo compounds, Biological activity of Azo compounds.

### Introduction: -

A series of N(1)-aryloxy (or thio) acetyl-3,5-dimethyl pyrazoles have been prepared in order to study their hypoglycemic and CNS activities [1,2]. A slight reduction of blood sugar was observed when these compounds were screened on rats at an oral dose of 250mg/kg body weight as well as some of these compounds were found CNS stimulant, relatively non-toxic, induced writhing and piloerections in albino mice<sup>(1)</sup>. Several N-substituted (3,5) or (2,3) -dimethyl

pyrazoles have recently been known to display significant hypoglycemic activity [3,5]. Some compounds having N-acyl and N-aryloxyacetyl [1] moieties have also evinced substantial hypoglycemic efficacy.

The present paper describes the preparation of a new azo compound derivative from (1-phenyl-2,3-dimethyl-4-amino-3-pyrozoline-5-one), in an attempt to introduce the azo (-N=N-) moiety in the structure of pyrozole ring to investigate the

\*Al-Mustansirya University, College of Science, Department of Chemistry, Baghdad, Iraq.

coordination behaviour of the new (PAD) ligand toward some transition metal ions {Co(II), Ni(II), Pd(II), Fe(III), Ru(III), Rh(III), Os(III), Ir(III), Pt(IV) and VO(IV)}, which have been chosen to react with (PAD) and to compare the biological activity of (PAD) ligand and their metal complexes with the main ring structure.

## Material and Methods:

### Physical measurements and analysis:-

Melting points were recorded on Gallenkamp melting point apparatus and were uncorrected. FT-IR spectra were recorded using FT-IR.8300 Shimadzu in the range of (4000-200)  $\text{cm}^{-1}$ , samples were measured as (CsI disc). Electronic spectra were obtained using UV-1650 PC Shimadzu Spectrophotometer at room temperature. The measurements were recorded using a concentration of  $10^{-3}\text{M}$  of the (PAD) ligand and their metal complexes in chloroform as a solvent. Microanalytical data for (C.H.N) were obtained using 1108CHN-O. The metal content was estimated Spectrophotometrically using Flame Atomic absorption Shimadzu A.A-670 Spectrophotometer. Conductivity measurements were obtained using Corning Conductivity Meter 220 with cell constant ( $1\text{cm}^{-1}$ ), using DMF as a solvent and concentration of  $10^{-3}\text{M}$  at  $25\text{C}^{\circ}$ . Magnetic susceptibility measurements were obtained at  $25\text{C}^{\circ}$  on the solid state applying Faraday's method using Bruker BM6 instrument.

### A-preparation of the AZO (PAD)ligand

(PAD) was prepared using the general procedure [6]. 1-phenyl-2,3-dimethyl-4-amino-3-pyrozoline-5-one (2.54gm, 0.0142mol) was dissolved in 100ml of water and 10ml of

concentrated hydrochloric acid and diazotized below  $5\text{C}^{\circ}$  with  $\text{NaNO}_2$  (1gm, 0.0142mol). The resulting diazonium chloride solution was mixed with N,N-dimethylaniline (1.72, 0.0142mol) dissolved in alcoholic NaOH solution (3gm, 100ml) below  $0\text{C}^{\circ}$ . The mixture was left in the refrigerator over night, the solid product was filtered off, washed with cold water 250ml and crystallized from ethanol and dried over  $\text{CaCl}_2$ .

### B-Preparation of the metal complexes (I-10)

The metal complexes were prepared by mixing the ethanolic solutions of the (PAD) ligand (0.01mol) with the appropriate metal ion salts [ $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$ ,  $\text{NiCl}_2 \cdot 6\text{H}_2\text{O}$ ,  $\text{PdCl}_2(\text{PhCN})_2$ ,  $\text{FeCl}_3 \cdot 9\text{H}_2\text{O}$ ,  $\text{RuCl}_3 \cdot \text{H}_2\text{O}$ ,  $\text{RhCl}_3 \cdot \text{H}_2\text{O}$ ,  $\text{OsCl}_3 \cdot \text{H}_2\text{O}$ ,  $\text{IrCl}_3 \cdot \text{H}_2\text{O}$ ,  $\text{H}_2\text{PtCl}_6 \cdot 6\text{H}_2\text{O}$  and  $\text{VOSO}_4 \cdot 5\text{H}_2\text{O}$ ], (0.01mol). The pH of the reaction mixture was adjusted to ~8 and it was refluxed for (2-3hr.) on a water bath. The metal chelates precipitated on refluxing the reaction mixture were filtered, washed thoroughly with hot water then ethanol several times, till the filtrate was colourless. The colored precipitates were finally washed with pet. ether ( $60-80^{\circ}$ ) and then dried under vacuum.

### C--Study of complex formation in solution

Complexes of (PAD) with metal ions were studied in solution using ethanol as a solvent, in order to determine [M : (PAD)] ratio in the complex following Molar ratio method [7]. A series of solutions were prepared having a constant concentration [10-3M] of the metal ion and (PAD). The [M : (PAD)] ratio was determined from the relationship between the absorption of the absorbed light and the mole ratio of [M : (PAD)].

**D- Stability constant of azo complexes**

The stability constant (**K**)[7] of the (1:1) or (1:2) [Metal: (PAD)] complex were evaluated as follows: Two sets of solutions were prepared, the first set of solutions were formulated to contain stoichiometric amount (1ml) of ( $10^{-3}$ M) ligand to (1 ml) of ( $10^{-3}$ M) of metal ion by placing in to a three series of 10ml volumetric flasks. The solutions of the coloured complexes were diluted to the mark with ethanol(**As**). The second set were formulated to contain five fold excess (5 ml) of ( $10^{-3}$ M) ligand, by placing in to a three series of (10ml) volumetric flasks followed by addition of (1ml) of ( $10^{-3}$ M) of metal ion solution, the volumes were then completed to the mark with ethanol(**Am**). The absorbance (**As** and **Am**) of the solutions, were measured at  $\lambda_{\text{max}}$  of maximum absorption . The stability constant (**K**) and the molar absorptivity ( $\epsilon_{\text{max}}$ ) have been calculated.

**E- Biological activity for (PAD) and its complexes**

The biological activity of the prepared (PAD) and their respective complexes were studied against selected types of bacteria which include[(*Klesbiella pneumonia*) as gram negative] and[(*Streptococcus faecalis*) as gram positive], were cultivated in Nutrient agar medium, as well as DMSO was used as a solvent and as a control , the concentrations of the compounds in this solvent were  $10^{-3}$ M, by using **Disc Sensitivity Test** [8,9], this method involves the exposure of the zone of inhibition toward the diffusion of micro-organism on agar plate. The plates were

incubated for 24hr. at 37 °C, the zone of inhibition of bacterial growth around the disc was observed.

In order to complete this study, the new (PAD) ligand and their metal complexes were tested for their *in vitro* growth inhibitory activity against a pathogenic fungi, i.e., (*Candida albicans* and *Aspergillus flavus*) on **Potato dextrose agar** medium and incubated at 30 C° for 72 hr., DMSO was used as a solvent and as a control, for both techniques, the concentration of the compounds in this solvent were  $10^{-3}$ M. The inhibition of fugal growth, expressed in percentage terms, were determined on the growth in test plates compared to the respective control plates, as given by the **Vincent equation** [8-10], Table (5).

**Results and Discussion:-****(A)-Elemental Analysis :-**

The analytical and physical properties of the (PAD) and its metal complexes are summarised in Table (1), indicate [1:1] [Metal:Ligand] stoichiometry in case [1- 4 and 10] complexes except [5-9] complexes where the molar ratio are [1:2] .The new (PAD) ligand was soluble in common organic solvents such as (ethanol, acetone and methanol) whereas the [1-10] of the new azo coloured crystallin solids complexes were soluble in ( $\text{CH}_2\text{Cl}_2$ ,  $\text{CHCl}_3$ , DMF and DMSO) They are thermally stable and unaffected by atmospheric oxygen and moisture . The elemental analyses data are in agreement with the proposed stoichiometry.

Table (1) Physical data for (PAD) and its metal complexes

Comp. No.	Color	Melting Point C°	Yield %	Elemental analyses				Suggested Formula for isolated precipitate
				Found (Calc.) %				
				C	H	N	M	
(PAD)	Orange	187-189	95	68.34 (68.05)	5.93 (6.27)	20.07 (20.89)	-	C <sub>19</sub> H <sub>21</sub> N <sub>5</sub> O
[1]	Greenish-Blue	208	84	46.95 (47.30)	4.58 (4.35)	15.26 (14.52)	12.33 (12.02)	[Co (PAD)Cl <sub>2</sub> ] . H <sub>2</sub> O
[2]	Reddish brown	211	72	44.71 (45.53)	3.44 (4.19)	13.82 (13.98)	10.77 (11.72)	[Ni (PAD)Cl <sub>2</sub> ] . 2 H <sub>2</sub> O
[3]	Brown	215	86	43.61 (44.49)	4.23 (4.09)	12.24 (13.66)	19.82 (20.76)	[Pd (PAD)Cl <sub>2</sub> ]
[4]	Dark brown	219	70	25.04 (25.21)	3.60 (2.32)	7.11 (7.74)	6.49 (6.17)	[Fe (PAD)Cl <sub>2</sub> (H <sub>2</sub> O) <sub>2</sub> ] . Cl.2H <sub>2</sub> O
[5]	Dull grey	233	78	52.88 (51.96)	4.13 (4.78)	16.27 (15.95)	10.69 (11.51)	[Ru (PAD) <sub>2</sub> Cl <sub>2</sub> ] Cl
[6]	Yellowish-red	230	80	50.07 (50.81)	3.91 (4.68)	15.41 (15.60)	11.58 (11.46)	[Rh (PAD) <sub>2</sub> Cl <sub>2</sub> ] Cl. H <sub>2</sub> O
[7]	Blackish brown	238	72	43.77 (44.67)	4.02 (4.11)	12.89 (13.71)	18.13 (18.63)	[Os (PAD) <sub>2</sub> Cl <sub>2</sub> ] Cl. 3H <sub>2</sub> O
[8]	Orangish-red	232	78	43.21 (42.99)	3.17 (3.95)	3.84 (3.95)	18.01 (18.12)	[Ir(PAD) <sub>2</sub> Cl <sub>2</sub> ] Cl. 2 EtOH
[9]	Dark red	238	81	46.19 (45.27)	4.24 (4.17)	14.46 (13.90)	19.31 (19.37)	[Pt (PAD) <sub>2</sub> Cl <sub>2</sub> ] Cl <sub>2</sub>
[10]	Dark green	220	83	45.68 (45.78)	3.36 (4.21)	14.29 (14.05)	11.12 (10.23)	[VO (PAD) SO <sub>4</sub> ]

**(B)-Infra red spectra :-**

The FT-IR.spectrum of the free (PAD) ligand was compared with the spectra of metal complexes and data is shown in Table (2).A strong bands is observed in the free (PAD) ligand at (1568cm<sup>-1</sup>) and (1775cm<sup>-1</sup>), which are assigned to  $\nu\text{N}=\text{N}$  and  $\nu\text{C}=\text{O}$  respectively [11,12].Coordination of the (PAD) ligand to the metal through the azo nitrogen atom and oxygen atom of the carbonyl group are expected to reduce the electron density in the azo and carbonyl links and lower the  $\nu\text{N}=\text{N}$  and  $\nu\text{C}=\text{O}$  absorptions frequency.In the spectra of all the new azo complexes ,the bands due to  $\nu\text{N}=\text{N}$  and  $\nu\text{C}=\text{O}$  were shifted to lower frequencies (1550-1556) cm<sup>-1</sup> and

(1763-1768) cm<sup>-1</sup> respectively as is shown in table (2). On the basis of this evidence, it is concluded that (PAD) acts as **bidentate** ligand in all the complexes . In the far-infrared region the azo complexes exhibited bands around (513 - 520) cm<sup>-1</sup>, (460 - 508) cm<sup>-1</sup> and (370 - 405) cm<sup>-1</sup>, which are assignable to  $\nu(\text{M} - \text{O})$ ,  $\nu(\text{M} - \text{N})$  and  $\nu(\text{M} - \text{Cl})$  modes respectively[13]. A strong band was observed at (978) cm<sup>-1</sup> due to ( $\nu \text{V} = \text{O}$ ) stretching mode in [10] complex[13], furthermore a band related to ( $\text{SO}_4$ ) sulfate anion in the [10] complex was observed at (1504) cm<sup>-1</sup>, which indicated a bidentate behavior[13,14].

Table (2) Characteristic Stretching Vibrational Frequencies (cm<sup>-1</sup>) located in the FT-IR of (PAD) and their metal complexes

Comp.No.	$\nu\text{N}=\text{N}$	$\nu\text{C}=\text{O}$	$\nu\text{M}-\text{O}$	$\nu\text{M}-\text{N}$	$\nu\text{M}-\text{Cl}$	Others
(PAD)	1568(s)	1775(s)	-	-	-	-
[1]	1553(m)	1763(m)	514(mw)	483(mw)	387(w)	3453( $\nu\text{O}-\text{H}$ of water)
[2]	1553(m)	1766(m)	516(mw)	488(mw)	401(w)	3456( $\nu\text{O}-\text{H}$ of water)
[3]	1556(m)	1763(m)	513(mw)	504(mw)	405(w)	-
[4]	1550(m)	1765(m)	520(mw)	508(mw)	398(w)	1615 of coordinated H <sub>2</sub> O 3456( $\nu\text{O}-\text{H}$ of water)
[5]	1552(m)	1763(m)	518(mw)	495(mw)	400(w)	-
[6]	1550(m)	1766(m)	516(mw)	508(mw)	403(w)	3458( $\nu\text{O}-\text{H}$ of water)
[7]	1554(m)	1766(m)	519(mw)	506(mw)	405(w)	3458( $\nu\text{O}-\text{H}$ of water)
[8]	1552(m)	1764(m)	516(mw)	508(mw)	400(w)	3511( $\nu\text{O}-\text{H}$ of water)
[9]	1555(m)	1768(m)	520(mw)	504(mw)	405(w)	-
[10]	1550(m)	1766(m)	518(mw)	500(mw)	-	1504 of (SO <sub>4</sub> )anion coordination

Where:-(s)=strong, (m)=medium, (mw)=medium weak, (w)=week

**(C)- Electroinc spectra , magnetic susceptibility and conductivity measurements :-**

The electronic spectrum of the (PAD) ligand {4-[(1-phenyl-2,3-dimethyl-3-pyrozoline-5-one)azo]-N,N-dimethyl aniline} shows an intense band at (40816)  $\text{cm}^{-1}$  with a weak shoulder at (36680)  $\text{cm}^{-1}$  followed by a strong intense band at (31125)  $\text{cm}^{-1}$  and (29411)  $\text{cm}^{-1}$ . The first three bands may be due to ( $\pi \rightarrow \pi^*$ ) transitions, while the fourth band is due to ( $n \rightarrow \pi^*$ ) transitions which arises due to greater delocalisation of the (C=C, C=O and N=N)  $\pi$  - electron resulting in a bathochromic shift [12,15].

The electronic spectra of the [1-10] metal complexes solutions in chloroform were recorded in the range (200 - 1100) nm and the data of these complexes are reported in Table(3).

[1]:-The greenish-blue cobalt(II) complex gave a magnetic moment value of (4.53)B.M, which indicates a high-spin type complex. Electronic spectrum in chloroform solvent exhibited a splitted band in the range of (17452-14184)  $\text{cm}^{-1}$  [16,17], Fig. (1). These bands can be assigned to the transition  ${}^4A_2 \rightarrow {}^4T_{1(g)}(v_3)$ . A broad band was observed at 3244  $\text{cm}^{-1}$  in infrared spectrum can be assigned to the  ${}^4A_2 \rightarrow {}^4T_{2(g)}(v_1)$ , while the transition of ( $v_2$ ) expected in the range (5000 - 6000)  $\text{cm}^{-1}$  can not be measured [18]. The various ligand field parameters ( $10Dq$ ,  $B$  and  $v_2$ ) have been calculated by referred to Tanaba-Sugano diagram for ( $d^7$ ) configuration [19-21], to be (3244, 720.8 and 5453) respectively, as well as the calculation of the spin-orbit coupling constant ( $\lambda$ ) was calculated. The resulting value ( $\lambda = -198.4$ ) show the present complex to be distorted tetrahedral [20,22]. The nephelauxetic factor ( $\beta$ ) was calculated and found to be (0.64) indicating high degree of

covalence in bonding of ligand donor atoms with cobalt (II) ion [21,22]. The molar conductance showed that the complex was nonelectrolyte, table(3).

[2]:-The nickel (II) complex is diamagnetic suggesting square planar geometry. The electronic spectrum of Ni(II) complex shows two main absorption bands, at (18520)  $\text{cm}^{-1}$  and (25000)  $\text{cm}^{-1}$ , which are assigned to  ${}^1A_{1g} \rightarrow {}^1A_{2g}$  and  ${}^1A_{1g} \rightarrow {}^1B_{1g}$  transition respectively, in a square planar disposition [17,19,23-25]. Conductivity measurement in (DMF) showed that the complex was non conducting behavior, table (3).

[3]:-The prepared brown Pd (II) complex showed a strong charge transfer band at (29630)  $\text{cm}^{-1}$ , as well as two ligand field bands at (22580)  $\text{cm}^{-1}$  and (26334)  $\text{cm}^{-1}$  which may be assigned to :-  ${}^1A_{1g} \rightarrow {}^1B_{1g}$  and  ${}^1A_{1g} \rightarrow {}^1E_g$ , respectively, in a square planar environment [19,22,26]. The magnetic moment value was (0.82)B.M. Conductivity measurement in (DMF) showed that the complex was non ionic, table (3).

[4]:-The Fe(III) complex displays three main absorption bands which are assigned to :-  ${}^6A_{1g} \rightarrow {}^4T_{1g}$ ,  ${}^6A_{1g} \rightarrow {}^4T_{2g}$ , (PAD)  $\rightarrow$  Fe (C.T) transition [16,19,23] respectively. The ligand field parameters calculated [19,21,24] for this complex are as follows : ( $10Dq = 6634 \text{cm}^{-1}$ ), ( $B = 620 \text{cm}^{-1}$ ) and ( $\beta = 0.45$ ). The magnetic moment is (5.83)B.M. with five unpaired electrons and an octahedral configuration [23,24]. Conductivity measurement showed that the complex was to be ionic, table(3).

[5]:- The spectrum of this complex exhibited four absorption bands at (15898, 19708, 27568 and 35144)  $\text{cm}^{-1}$ . These bands correspond to those of octahedral Ru (III) complexes [19,21,26-28], and were assigned to :-

${}^2T_{2g} \rightarrow {}^4T_{1g}$  ,  ${}^2T_{2g} \rightarrow {}^4T_{2g}$  ,  ${}^2T_{2g} \rightarrow {}^2E_g$  and  ${}^2T_{2g} \rightarrow {}^2A_{1g}$  transitions [16,19] respectively. The value of ( $10Dq$  and  $\mu_{\text{eff}}$ ) came out (27568 and 1.69) respectively, revealing inner-orbital low-spin distorted octahedral stereochemistry of the ligand around Ru(III) ion [29]. Conductivity measurement showed that complex was ionic, table (3).

[6]:-Electronic spectrum of the yellowish – red [6] complex in chloroform solution, showed four absorption bands, the first band as a weak shoulder at (18203)  $\text{cm}^{-1}$ , a second band as a broad one at (20812)  $\text{cm}^{-1}$  and the last two absorption bands as a shoulder at (23122) and (25068)  $\text{cm}^{-1}$ , which might be assigned to the transitions :-  ${}^1A_{1g} \rightarrow {}^3T_{1g}$  ,  ${}^1A_{1g} \rightarrow {}^3T_{2g}$  ,  ${}^1A_{1g} \rightarrow {}^1T_{1g}$  and  ${}^1A_{1g} \rightarrow {}^1T_{2g}$  respectively, which came in accordance with the published data for octahedral **Rh(III)** complexes [22,26,28]. The band at 23122  $\text{cm}^{-1}$  can be taken as ( $10Dq$ ) value [18,19,22]. Magnetic moment of solid complex, table(3), showed a higher orbital contribution. Conductivity measurement in DMF showed that the complex was ionic, table (3).

[7]:-The spectral behavior of this complex, table(3), is identical with that of [4] complex, which agree with octahedral geometry around **Os(III)** complex [19,24,30]. The magnetic moment that was measured at room temperature was found to be (1.98)B.M. this value refer to high spin complex, which is the only case in octahedral environment [24,30]. The conductance measurement indicate that the complex to be ionic, table (3).

[8]:-The electronic spectrum of the prepared orange-red **Ir(III)** complex in chloroform solution, showed three absorption bands at (17625, 22280 and 25461)  $\text{cm}^{-1}$ , which might be assigned to the transitions :-  ${}^1A_{1g} \rightarrow {}^3T_{1g}$  ,

${}^1A_{1g} \rightarrow {}^1T_{1g}$  and  ${}^1A_{1g} \rightarrow {}^1T_{2g}$  respectively, these indicate an octahedral geometry [19,31]. The different ligand parameters ( $10Dq$ ,  $B$  and  $\beta$ ) have been calculated using **Tanaba-Sugano** diagram of ( $d^6$ ) system and found to be (24160, 270 and 0.41) respectively [19-21]. The observed high magnetic moment value (1.83B.M.) of the present Ir(III) ( $d^6$ ) configuration, which agree with octahedral geometry around Ir(III), this result indicate a higher orbital contribution around metal ion [19,31]. Conductivity measurement in DMF showed that the complex was to be ionic, table (3).

[9]:-The spectral and magnetic moment behaviors of this complex is identical with that of [6] and [8] complexes, which agree with octahedral geometry around **Pt(IV)** complex [18,19,21], table(3). Conductivity measurements in DMF (182.18  $\mu\text{s}\cdot\text{cm}^{-1}$ ) showed electrolytic nature of 1:2.

[10]:-The electronic spectrum of vanadium (IV) complex, showed three prominent (d-d) absorption bands, the first one as a strong broad band at (11130)  $\text{cm}^{-1}$ , and the second and third bands as a weak shoulders at (19053 and 24112)  $\text{cm}^{-1}$ , these bands can be attributed to the following transitions :-  ${}^2B_{2g} \rightarrow {}^2E_g$  ,  ${}^2B_{2g} \rightarrow {}^2B_{1g}$  , and  ${}^2B_{2g} \rightarrow {}^2A_{1g}$  respectively [14,16,19,22]. The magnetic moment (1.98B.M) is higher than spin value of the vanadium metal only, this result indicates a higher orbital contribution [19,22]. These transitions and magnetic moment value came in accordance with the published data for **square pyramidal** VO(IV) complexes [14,19,22]. Conductivity measurement in DMF showed that the complex was non-ionic, Fig. (2), table(3).




Fig. (1): -UV-Vis.- Spectrum of [1] complex




Fig. (2): -UV-Vis.- Spectrum of [10] complex

**Table (3) Electronic Spectra (CHCl<sub>3</sub>), Conductance (in DMF), and Magnetic moment (B.M) for Azo metal complexes**

Comp No.	Bands cm <sup>-1</sup>	Assignment	Molar cond. $\mu$ S.cm <sup>-1</sup> .mol <sup>-1</sup>	$\mu$ eff. B.M	Suggested Structure
[1]	3244 5453 (cal) 15733(av.)	${}^4A_2 \longrightarrow {}^4T_2$ (F) ${}^4A_2 \longrightarrow {}^4T_1$ (F) ${}^4A_2 \longrightarrow {}^4T_1$ (P)	13.02	4.53	Tetrahedral
[2]	18520 25000	${}^1A_{1g} \longrightarrow {}^1A_{2g}$ ${}^1A_{1g} \longrightarrow {}^1B_{1g}$	18.32	0.36	Square planer
[3]	22580 26334 29630	${}^1A_{1g} \longrightarrow {}^1B_{1g}$ ${}^1A_{1g} \longrightarrow {}^1E_g$ (PAD) $\longrightarrow$ Pd (C.T)	11.68	0.14	Square planer
[4]	15084 17811 29344	${}^6A_{1g} \longrightarrow {}^4T_{1g}$ ${}^6A_{1g} \longrightarrow {}^4T_{2g}$ (PAD) $\longrightarrow$ Fe (C.T)	68.58	5.83	Octahedral
[5]	15898 19708 27568 31144	${}^2T_{2g} \longrightarrow {}^4T_{1g}$ ${}^2T_{2g} \longrightarrow {}^4T_{2g}$ ${}^2T_{2g} \longrightarrow {}^2E_g$ ${}^2T_{2g} \longrightarrow {}^2A_{1g}$	70.15	1.69	Octahedral
[6]	18203 20812 23122 25068	${}^1A_{1g} \longrightarrow {}^3T_{1g}$ ${}^1A_{1g} \longrightarrow {}^3T_{2g}$ ${}^1A_{1g} \longrightarrow {}^1T_{1g}$ ${}^1A_{1g} \longrightarrow {}^1T_{2g}$	70.15	1.69	Octahedral
[7]	16811 18238 31480	${}^6A_{1g} \longrightarrow {}^4T_{1g}$ ${}^6A_{1g} \longrightarrow {}^4T_{2g}$ (PAD) $\longrightarrow$ Os (C.T)	75.39	1.98	Octahedral
[8]	17625 22280 25461	${}^1A_{1g} \longrightarrow {}^3T_{1g}$ ${}^1A_{1g} \longrightarrow {}^1T_{1g}$ ${}^1A_{1g} \longrightarrow {}^1T_{2g}$	70.73	1.83	Octahedral
[9]	18998 24023 30158	${}^1A_{1g} \longrightarrow {}^3T_{1g}$ ${}^1A_{1g} \longrightarrow {}^1T_{2g}$ (PAD) $\longrightarrow$ Pt (C.T)	182.18	2.04	Octahedral
[10]	11130 19053 24112	${}^2B_{2g} \longrightarrow {}^2E_g$ ${}^2B_{2g} \longrightarrow {}^2B_{1g}$ ${}^2B_{2g} \longrightarrow {}^2A_{1g}$	10.25	1.98	Square pyramideal

**Molar ratio and Stability constant for Azo metal complexes :-**

The molar ratio method was follow to determine the [M:(PAD)]

ratio . The results of complex in ethanol as a solvent, table (4), suggest that the metal to ligand ratio was [1:1] for [1-4 and 10] complexes, while [1:2]

for [5-9] complexes, which were comparable to those obtained from solid state study, table(1).

Chelate stability constant (**K**) {eq.(1) and (2)} ,as well as molar absorptivity ( $\epsilon_{max}$ ) {eq.(3)} for all complexes were obtained spectrophotometrically by using the following equations :-

$$K = (1 - \alpha / \alpha^2 C) \text{ Where } [1:1] \text{ of } [M:(PAD)] \dots (1)$$

$$K = (1 - \alpha / 4 \alpha^3 C) \text{ Where } [1:2] \text{ of } [M:(PAD)] \dots (2)$$

$$A_m = (\epsilon_{max} bC) \dots (3)$$

The degree of dissociation ( $\alpha$ ) for the complex is obtained from the relation ship (4).

$$\alpha = (A_m - A_s / A_m) \dots (4)$$

Where (**As**) and (**Am**) are the absorption of the partially and fully formed complex respectively at optimum concentration and ( $\lambda_{max}$ ) of maximum absorption, Fig. (3), table (4).

The results in Table(4) ,show that mole ratio of (1:2) for [5-9] complexes yielded rather high, this probably due

to the presence of a bulk aromatic groups and a higher oxidation state of the metal ions. As well as the developed color for all complexes become stable after one hour up to six hours.




Fig. (3): - Mole ratio plot of (PAD) complexes (5 and 9)

Table (4) Molar ratio, Stability constant and Molar absorptivities of Azo metal complexes at room temperature


Complex	[M:(PAD)]	$A_s$	$A_m$	$K$ $L. mol^{-1}$	$\epsilon_{max}$ $L. mol^{-1} cm^{-1}$	$\alpha$	$\lambda_{max}$ nm
[1]	1:1	0.278	0.308	$9.59 * 10^5$	3081	0.097	571
[2]	1:1	0.373	0.428	$6.32 * 10^{10}$	4281	0.128	660
[3]	1:1	0.398	0.453	$6.00 * 10^5$	4532	0.121	403
[4]	1:1	0.356	0.409	$2.42 * 10^5$	4093	0.130	486
[5]	1:2	0.495	0.551	$2.18 * 10^{10}$	5511	0.101	408
[6]	1:2	0.294	0.336	$4.48 * 10^{10}$	3363	0.125	432
[7]	1:2	0.355	0.403	$1.62 * 10^{10}$	3090	0.119	443
[8]	1:2	0.259	0.309	$5.03 * 10^9$	3106	0.161	418
[9]	1:2	0.374	0.418	$1.93 * 10^{10}$	6682	0.180	515
[10]	1:1	0.324	0.378	$4.25 * 10^5$	5720	0.3785	756

Structure of [PAD] and its Complexes [1-10]:-


According to the results obtained ,

the structure of the above mentioned [PAD] and [1-10] their complexes can be illustrated as follow :-







Ligand [ PAD ]




Complex [ 1 ]




Complexes [ 2,3 ]



Complex [ 4 ]



Complexes [ 5,6,7,8, and 9 ]



Complex [ 10 ]

**Antimicrobial activities:-**

The azo (PAD) ligand and [1-10] their metal chelates were screened *in vitro* in order to evaluate their

antibacterial action against [*Klesbiella pneumonia*] as gram negative and [*Streptococcus faecalis*] as gram

positive] species as well as antifungal activity against two micro-organisms (*Candida albicans* and *Aspergillus flavus*) ,table (5) at two different concentrations. The results from these studies ,showed the followings:-

1) All the **azo metal complexes** are more toxic as compared with their [PAD] parent ligand against the same micro-organism and under the identical experimental conditions , table (5).

2) The increase in the antibacterial and antifungal activities of metal chelates may be due to the effect of the metal ion on the normal cell process . The toxicity also increases with the increasing concentration of test solution containing the new azo complexes .These activities may be explained by **Tweedy,s Chelation Theory** [34,35] .

3). The [1,2 and 10] complexes showed a moderate activities ,while [3-9] complexes exhibited a great enhancement of activity against types of bacteria, Table (6). This difference in synergistic effect between the metal ion and the ligand molecule may be attributed to many factors, of these in that [3-9] complexes are considered to be a hard metal ions making their complexes to be less lipophilic, which relatively retard their permeation through the lipid part of the cell membrane. On the other hand the [1 and 2] are considered to be soft metal ions, which render their complexes to be more lipophilic and will facilitate the penetration through the cell wall and affect the environment of cells.

**Table (5) Antibacterial and antifungal activities data for the Azo metal complexes and their parent (PAD) ligand**

Compo.	<i>Klesbiella pneumonia</i>		<i>Streptococcus faecali</i>		<i>Aspergillus flavus</i>		<i>Candida albicans</i>	
	100 ppm	200 ppm	100 ppm	200 ppm	100 ppm	200 ppm	100 ppm	200 ppm
Control (DMSO)	-	-	-	-	-	-	-	-
(PAD)	6	10	-	6	35	27	40	30
[1]	6	12	8	10	30	18	20	15
[2]	8	12	10	16	31	20	25	20
[3]	10	16	8	10	27	18	23	14
[4]	6	14	10	16	24	12	20	11
[5]	6	18	6	10	22	16	27	15
[6]	8	10	6	12	30	18	22	16
[7]	6	8	8	18	25	12	31	23
[8]	8	12	10	16	22	16	18	13
[9]	10	16	6	20	20	18	20	11
[10]	6	8	8	10	30	23	21	18
Where :[6-8: (+),8-10: (++) ,>10: (+++)]					30-40: (+++) , 20-30 :(+++) , 10-20: (++++)			

## References:-

- Husain,M.I. and Srivastava,G.C., 1979. «*Synthesis of N-aryloxy (or thio) acetyl acetamido pyrazoles*» ,J.Ind.Chem.Soc.,LVI(3),14-18.
- Siddiqui,Z.A., 1990.«*Synthesis and Characterization of the complexes of 5-amino-4-cyanopyrazole with Cr(III), Mn(II), Zn(II),Cd(II) and Hg(II)*»,Ind.J.Chem.,29A(1),111-117.
- Cotton,F.A. and Wilkinson,G., 1986. «*Advanced Inorganic Chemistry*» ,4<sup>th</sup>Ed.,Wiley Eastern Private Ltd.,pp198.
- Eichhorn,G.L. and marzilli, L.G., 1998. «*Advanced Inorganic Biochemistry,models in inorganic chemistry*»,PTR prentice-Hall,Inc., pp92.
- Al-Hamadani, A.A.S., 2005. «*Synthesis and Characterization of Co(II),Ni(II), Cu(II) and Zn(II) Schiff base complexes with o-hydroxy benzylidene-1-phenyl-2,3-dimethyl-4-*

- amino-3-pyrazolin-5-one* », J. Um-Salama for Sci.,12(2),66-70.
6. Vogel,A.I., 1956. «*A Text book of practical organic chemistry* », 3<sup>rd</sup>Ed.,Longman,pp62.
  7. Skoog,D.A., 1988. «*Fundamentals of analytical chemistry*», 5<sup>th</sup> Ed., John-Wiely and Sons, New York, London, pp186
  8. Collins,C.H.,and Lyne,P.M., 1970. «*Microbiological methods*»,3<sup>rd</sup>Ed., Butter Worth and CO.Ltd.,pp203.
  9. Atlas,M. R. , Alfres, E. ,Alfres, B., and Lawrence, C.P., 1995. «*Laboratory manual Experimental Microbiology*», Mosby- year Book, Inc., pp96.
  10. Mishra,L.,and Singh,V.K., 2000. «*Synthesis and antimicrobial studies of some metal complexes*», Ind.J. Chem.,32(4),98-104.
  11. Baker,A.J.,and Cairns,T., 1967. «*Spectroscopic techniques in organic chemistry*», Vol.(2), Heyden and Son.Ltd.,pp233.
  12. Silverstein ,R.M., Bassler, G.C., and Morrill, T.G., 1981. «*Spectrometric Identification of organic Compounds*»,4<sup>th</sup> Ed. , John-Wiely and Sons,Inc., New York, London, pp188.
  13. Nakamoto,K., 1997. «*Infrared of Inorganic and Coordination Compounds*», 6<sup>th</sup> Ed. , John-Wiely, Inc., New York, London, pp245.
  14. Kumar ,U., Singh ,C.,and Mishra, K. , 1988.«*Oxovanadium (IV) complexes with thiohydrazide derivatives* », J.Ind.Chem. Soc., LVII,76-81.
  15. Al-Adely,K.J., 2000. «*Synthesis and spectroscopic study of group (IIB) and (IIIB) metal ion complexes with new azo compounds and with 1-(2-pyridylazo)2-naphthole*», Ph. D. Thesis, Baghdad university, Iraq.
  16. Carl Ballhausen, I., 1962. «*Introduction to ligand Field Theory*», New York,Toronto-London, pp157.
  17. Nicholis, D., 1984. «*Complexes of First-Row transition elements*», Translated by Dr.W.I.Azeez,Mosul university, Iraq, pp89.
  18. Jassim,A.H., 1993. «*Synthesis and study of transition metal Complexesof chelating heterocycles towards the design of Biologically active compounds* », Ph. D. Thesis, Al- Nahrain university, Baghdad, Iraq.
  19. Figgis,B.N., 1966. «*Introduction to ligand Field*», John-Wiely and Sons,Inc.,New York, London, pp166.
  20. Sutton,D., 1968. «*Electronic Spectra of Transition metal Complexes*», Mc Graw-Hill publishing Compony ltd ,pp89.
  21. Greenwood ,N.N.,and Earnshaw,A. , 1998. «*Chemistry of the Elements*», 2<sup>nd</sup> Ed., Pergaman Press,pp1199.
  22. Al-Hasani, R.A.M., 2004. «*Synthesis and study of some transition metal Complexes with mannich bases derived from-1,2,4-triazole ring* », Ph. D. Thesis, Al-Nahrain university, Baghdad, Iraq.
  23. Lever,A.B.P., 1968. «*Inorganic Electronic Spectroscopy*», Elsevier Publishing Company New York, London, pp183.
  24. Rao,P.,and Ganorkar,M., 1988. «*Synthesis and structural studies of Os(III),Mn(II),Fe(III), Co(II), Ni(II),and Cu(II)complexes of N-(4-methyl -8-acetoumbeliferonylidene)-N<sup>-</sup>(isonicotinoyl) hydrazine*», Ind.J. Chem.,27A,44-51.
  25. Raman,N., Esthar, S., and Thangaraja, C., 2004. «*A new mannich base and its transition metal (II) complexes - synthesis , structural ,characterization and electrochemical study*», J.Chem. Sci.,16(4),148-152.
  26. Sarga,W.J.,and Lenarcik,B., 1995. «*Coordination compounds of 2-mercapto-1-methylimidazol with Pt(II),Pd(II),Rh(III) and Ru(III)* », Trans.Met.Chem.,15(8),69-74.

27. Gupta, H.K., and Dikshit, S.K., 2000. «Reactions of Ru(II), Ru(III), Pt(II) and Pt(IV) with 2-mercapto-3-phenyl-4-quinazolinone in the presence and absence of various N-Heterocyclic base», J.Chem. Sci., 104(2), 117-122.
28. Imran, M., Iqbal, J., and Latif, S., 2006. «Synthesis, Characterization and invitro screening of quinoline derivative and its complexes with Ru(III), Rh(III), and Pt(IV)», J.Bio. Sci., 6(5), 98-108.
29. Saxena, R.C., Jain, C.L., and Benjamin, R., 1988. «Magnetic and spectral behaviour of Pd(II), Pt(IV) and Ru(III) complexes with some tridentate substituted thiosemicarbazones», J.Ind. Chem. Soc., LXIII, 69-78.
30. Agarwal, S.K., and Saxena, R.C., 1979. «Complexes of Cu(II), Ni(II), Co(II), Fe(III) and Os(III) with 4-p-dimethylamino-anilino-3-penten-2-one», J.Ind. Chem. Soc., LVI, 101-112.
31. Gajendr, M.R., and Agarwala, U.A., 2000. «Complexing behaviour of 1,3,4-thiadiazole-2-thiol-5-amino with Fe(II), Co(II), Ru(II), Ru(III), Pd(IV), Ir(III) and Pt(IV)», Inorg.Nucl.Chem., 37(3), 88-94.
32. Anjane, Y., and Rao, R., 1986. «Synthesis and antimicrobial studies of some metal complexes», Synth.React.Inorg.Met.Org.Chem., 16(3), 92-101.
33. Mishra, L., and Singh, V., 1998. «Synthesis and antimicrobial studies of some metal complexes», Ind.J. Chem., 32(8), 32-38.
34. Tweedy, B.G., 1985. «Phytopathology and Antibiotic ascientific approach» 1<sup>st</sup> Ed., Mir Publishers, pp208.
35. Dharmaraj, N., 2001. «Ruthenium (II) complexes containing bidentate schiff bases and their antifungal activity», Trans.Met. Chem., 26(10), 53-59.

### تحضير ودراسة كلاً من تراكيب و الفعالية البايولوجية لـ [4-(1-فنيـل-2و3-ثنائي المثل-3-بايروزولين-5-أون)أزو]-N,N-ثنائي مثيل- أنيلين مع بعض أيونات العناصر الانتقالية

رحاب عبد المهدي الحسني\*

\*الجامعة المستنصرية/ كلية العلوم / قسم الكيمياء

#### الخلاصة:-

تم في هذا البحث تحضير و تشخيص معقدات الليكاند الجديد :-

4-[(1-phenyl-2,3-dimethyl-3-pyrozoline-5-one)azo]-N,N-dimethyl anline (PAD) للعناصر الإنتقالية {الكوبلت (II) و النيكل (II) و البلاديوم (II) والكروم (III) و الحديد (III) و الروثينيوم (III) و الروديوم (III) و الأوزميوم (III) و الأريديوم (III) و البلاتينيوم (IV) و الفناديوم (IV)}. درس (PAD) وشخص بطرق تحليل طيف الأشعة تحت الحمراء (FT-IR) وطيف الأشعة فوق البنفسجية (UV) والتحليل الدقيق للعناصر (C.H.N)، في حين شُخصت المعقدات المحضرة وتم إقتراح الشكل الهندسي لها بالحالة الصلبة باستخدام تقنية الامتصاص الذري للهبتي للعناصر والتحليل الدقيق للعناصر (C.H.N) وطيف الأشعة تحت الحمراء (FT-IR) وطيف الأشعة فوق البنفسجية- المرئية (UV-Vis) فضلاً عن قياسات الحساسية المغناطيسية والتوصيلية الكهربائية. تم كذلك دراسة طبيعة المعقد المتكون في محلول الايثانول باتباع طريقة النسبة المولية ، وقد اعطت هذه الدراسة نتائجاً مطابقة مع تلك التي تم الحصول عليها في الحالة الصلبة . كما درست ثوابت الاستقرار للمعقدات المتكونة مع الزمن وكانت مدة ثبات اللون اكثر من (6) ساعات . فضلاً عن ذلك تم حساب قيمة الممتصية المولارية للمعقدات . تم تقويم الفعالية المضادة للبكتريا للليكاند (PAD) ومعقداته وأختير نوعان من من البكتريا و ( *Klesbiella pneumonia subtilus* ) ، سالية الصبغة) و (*Streptococcus faecalis*) ، موجبة الصبغة) لهذا الغرض، بالإضافة إلى ذلك تم إجراء التقويم الحيوي للليكاند ومعقداته ضد نوعين من الفطريات (*Candida albicans*) و (*Aspergillus flavus*) .