

Molecular characterization and microbial resistance of different bacterial isolates in some dairy products

W.K. Alkhafaje¹ , Z.A. Olama¹ and S.M. Ali²

¹Botany and Microbiology Department, Faculty of Science, Alexandria University, ²Nucleic Acid Research Department, Genetic Engineering and Biotechnology Research Institute, City of Scientific Research and Technological applications, Alexandria, Egypt

Article information

Article history:

Received June 10, 2021
Accepted August 14, 2021
Available online March 1, 2022

Keywords:

Multidrug resistance
Gene
Bacteria
Milk
Dairy products

Correspondence:

S.M. Ali
safaa.mohamedali@yahoo.com

Abstract

Bacterial contamination of milk and dairy products is a common problem. In the last two years, the foodborne diseases caused by the intake of milk and dairy products have been mostly disturbed with *Salmonella enterica*, *Listeria monocytogenes*, *Escherichia coli* O157:H7 and *Campylobacter jejune*. The study aims to isolate multidrug resistance (MDR) bacteria in dairy products and study of the molecular characterization of that isolates. MDR bacteria were found in 30 out of the 131 bacterial isolates. The incidence of MDR bacterial isolates revealed the abundance of *Staphylococcus* sp. with 43.3%, *Bacillus* sp 16.7%, *Salmonella* 13.3%, *E. coli* 10 %, *Enterococcus* 6.7 % *Pseudomonas* 3.3 %, *Shigella* 3.3 % and *Proteus* 3.3 %. Molecular studies of genes presence or absence for class A contain TEM, CTX and BSHV, class B contain VIM, IMP, KPC and NDM, class C contain FOX and class D contain OXA-10, OXA-24 and OXA-58 were tested. NDM, TEM, CITM and OXA -10 genes were the most abundant the selected bacterial isolates. The results of this study indicate that cheese made from unpasteurized milk can pose a significant risk to consumers. Product manufacturing processes should be subject to health control-to-control pathogens. The novelty in this work depend on screening of gene responsible of the resistance from the bacteria isolated from dairy product using the molecular technique.

DOI: [10.33899/ijvs.2021.130206.1764](https://doi.org/10.33899/ijvs.2021.130206.1764), ©Authors, 2022, College of Veterinary Medicine, University of Mosul.
This is an open access article under the CC BY 4.0 license (<http://creativecommons.org/licenses/by/4.0/>).

Introduction

The spoiling milk products worldwide is a huge economic problem. The microbial capacity and incidence of the bacterial pathogens in foods are indicators of the quality and safety of food. In addition, the education of the food handlers about the personal hygiene is importance from the food safety point of view (1). The highly nutritious nature of the dairy products makes them especially good media for the growth of microorganisms (2). The microbial contamination is one of the leading causes of the food spoilage worldwide (3). The contamination of food with microbes can occur at any stage of the food chain (4). The large number of diseases caused by food borne pathogens with significant effects on economy and human health. The bacterial pathogens use a

variety of different motility modes, including swimming, twitching, and swarming (1). *Staphylococcus aureus* is commonly food born pathogen of the great importance of animal and human concern. It is responsible for contaminate dairy products, kariesh cheese and ice cream from the different sources during their production, processing and handling that make them unfit for human consumption or even a dangerous source of the infection among customers establishing a potential health hazard (5). Also, *S. aureus* known for its ability to secrete a lot of toxins to aid in host tissue infiltration and a acquire nutrients (6). *Salmonella*, *Shigella*, *Listeria monocytogenes*, *Bacillus cereus*, *Campylobacter*, *Clostridium*, *Cronobacter*, *Escherichia* (*E.*) *coli*, *Staphylococcus* (*S.*) *aureus*, *Vibrio*, *Yersinia* (*Y.*) *enterocolitica*, viruses such as hepatitis A and Noro viruses,

and parasites *Cyclospora (C.) cayetanensis*, *Toxoplasma (T.) gondii*, and *Trichin* (7). The antimicrobial resistance (AMR): has the ability of kill microbes or limit their growth to grow in the presence of a drugs (8). The AMR complicates infection treatment and linked to increased mortality and morbidity. The emergence and spread of resistant and multidrug-resistant (MDR) bacteria has the enormous implications for worldwide healthcare delivery and the population health (9). The virulence functions encoded on the large extrachromosomal plasmids by pathogenic bacteria. These plasmids maintained at the low copy number to reduce the metabolic burden on their host (10). The widespread use of extended-spectrum cephalosporin creates a reservoir of resistant bacteria. Moreover, multi-resistance frequently associated with strains carrying (Extended-spectrum beta-lactamase) ESBLs, which could dramatically reduce the treatment options. The increasing number of *Enterobacteriaceae* with ESBLs that also contain (Metallo- β -lactamases) MBLs or AmpCs and other new mechanisms of resistance to fluoroquinolones or aminoglycosides indicate that recent increase of ESBLs -producing bacteria in Europe constitutes a complex problem (11).

ESBLs are worthy of the scientific community's attention over the past decades among the β -lactamases. ESBL's older and classical definition includes TEM-1, TEM-2, or SHV derivatives. ESBL divided into three main groups by the most recent definition. (I) ESBLA (class A ESBLs): CTX-M, SHV and TEM enzymes. (II) ESBLM (miscellaneous ESBLs) are sectioned into ESBLM-C (class C, plasmid mediated AmpC) and ESBLM-D (class D). (III) ESBLCARBA (ESBLs that degrade carbapenems) are divided into ESBLCARBA-A, ESBLCARBA-B, and ESBLCARBA-D. More than 500 β -lactamases reported to date produced by diverse bacteria. Beta-lactamases thought to be the most common resistance mechanism that contributes to widespread resistance among Gram-negative microbes also transmission of resistance occur between microorganisms (12). NDM-1 producing by *E. coli* infects the host by commonly invading sites like, urinary tract, blood, lungs, and wounds, leading to the urinary tract infections, septicemia, pulmonary infections, diarrhea, peritonitis, device-associated infections and soft tissue infections (13). The study's goal is to identify MDR bacteria from dairy products and examine their molecular characteristics.

Materials and methods

Sample collection and analysis of microbiology

Dairy products: samples from different locations in Alexandria collected during 2018: white cheese, white cheese produced, old cheese and milk. The sample collected for further use in sterile containers. To evaluate the incidence of isolates of bacterial ESBL and CR isolated from the certain dairy products at the various locations in Alexandria, Egypt. 100 sample were collected that were distributed as

Kareish cheese, Industrial white cheese, Old cheese, Roomy cheese and milk 40, 28, 15, 10 and 7% respectively.

Assessment of isolated bacteria's resistance prevalence

According to the modified Kirby-Bauer Disc Diffusion method, all the isolated bacteria subjected to the antibiotics resistance by using the disk diffusion method.

Microorganisms and molecular identification

The antibiotic resistant isolates subjected to phenotypic identification using the cultural characteristics in a trail to identify. Gram staining and analysing biochemistry. The region of 16S rRNA was amplified by using the universal primers F: AGAGTTTGATCMTGGCTCAG and R: TACGGYACCTTGTTACGACTT. The PCR reaction, was performed for 4 min at 95°C followed by 40 cycles each of (Denaturation: 40 second at 94°C, annealing: 50 second at 58°C and extension: 50 second at 72°C), followed by a supplementary 10min at 72°C. Sequences of the 16S rRNA genes obtained from the NCBI database. Multiple alignments based on the most closely related sequences and the similarity levels carried out by using the BLAST program1. A phylogenetic tree reconstructed using the Mega 5 software.

Bacterial resistance determination using molecular techniques

The fresh bacterial cells used to extract DNA by using the GeneJET Genomic DNA Purification Kit. GEBRI kit removed plasmid from the bacteria's selected isolates.

Virulence gene detection using PCR assay

Eighteen primers used in this study; the genus used for screening of ESBL class A were TEM, CTX BETA and BSHV. ESBL class B carbapenemase encoding genus were CITM, VIM, IMP, KPC and NDM. ESBL class C encoding genus ACCM. ESBL class D encoding genus was OXA-10, OXA-55, OXA58, OXA 60 and OXA 69. Screening for coding sequence of genes DHAM, EBCM, MEC A, VAN A, and TOHO1 evaluated. PCR reaction achieved for 4 min at 95°C followed by 35 cycles each of (Denaturation: 40 second at 94°C, annealing: 50 second at 50-60°C and extension: 50 second at 72°C), followed by a supplementary 10min at 72°C. After amplification by PCR assay, the products checked in 2% agarose gel electrophoresis. The genetic miscellany was determined as the experimental number of differentiation (Table 1).

Results

Assessment of isolated bacteria's resistance prevalence

The multiantibiotic resistance shown by the biggest promising isolates. Seven bacteria isolates were isolated from milk but show sensitive effect towards the antibiotic used for detection of MDR bacteria. All the selected MDR isolates were resistance to metronidazole (MTZ). Incidence

of antibiotic sensitivity of multi-antibiotic resistance bacterial isolates (Table 2). Distribution of the MDR bacterial isolates were 14 isolates from Kariesh cheese (out of 52 bacterial isolates), 11 isolates from industrial white

cheese (out of 19 bacterial isolates), 3 isolates from old cheese (out of 37 bacteria isolates) and 2 isolates from roomy cheese (out of 13 bacteria isolates) (Figure 1).

Table 1: primers used in the work-study

Gene	Primer	Sequence (5- 3)	Size [bp]	Pcr program	Reference
NDM	NDM-1	GGTTTGGCGATCTGGTTTTTC	580	55	14
	NDM-2	CGGAATGGCTCATCACGATC			
KPC	KPC 1	CGTCTAGTTCTGCTGTCTTG	750	56	15
	KPC 2	CTTGTCATCCTTGTTAGGCG			
VIM	VIM-1	GATGGTGTGTTGGTCGCATA	780	54	16
	VIM-2	CGAATGCGCAGCACCAG			
TEM	TEM-1	AAAATTCTTGAAGACG	1000	60	17
	TEM-2	TTACCAATGCTTAATCA			
DHAM	DHAM-1	AACTTTCACAGGTGTGCTGGG	500	60	18
	DHAM-2	TCCGTACGCATACTGGCTTTGC			
MECA	MECA-1	AAAATCGATGGTAAAGGTTGGC	550	59	19
	MECA-2	AGTTCTGCAGTACCGGATTTTG			
CTX_beta	CTX beta-1	TTTGCGATGTGCAG(C/T)ACCAG	650	58	Designed
	CTX beta-2	CGCGATATC(A/G)TTGGTGGTGCCATA			
OXA 58	OXA 58-1	AAGTATTGGGGCTTGTGCTG	600	58	20
	OXA 58-2	CCCCTCTGCGCTCTACATAC			
CITM	CITM-1	TTT CTC CTG AAC GTG GCT GGC	500	55	21
	CITM-2	TGG CCA GAA CTG ACA GGC AAA			
CTX	CTX-1	TTAGGAAATGTGCCGTGTA	900	56	22
	CTX-2	CGATATCGTTGGTGGTACCAT			
ACCM	ACCM-1	AACAGCCTCAGCAGCCGGTTA	750	53	23
	ACCM-2	TTCGCCGCAATCATCCCTAGC			
OXA-10	OXA-10-1	GCCATGAAAACATTTGCCGC	350	53	24
	OXA-10-2	GCCACCAATGATGCCCTCAC			
IMP	IMP-1	GTGGTTCTTGTAATGCTGAGG	650	55	25
	IMP-2	CCGCCTGCTCTAATGTAAGT			
EBCM	EBCM-1	TCGGTAAAGCCGATGTTGCGG	500	56	26
	EBCM-2	CTTCCACTGCGG CTG CCA GTT			
OXA-69	OXA-69-1	CTAATAATTGATCTACTCAAG	500	60	27
	OXA-69-2	CCAGTGGATGGATGGATAGATTATC			
TOHO-1	TOHO1-1	GCGACCTGGTTAACTACAATCC	600	60	28
	TOHO1-2	CGGTAGTATTGCCCTTAAGCC			
BSHV	BSHV-1	ATGCGTTATATTCGCCTGT	500	56	29
	BSHV-2	TGCTTTGTTATTCGGGCCAA			
VAN A	VANA-1	GGGAAAACGACAA TTGC	750	50	30
	VANA-2	GTACAATGCGGCCGTTA			

Identification of the bacterial isolates

The most commonly isolated pathogens were evaluated for phenotypic characterization and molecular identification using 16s rDNA. The amplified fragment sequences of 1500 bp identified. A dominant tool for identifying and classifying prokaryotes was the sequences of the different types of the strains from the gene bank. GenBank deposited the sequence of the most promising isolates and had the accession. From all the examined samples were *staphylococcus spp*, *Bacillus spp*, *Salmonella spp*, *E. coli*, *Enterococcus spp*,

Pseudomonas spp, *Shigella spp* and *Proteous spp*. with 43.3, 16.7, 13.3, 10, 6.7, 3.3, 3.3 and 3.3 % respectively. 23.1% out of the total isolates 130 were pathogens.

Resistance determination using molecular technique

The variation between the isolated MDR bacteria shown by plasmid extraction of the selected bacterial isolates. The results showed that isolates 2, 9, 16, 19, 20, 24, 25, 27, 29, 30, 39, 40 and 41 were highly antibiotic-resistant positive plasmid with high copy numbers (Figure 2).

Table 2: Resistance prevalence of the bacterial isolates

Antibiotics	Resistance %	Antibiotics	Resistance %
Colistin (CT)	66.7	Ceftazidime (CAZ)	90
Ampicilin +Sulbactam (A/S)	90	Amikacin (A/K)	70
Penicilin (P)	83.3	Clindamycin (DA)	93.3
Chloramphenicol (C)	76.7	Rifampine (RA)	86.7
Ofloxacin (OFX)	83.3	Nitrofurantion (F)	76.7
Levofloxacin (LEV)	86.7	Cefoperazone (CFP)	86.7
Cefepime (CPM)	93.3	Azetreonium (ATM)	96.7
Meropeneme (MEM)	70.7	Tetracycline (TE)	83.3
Metronidazole (MTZ)	100	Streptomycin (S)	83.3
Doxycycline (DO)	83,3	Trimethoprim-sulfamethoxaz (STX)	80




Figure 1: MDR incidence distribution in dairy product samples under test.




Figure 2: Agarose gel electrophoresis plasmid profiles of different bacterial isolates.

Detection of the virulence gene using PCR assay (Detection of β -lactamase gene class A, B, C and D)

Collected data of genes presence or absence for class A (TEM, CTX and BSHV) (Figure 3), class B (CITM, VIM, IMP, KPC and NDM) (Figure 4), class C (ACCM) (Figure 5), and class D (OXA-10, OXA-55, OXA-58 and OXA 69) (Figure 6). Screening of DHAM, EBCM, MEC, VAN and TOHA gene tested by using the selected bacterial isolates (Figure 7). The fragment size of amplified PCR product was calculated by using the software of Gel Documentation Analysis System (Alpha Imager TM 1220). In addition, the results of the present study showed that the average percentage of positive genes in all the selected MDR (30

isolates). Distribution of presence or absence of the 18 tested virulence genes among the MDR bacterial isolates (Figure 8).




Figure 3: Agarose gel electrophoresis of the amplified PCR fragment for Class A, no. refers to the code of positive isolate.




Figure 4: Agarose gel electrophoresis of the amplified PCR fragment for Class B, no. refers to the code of positive isolate.




Figure 5: Agarose gel electrophoresis of the amplified PCR fragment for Class C (ACCM gene), no. refers to the code of positive isolate.




Figure 6: Agarose gel electrophoresis of the amplified PCR fragment for Class D, no. refers to the code of positive isolate.




Figure 7: Agarose gel electrophoresis of the amplified PCR fragment for (EBCM, MEC, DHAM, VAN AND TOHA gene), no. refers to the code of positive isolate.




Figure 8: Average percentage of positive genes in all the selected MDR (30 isolates).

Discussion

Studies previously conducted in different countries revealed a wide *S. aureus* diversity in milk and dairy products. The prevalence of *S. aureus* was 43.3% and it was differing than which carried out from Iran and Italy on different dairy products revealed a lower *S. aureus* percentage. Lower prevalence noted in studies which detected *S. aureus* with 10, 11.3, 5 and 9.1% respectively for the kariesh cheese (31). Greater incidence of *S. aureus* was 72, 50, 70, 93 and 68 % respectively for the kariesh cheese (32). Percentage of *Bacillus spp.* in our study was 16.7% which were isolated from the industrial white cheese and the kariesh cheese where it was higher percentage in industrial white cheese. It was differed than the study by EFSA 2005 which appeared the incidence of *B. cereus* was highest in kariesh cheese 25 to 80% during all the seasons followed by koshary 45 to 70% and cornsnacks 20 to 50%. Percentage of *Salmonella sp.* in our study was 13.3% that were isolated from the industrial white cheese and the kariesh cheese 10 and 3.3%. Relatively lower results were obtained by Abo Zeed (33,34) where *Salmonella spp.* could not be identified in the surveyed of the white cheese samples. Ghada and colleagues (35) identified two samples polluted with *Salmonella spp.*

In our study, the *E. coli* percentage was 10% isolated from roomy, kariesh and industrial white cheese compared to other bacterial isolates. In study by Abdelrahman *et al.*, (36) found that, the incidence of *E. coli* in Kariesh, and Domiati cheese were 37.1% and 2.8%, respectively. While, could not be detect in Tallaga cheese. Naj and Ghanbarpour (37) who isolated *E. coli* with percentage of 98.7 % from the domestic soft cheese. A common principal of the resistance markers in all varieties estimated and associated antimicrobials such as tetracycline, B-lactams, sulfonamide and quinolones were detected (38). An aggregate number of extended-spectrum β -lactamases (ESBLs) have been predictable in Enterobacteriaceae through the latest few years. SHV types of enzymes shown to carry the SHV-1 gene within the chromosome (39). NDM-1 producers were resistant to imipenem, meropenem, ertapenem, gentamicin, amikacin, tobramycin, and ciprofloxacin, whereas, isolates were susceptible to colistin (40). The high prevalence of tetracycline- and penicillin-resistant *S. aureus*, *Salmonella* and *E. coli* observed in the current study, is in agreement with earlier findings. Gram-negative bacteria mostly preset TEM, almost 90% of the resistance against ampicillin in Gram-negative bacteria are due to TEM encoded genes (41).

Conclusion

The results of this study indicate that cheese made from unpasteurized milk can pose a significant risk to consumers. This risk varies depending on the geographical location of the study area, the state of education of the population, the extent of attention to hygiene and the method of preparing

and packaging dairy products. Product manufacturing processes should be subject to health control-to-control pathogens. Dairy markets monitored for pathogens. Reducing the surface area exposed to air reduces harmful microbial growth in dairy products.

Conflict of interest

No conflict.

Acknowledgments

The authors acknowledge City of Scientific Research and Technological Applications.

References

- Mahendra P, Selamawit M, Muluken T, Suneeta VP, Prajapati JP. Bacterial contamination of dairy products. *Beverage Food World*. 2016;43:40-43. [\[available at\]](#)
- Pal M, Bekele T, Feleke A. Public health significance of pasteurized milk. *Beverage Food World*. 2012;39: 55-56. [\[available at\]](#)
- Pal M. Spoilage of dairy products due to fungi. *Beverage Food World*. 2014;41: 37-38. [\[available at\]](#)
- Vrdoljak J, Dobrani V, Filipovi I, Zdolec N. Microbiological quality of soft, semi-hard and hard cheeses during the shelf-life. *J Macedonian Vet Rev*. 2016;39:59-64. DOI: [10.1515/macvetrev-2015-0068](#)
- Khalil II, AlDabbagh SYA, Shareef AM. Isolation, identification and detection of some virulence factors in yeasts from local cheese in Mosul city. *Iraqi J Vet Sci*. 2018;32(1):81-85. [\[available at\]](#)
- Kong C, Neoh H, Nathan S. Targeting *Staphylococcus aureus* toxins: A potential form of anti-virulence therapy. *Toxins*. 2016;8(3):72. DOI: [10.3390/toxins8030072](#)
- Bintsis T. Foodborne pathogens. *AIMS Microbiol*. 2017;3(3):529. DOI: [10.3934/microbiol.2017.3.529](#)
- World Health Organization. Antimicrobial resistance: Global report on surveillance. Switzerland: World Health Organization; 2014. [\[available at\]](#)
- Liu YY, Wang Y, Walsh TR, Yi LX, Zhang R, Spencer J, Doi Y, Tian G, Dong B, Huang X. Emergence of plasmid-mediated colistin resistance mechanism MCR-1 in animals and human beings in China: A microbiological and molecular biological study. *Lancet Infectious Dis*. 2016;16(2):161-168. DOI: [10.1016/S1473-3099\(15\)00424-7](#)
- Mustafa JY. Study the effect of cloned pET-32a (+) plasmid by Lysostaphin gene against *Staphylococcus aureus*. *Iraqi J Vet Sci*. 2020;35(2):233-238. DOI: [10.33899/ijvs.2020.126698.1362](#)
- Urumova V. Extended spectrum beta lactamase producing animal enterobacteriaceae isolates as potential risk to public health - review. *Valentina*. 2015;166:192-207. [\[available at\]](#)
- Walsh TR, Weeks J, Livermore DM, Toleman MA. Dissemination of NDM-1 positive bacteria in the New Delhi environment and its implications for human health: An environmental point prevalence study. *Lancet Infect Dis*. 2011;11:355-62. DOI: [10.1016/S1473-3099\(11\)70059-7](#)
- Hu H, Hu Y, Pan Y, Liang H, Wang H, Wang X, Hao Q, Yang X, Yang X, Xiao X. Novel plasmid and its variant harboring both a bla (NDM-1) gene and type IV secretion system in clinical isolates of *Acinetobacter lwoffii*. *Antimicrob Agents Chemother*. 2012;56:1698-1702. DOI: [10.1128/AAC.06199-11](#)
- Nor Zanariah Z, Anita S, Hanafiah A, Najihan AS. Molecular detection of the New Delhi metallo- β -lactamase-1 gene in Enterobacteriaceae isolates in a tertiary medical centre. *Malaysian J Pathol*. 2015;37(3):227-232. [\[available at\]](#)
- Patrice, N, Thierry, N, Laurent, P. Global Spread of Carbapenemase-producing Enterobacteriaceae. *Emerg Infect Dis*. 2011. 17(10):1791-1798. DOI: [10.3201/eid1710.110655](#)
- Jyoti S, Meera S, Pallab R. Detection of TEM and SHV genes in *Escherichia coli* and *Klebsiella pneumoniae* isolates in a tertiary care hospital from India. *Indian J Med Res*. 2010. 132: 332-336. [\[available at\]](#)
- Hosseini S, Maleki A, Ghafourian S, Reza M, Fadavi HV, Shahmir P, Sadeghifard N. Molecular Characterization of AmpC β -Lactamases among *Klebsiella pneumoniae* isolated from Ilam and Tehran hospitals, from Iran. *J Pure Appl Microbiol*. 2014;8(3):2217-2220. [\[available at\]](#)
- Taweeporn S, Chariya C, Kunyaluk C, Temduang L, Chaisiri W. Evaluation of different primers for detecting Meca gene by PCR in comparison with phenotypic Methods for discrimination of Methicillin-resistant *Staphylococcus aureus*. *Southeast Asian J Trop Med Public Health*. 2002;15(1):34-40. [\[available at\]](#)
- Khorsi K, Messai Y, Hamidi M, Ammari H, Bakour R. High prevalence of multidrug-resistance in *Acinetobacter baumannii* and dissemination of carbapenemase-encoding genes blaOXA-23-like, blaOXA-24-like and blaNDM-1 in Algiers hospitals. *Asian Pacific J Trop Med*. 2015;8(6):438-446. DOI: [10.1016/j.apjtm.2015.05.01](#)
- Sana J, Mohd S, Sobia F, Anuradha S, Haris MK. Molecular characterization of genes encoding AmpC beta-lactamases in clinical isolates of *Pseudomonas* and *Acinetobacter* species. *J Appl Pharmaceut Sci*. 2015;5(10):048-051. DOI: [10.7324/JAPS.2015.501009](#)
- Mohammad SN, Sreedevi B, Chaitanya RK, Sreenivasulu D. Betalactamase antimicrobial resistance in Klebsiella and Enterobacter species isolated from healthy and diarrheic dogs in Andhra Pradesh, India. *Vet World*. 2017;2231-0916. [\[available at\]](#)
- Kolar M, Bardon J, Chroma M, Hricova K, Stosova T, Sauer P, Koukalova D. ESBL and AmpC beta-lactamase-producing Enterobacteriaceae in poultry in the Czech Republic. *Vet Med*. 2010;55(3):119-124. DOI: [10.17221/165/2009-VETMED](#)
- Frédéric B, Catherine B, Nicole LZ. Identification of PSE and OXA β -lactamase genes in *Pseudomonas aeruginosa* using PCR-restriction fragment length polymorphism. *Brit Soci Antimicrobiol Chemo*. 2002;50:11-18. [\[available at\]](#)
- José AD, Alejandra B, Juan A, Cynthia R, Ángela F, Gabriel OG. Blactamases produced by amoxicillin-clavulanate-resistant Enterobacteria isolated in Buenos Aires, Argentina: A new blaTEM gene. *Rev Argent Microbiol*. 2014; 46(3):210-217. DOI: [10.1016/S0325-7541\(14\)70075-6](#)
- Patrik M, Magdalena R, Milan K. Research article primer evaluation for PCR and its application for detection of carbapenemases in enterobacteriaceae Jundishapur. *J Microbiol*. 2016; 9(1):e29314. DOI: [10.5812/jjm.29314](#)
- Chih-Chuan K, Meei-Fang L, Chin-Fu L, Yi-Ching H, Po-Yu L, Ching-Wen C, Zhi-Yuan S. Antimicrobial susceptibility and multiplex PCR screening of AmpC genes from isolates of *Enterobacter cloacae*, *Citrobacter freundii*, and *Serratia marcescens*. *J Microbiol Immunol Infect*. 2010;43(3):180-187. DOI: [10.1016/S1684-1182\(10\)60029-1](#)
- Evans BA, Hamouda A, Towner KJ, Amyes SB. OXA-51-like β -lactamases and their association with particular epidemic lineages of *Acinetobacter baumannii*. *Clin Microbiol Infect*. 2008;14(3):268-275. DOI: [10.1111/j.1469-0691.2007.01919.x](#)
- Tadesse E, Josephine B, Daniel A, Moses NN, Joyce N, Wondwossen AG, John SG, Appolinaire D, Ephrem E. Genetic markers associated with resistance to beta-lactam and quinolone antimicrobials in non-typhoidal Salmonella isolates from humans and animals in central Ethiopia. *Antimicrobiol Resist Infect Cont*. 2017;10(8):189-196. DOI: [10.1186/s13756-017-0171-6](#)
- Nashwa MA, Maysaa EZ. Molecular study of *Acinetobacter baumannii* isolates for metallo- β -lactamases and extended-spectrum- β -lactamases genes in intensive care unit, Mansoura university hospital, Egypt. *Inter J Microbiol*. 2017;3925868. DOI: [10.1155/2017/3925868](#)
- Chimanjita P, Mangala L, Swapnil R, Kandarpa KS. Emergence of va 17 gene among vancomycin-resistant enterococci in a tertiary care hospital of North - East India. *Indian J Med Res*. 2016;143(3):357-361. DOI: [10.4103/0971-5916.182627](#)
- Mahmood MA, Essa MA. Antimicrobial activity of peptides extracted from camels' blood neutrophils against some pathogenic bacteria. *Iraqi J Vet Sci*. 2021;35(1):33-37. DOI: [10.33899/ijvs.2020.126239.1270](#)

32. Salama EM, Saad AH, Enan GA, Suzan IY. Incidence and biocontrol of *Staphylococcus aureus* in some milk products. Second Conference of Food Safety, Suez Canal University, Faculty of Veterinary Medicine; 2015. 29-35 p.
33. El-Gamal MS, El Dairouty RK, Okda AY, Sahar HS, El-Shamy SM. Incidence and interrelation of *Cronobacter sakazakii* and other foodborne bacteria in some milk products and infant formula milks in Cairo and Giza Area. World Applied Sci. J. 2013;26:1129-1141. DOI: [10.5829/idosi.wasj.2013.26.09.13542](https://doi.org/10.5829/idosi.wasj.2013.26.09.13542)
34. Abo Zeed BA. prevalence of biological hazards in milk and some dairy products in Egyptian markets [PhD dissertation]. Cairo: Faculty of Veterinary Medicine, Cairo University; 2014.
35. Ghada ZAA, Alia MH, Soha Al-S, Magdy NA, Mohammed FS. Chemical, nutritional and microbiological evaluation of some Egyptian soft cheeses. Egyptian J Hospital Med. 2004;17:44-57. [\[available at\]](#)
36. Abdelrahman M, El Bagoury HS, Hadeer S. Incidence of *Escherichia coli* and *Salmonella* species with special reference to antibiotic resistant pathogenic *E. coli* isolated from locally produced cheeses in Egypt. Alexandria J Vet Sci. 2019;10:5455. DOI: [10.5455/ajvs.21944](https://doi.org/10.5455/ajvs.21944)
37. Najand LM, Ghanbarpour R. A study on enteropathogenic *Escherichia coli* isolated from domestic Iranian soft cheese. Vet Arh. 2006;76:531-536. [\[available at\]](#)
38. Fontes CO, Silva VL, de Paiva MRB, Garcia RA, Resende JA, Ferreira-Machado AB, Diniz CG. E-negative *Staphylococci* isolated from soft cheese in Brazil. J Food Sci. 2013;78:594-599. DOI: [10.1111/1750-3841.12088](https://doi.org/10.1111/1750-3841.12088)
39. Ahmed IM. Detection of CTX-M gene in extended spectrum β -lactamases producing Enterobacteriaceae isolated from bovine milk. Iraqi J Vet Sci. 2021;35(2):397-402. DOI: [10.33899/ijvs.2020.126909.1412](https://doi.org/10.33899/ijvs.2020.126909.1412)
40. Kumarasamy KK, Toleman MA, Walsh TR, Bagaria J, Butt F, Balakrishnan R. Emergence of a new antibiotic resistance mechanism in India, Pakistan, and the UK: A molecular, biological, and epidemiological study. Lancet Infect Dis. 2010;10:597-602. DOI: [10.1016/S1473-3099\(10\)70143-2](https://doi.org/10.1016/S1473-3099(10)70143-2)
41. Sadeeq R, Tariq A, Ijaz A, Nazir AK, Bo H, Jian Gao. The growing genetic and functional diversity of extended spectrum beta-lactamases. Res Inter. 2018. DOI: [10.1155/2018/9519718](https://doi.org/10.1155/2018/9519718)

التوصيف الجزيئي والمقاومة الميكروبية لل عزلات البكتيرية المختلفة في بعض منتجات الألبان

وليد خالد الخفاجي^١، زكية أحمد علما^١ و صفاء محمد علي^٢

^١ قسم النبات والميكروبيولوجي، كلية العلوم، جامعة الإسكندرية، أقسم بحوث الأحماض النووية، معهد الهندسة الوراثية، مدينة الأبحاث العلمية والتطبيقات التكنولوجية، الإسكندرية، مصر

الخلاصة

يعد التلوث الجرثومي للحليب ومنتجاته إحدى أهم المشاكل شائعة الحدوث. في العامين الماضيين، تحدث الأمراض التي تنتقل عن طريق الغذاء نتيجة تناول الحليب ومنتجاته إلى اضطراب توزيع لبكتيريا السالمونيلا انتيريكا واليسيريا وايشيريشيا كولاي وكمبيلاباكتريجيونا. فكان الهدف الرئيسي من الدراسة هو عزل بكتيريا التي لها قدرة على مقاومة العديد من المضادات الحيوية التي عزلت من الحليب ومنتجات الألبان ودراسة التوصيف الجزيئي لتلك العزلات. إن ٣٠ عزلة بكتيرية من أصل ١٣١ عزلة كانت مقاومة للعديد من المضادات الحيوية. كشفت الدراسة إن البكتيريا من نوع الاستافيلوكوكس هي أكثر الأنواع الموجودة بين تلك العزلات بنسبة ٤٣,٣٪، ١٦,٧٪ من نوع الباسيلس، السالمونيلا كانت ١٣,٣٪، الإشيريشيا كولاي كانت ١٠٪، الانتيريوكوكاس ٦,٧٪ و ٣,٣٪ للسيدوموناس، الشغيلة كانت ٣,٣٪ والبروتيويس كان ٣,٣٪. لقد تمت الدراسات الجزيئية لوجود الجينات أو غيابها للفئة أ (TEM) و CTX و BSHV) والفئة ب (VIM و IMP و KPC و NDM) والفئة ج (FOX) والفئة د (OXA-10 و OXA-24 و OXA-58). كانت جينات NDM و TEM و CITM و (OXA-10) الأكثر وفرة في العزلات البكتيرية المختارة. تشير نتائج هذه الدراسة إلى أن الجبن المصنوع من الحليب غير المبستر يمكن أن يشكل خطراً كبيراً على المستهلكين. يجب أن تخضع عمليات تصنيع منتجات الألبان إلى الرقابة الصحية لمنع مسببات الأمراض. تعتمد الحداثة في هذا العمل على فحص الجين المسؤول عن المقاومة من البكتيريا المعزولة من منتجات الألبان باستخدام تقنية البيولوجيا الجزيئية.

