

## L-pre- and L-semi-P- compact Spaces

*Nerjis Abdul Jabbar Dawood\**

*Suaad Gedaan Gasim\**

Received 4, August, 2010

Accepted 14, January, 2011

### Abstract:

The purpose of this paper is to study a new types of compactness in the dual bitopological spaces. We shall introduce the concepts of L-pre- compactness and L-semi-P- compactness .

**Key words:** L-pre-compact ,L-semi-p-compact ,L-pre-open,L-semi-p-open.

### Introduction:

The concepts of bitopological space was initiated by Kelly[1].A set  $X$  equipped with two topologies  $\tau_1$  and  $\tau_2$  is called a bitopological space denoted by  $(X, \tau_1, \tau_2)$ . Navalagi [2] introduced the concepts of pre-open and semi-P-open sets. A subset  $A$  of a topological space  $(X, \tau)$  is said to be “pre-open” set if and only if  $A \subseteq \text{int } cl(A)$ , the family of all pre open subsets of  $X$  is denoted by  $PO(X)$ .The complement of a pre-open set is called pre-closed set, the family of all pre- closed subsets of  $X$  is denoted by  $PC(X)$  [2].The smallest pre- closed subset of  $X$  containing  $A$  is called “pre-closure of  $A$ ” and is denoted by  $pre-cl(A)$ [3].

Let  $(X, \tau)$  be a topological space, a subset  $A$  of  $X$  is said to be “semi-P-open” set if and only if there exists a pre-open subset  $U$  of  $X$  such that  $U \subseteq A \subseteq pre-cl(U)$ , the family of all semi –p-open subsets of  $X$  is denoted by  $SPO(X)$ .The complement of a semi-p-open set is called “semi-p-closed” set, the family of all semi-p-closed subsets of  $X$  is denoted by  $SPC(X)$ . The smallest semi-p-closed

set containing  $A$  is called semi-p-closure of  $A$  denoted by  $semi-p-cl(A)$ [4]. [3] shows that every open set is a pre-open and the union of any family of pre-open subsets of  $X$  is a pre-open set, but the intersection of any two pre-open subsets of  $X$  need not be apre-open set.[4] shows that every pre-open set is a semi –p-open and consequentiy every open set is a semi-p-open. Also she shows that the union of any family of semi-p-open subsets of  $X$  is a semi-p-open set, but the intersection of any two semi-p-open subsets of  $X$  need not be a semi-p-open set.

L-open set was studied by Al-Talkhany [5], a subset  $G$  of a bitopological space  $(X, \tau_1, \tau_2)$  is said to be “L –open” set if and only if there exists a  $\tau_1$ -open set  $U$  such that  $U \subseteq G \subseteq cl_{\tau_2}(U)$ , the family of all L-open subset of  $X$  is denoted by  $L-O(X)$ .The complement of an L-open set is called “L-closed” set, the family of all L-closed subsets of  $X$  is denoted by  $L-C(X)$ .In a bitopological space  $(X, \tau_1, \tau_2)$  every  $\tau_1$ -open set is an L-open set[5].The union of any family of L-open subsets of  $X$  is an L-open set, but the intersection of any two L-open

---

\*Department of Mathematics-Ibn-Al-Haitham College of Education - University of Baghdad

subsets of X need not be L-open set[5].

A collection of sets is said to have the finite intersection property (FIP) if and only if the intersection of each finite subcollection of it is non empty.[6]

**2-L-pre - and L-semi-p - compact spaces**

In this section we shall introduce a new typ of compactness namely L-pr – (L-semi-p-) compactness. We start with definition of L-pre-(L-semi-p-) open set.

**Definition (2.1):**

Let  $(X, \tau_1, \tau_2)$  be a bitopological space and let G be a subset of X. then G is said to be:

- 1- “L-pre-open” set if and only if there exists a  $\tau_1$ -pre-open set U such that  $U \subseteq G \subseteq cl\tau_2(U)$ .The family of all L-pre-open sub sets of X is denoted by  $L - PO(X)$ .
- 2- “L-semi-P-open” set if and only if there exists a  $\tau_1$ - semi-P-open setU such that  $U \subseteq G \subseteq cl\tau_2(U)$ .The family of all L- semi-P-open sub sets of X is denoted by  $L - SPO(X)$ .

**Definition (2.2):**

Let  $(X, \tau_1, \tau_2)$  be a bitopological space and let A be a subset of X .

- 1. By an “L-open cover of A” we mean a subcollection of the family L-O(X) which covers A .
- 2. By an “L -pre-open cover of A” we mean a subcollection of the family L-PO(X) which covers A.
- 3. By an “L -semi-p-open cover of A” we mean a subcollection of the family L-SPO(X) which covers A.

**Remark (2.3):**

- 1- Every L-open cover is an L- pre-open.

- 2- Every L-pre-open cover is an L-semi-P-open.

- 3- Every L-open cover is an L-semi-P-open.

The converse of each case of remark (2.3) is not true in general as the following example shows:

**Example (2.4):**

Let  $X = \{a, b, c, d\}$

$$\tau_1 = \{X, \emptyset, \{a\}, \{b, c\}, \{a, b, c\}\}$$

$\tau_2 = D$ =The discrete topology=The power set of X

$$L - O(X) = \{X, \emptyset, \{a\}, \{b, c\}, \{a, b, c\}\}$$

$$L - PO(X) = \left\{ X, \emptyset, \{a\}, \{b\}, \{c\}, \{a, b\}, \{a, c\}, \{b, c\}, \{a, b, c\}, \{a, b, d\}, \{a, c, d\} \right\}$$

$$L - SPO(X) = L - PO(X) \cup$$

$$\{\{a, d\}, \{c, d\}, \{b, c, d\}\}$$

Let  $C = \{\{c\}, \{a, b, d\}\}$  and

$B = \{\{a, d\}, \{b, c\}\}$ , clear that B is an L-semi-p- open cover, but it is neither L-pre-open nor L-open, and C is an L-pre-open cover, but it is not L-open cover.

**Remark (2.5):**

Every  $\tau_1$ -pre-open( $\tau_1$ -semi-p-open) cover of a sub set of a bitopological space  $(X, \tau_1, \tau_2)$  is an L-pre-open “L-semi-p-open” respectively.

The opposite direction of remark (2.5) is not true in general as the following example show:

**Example (2.6):**

Let

$$X = \{a, b, c, d\} \quad \tau_1 = \{X, \emptyset, \{a\}, \{b, c\}, \{a, b, c\}\}$$

$$\tau_2 = I$$
 =the indiscrete topology

$$\tau_1 - PO(X) = \left\{ X, \emptyset, \{a\}, \{b\}, \{c\}, \{a, b\}, \{a, c\}, \{a, b, d\}, \{a, c, d\}, \{a, b, c\} \right\}$$

$$\tau_1 - SPO(X) = \tau_1 - PO(X) \cup \{\{a, d\}, \{b, c, d\}\}$$

$$L - PO(X) = \tau_1 - PO(X) \cup$$

$$\{\{a, d\}, \{b, d\}, \{c, d\}, \{b, c, d\}\}$$

$$L - SPO(X) = L - PO(X)$$

If  $C = \{\{a, c\}, \{b, d\}\}$ , then C is an L-pre-open and L-semi-p-open cover, but it is neither  $\tau_1$ -pre-open nor  $\tau_1$ -semi-p-open cover.

**Definition (2.7):**

A bitopological space  $(X, \tau_1, \tau_2)$  is said to be :

- 1- “L-pre-compact space ” if and only if every L-pre-open cover of X has a finite sub cover.
- 2- “L-semi-p-compact space ” if and only if every L-semi-p-open cover of X has a finite sub cover.

**Proposition (2.8):**

- 1- Every L-semi-p-compact space is an L-pre-compact.
- 2- Every L-pre-compact space is an L-compact.
- 3- Every L-semi-p-compact space is an L-compact.

**Proof:**

Follows from remark (2.3).

**Remark (2.9):**

The opposite direction of each case in proposition (2.8) is not true in general. As the following two examples show:

- 1- Let X be an infinite set with two topologies  $\tau_1 = I$  and  $\tau_2 = D$

$$L-O(X) = \{X, \phi\}, L-PO(X) = \mathbb{P}(X) \text{ and } L-SPO(X) = \mathbb{P}(X)$$

Note that  $(X, \tau_1, \tau_2)$  is an L-compact space but it is neither L-pre-compact space nor L-semi-p-compact.

Let  $X=N$  with two topologies  $L-SPO(N) = \mathbb{P}(N)$

Note that  $(N, \tau_1, \tau_2)$  is an L-pre-compact space, but it is not L-semi-p-compact.

**Proposition (2.10):**

Let  $(X, \tau_1, \tau_2)$  be abitopological

$$\tau_1 = \{u \subseteq N : 2 \notin u\} \cup \{N\}$$

$$\tau_2 = D$$

$$L-O(N) = \tau_1$$

$$L-PO(N) = \tau_1$$

space. If

- 1- X is an L-pre-compact space, then  $(X, \tau_1)$  is pre-compact space.
- 2- X is an L-semi-p-compact space, then  $(X, \tau_1)$  is semi-p-compact space.

**Proof:**

follows from remark (2.5).

**Remark (2.11):**

The opposite direction of each case in proposition (2.10) is not true in general.

As the following example show:

Let  $X = N =$  The set of natural numbers

$$\tau_1 = \{u \subseteq N : 1 \notin u\} \cup \{N\}$$

$$\tau_2 = I$$

$$\tau_1-PO(N) = \tau_1$$

$$L-PO(N) = \mathbb{P}(N) \setminus \{1\}$$

Note that  $(N, \tau_1)$  is pre-compact space, but  $(N, \tau_1, \tau_2)$  is not L-pre-compact space.

**Proposition (2.12):**

An L-pre-closed (L-semi-p-closed) subset of an L-pre-compact (L-semi-p-compact) space is an L-pre-compact (L-semi-p-compact) set respectively

**Proof:**

Let A be an L-pre-(L-semi-p-) closed subset of an L-pre- (L-semi-p-) compact space  $(X, \tau_1, \tau_2)$  and let  $\{G_\alpha : \alpha \in \Lambda\}$  be an L-pre-(L-semi-p-) open cover of A. Then  $\{G_\alpha : \alpha \in \Lambda\} \cup A^c$  forms an L-pre-(L-

semi-p-) open cover of  $X$  which is L-pre- (L-semi-p-) compact space. So there are finitely many elements  $\alpha_1, \alpha_2, \dots, \alpha_n$  such that  $X = \bigcup_{i=1}^n G_{\alpha_i} \cup A^c$ , it follows that  $A \subseteq \bigcup_{i=1}^n G_{\alpha_i}$ . Hence  $A$  is an L-pre-(L-semi-p-) compact.

**Corollaries (2.13):**

- 1- An L-pre-closed (L-semi-p-closed) subset of an L-pre- compact(L-semi-p-compact) space is an L- compact.
- 2- An L-semi-p-closed subset of an L-semi -p- compact space is an L-pre-compact.

**Proof :**

follows from propositions (2.12) and (2.8).

**Corollaries (2.14):**

- 1- An L-pre-closed (L-semi-p-closed) subset of an L-pre- compact (L-semi-p-compact) space is a  $\tau_1$ -pre-compact( $\tau_1$ -semi-p-compact) respectively.
- 2- An L-semi-p-closed subset of an L-semi -p- compact space is a  $\tau_1$ -pre-compact.
- 3- An L-pre-closed (L-semi-p-closed) subset of an L-pre- compact(L-semi-p-compact) space is a  $\tau_1$ - compact.

**Proof :**

follows from proposition (2.12),remarks (2.3) and (2.5).

**Definition (2.15):**

A bitopological space  $(X, \tau_1, \tau_2)$  is said to be :

- 1. “L- $T_2$  -space” if and only if for each pair of distinct points  $x$  and  $y$  in  $X$ ,there exist two disjoint L-open subset  $G$  and  $H$  of  $X$  such that  $x \in G$  and  $y \in H$  .[5]

- 2. “L-pre- $T_2$ -space” if and only if for each pair of distinct points  $x$  and  $y$ ,there are two disjoint L-pre-open subsets  $U$  and  $V$  of  $X$  such that  $x \in U$  and  $y \in V$  .

- 3. “L-semi-p- $T_2$ -space” if and only if for each pair of distinct points  $x$  and  $y$ ,there are two disjoint L-semi-p-open subsets  $U$  and  $V$  of  $X$  such that  $x \in U$  and  $y \in V$  .

**Remark (2.16):**

An L-pre- compact subset of an L-pre - $T_2$ -space need not be L-pre-closed.

**For example:**

$$X = \{1,2,3\}$$

$$\tau_1 = \{X, \phi, \{1,2\}\}$$

$$\tau_2 = \{X, \phi, \{1\}, \{3\}, \{1,3\}\}$$

$$L - O(X) = \{X, \phi, \{1,2\}\}$$

$$L - PO(X) = L - O(X) \cup \{\{1\}, \{2\}, \{2,3\}, \{1,3\}\}$$

Note that  $(X, \tau_1, \tau_2)$  is an L-pre - $T_2$ -space.

Let  $A = \{1,2\}$ , clear that  $A$  is an L-pre- compact subset of  $X$ , but it is not L-pre-closed.

**Remark (2.17):**

An L-semi -p- compact subset of an L-semi-p- $T_2$ -space need not be L-semi-p-closed.

**For example:**

Note that  $(X, \tau_1, \tau_2)$  is an L-semi-p - $T_2$ -space.

Let  $A = \{1,2,4\}$ , clear that  $A$  is an L-semi-p- compact subset of  $X$ , but it is not L -semi-p-closed.

**Definition (2.18):**

$$\begin{aligned}
 X &= \{1,2,3,4\} \\
 \tau_1 &= \{X, \phi, \{1\}, \{2\}, \{1,2\}\} \\
 \tau_2 &= D \\
 L-O(X) &= \{X, \phi, \{1\}, \{2\}, \{1,2\}\} \\
 L-PO(X) &= \{X, \phi, \{1\}, \{2\}, \{1,2\}, \{1,2,3\}, \{1,2,4\}\} \\
 L-SPO(X) &= L-PO(X) \cup \\
 &\quad \left\{ \{2,3,4\}, \{1,3\}, \{1,4\}, \{2,3\}, \{2,4\}, \{1,3,4\} \right\}
 \end{aligned}$$

$$\text{Let } f : (X, \tau_1, \tau_2) \rightarrow (Y, \tau_1', \tau_2')$$

be any function, then f is said to be:

1. "L-continuous" function if and only if the inverse image of any L-open subset of Y is an L-open subset of X.[5]
2. "L-pre-irresolute" function if and only if the inverse image of an L-pre-open subset of Y is an L-pre-open subset of X .
3. "L-semi-p-irresolute" function if and only if the inverse image of an L-semi-p-open subset of Y is an L-semi-p-open subset of X .

**Proposition (2.19):**

The L-pre-irresolute (L-semi-p-irresolute ) image of an L-pre-compact (L-semi-p-compact) space is an L-pre-compact (L-semi-p-compact) respectively.

**Proof:**

Suppose that  $f : (X, \tau_1, \tau_2) \rightarrow (Y, \tau_1', \tau_2')$  is an L-pre-(L-semi-p-) irresolut and onto function and  $X$  is an L-pre-(L-semi-p-) compact space. Let  $\{G_\alpha : \alpha \in \Delta\}$  be an L-pre-(L-semi-p-) open cover of  $Y$ , it follows that  $\{f^{-1}(G_\alpha) : \alpha \in \Delta\}$  is an L-pre-(L-semi-p-) open cover of  $X$  which is L-pre-(L-semi-p-) compact. So there are finitely many elements  $\alpha_1, \alpha_2, \alpha_3, \dots, \alpha_n$  such that  $X = \bigcup_{i=1}^n f^{-1}(G_{\alpha_i}) = f^{-1}(\bigcup_{i=1}^n G_{\alpha_i})$

.Therefore  $Y = \bigcup_{i=1}^n G_{\alpha_i}$  Hence  $Y$  is an Lpre-(L-semi-p-) compact.

**Proposition(2.20):**

The L-continuous image of an L-compact space is an L-compact.

**Proof:**

Suppose that  $f : (X, \tau_1, \tau_2) \rightarrow (Y, \tau_1', \tau_2')$  is an L-continuous and onto function and  $X$  is an L-compact space. Let  $\{G_\alpha : \alpha \in \Delta\}$  be an L-open cover of  $Y$ , it follows that  $\{f^{-1}(G_\alpha) : \alpha \in \Delta\}$  is an L-open cover of  $X$  which is L-compact. So there are finitely many elements  $\alpha_1, \alpha_2, \alpha_3, \dots, \alpha_n$  such that  $X = \bigcup_{i=1}^n f^{-1}(G_{\alpha_i}) = f^{-1}(\bigcup_{i=1}^n G_{\alpha_i})$  .Therefore  $Y = \bigcup_{i=1}^n G_{\alpha_i}$  ,hence  $Y$  is an L-compact.

**Proposition (2.21):**

The L-continuous image of an L-pre-compact (L-semi-p-compact) space is an L-pre-compact

**Proof:**

follows from proposition (2.20) and (2.8).

**Proposition (2.22) :**

The L-pre-irresolute image of an L-semi-p-compact space is an L-pre-compact.

**Proof:**

follows from proposition (2.19) and (2.8).

**Theorem (2.23):**

Let  $(X, \tau_1, \tau_2)$  be abitopological space and let A be a subset of X. A point x in X is an L-pre-closure (L-semi-p-closure) point of A if and only if every L-pre-neighbourhood (L-semi-p-neighbourhood) of x intersects A.

**Proof:**

Assum that  $x$  is an L-pre-closure (L-semi-p-closure) of  $A$ , then

$$x \in \mathfrak{S} = \bigcap \left\{ \begin{array}{l} F \subseteq X : A \subseteq F \\ \text{and } F \text{ is an L-pre-closed} \\ (L\text{-semi-p-closed}) \end{array} \right\}.$$

Suppose that there exists an L-pre-neighbourhood (L-semi-p-neighbourhood)  $M$  of  $x$  such that  $M \cap A = \emptyset$ , that is, there exists an L-pre-open(L-semi-p-open) set  $G$  such that  $x \in G \subseteq M$ , then such that  $A \subseteq M^c \subseteq G^c$ , but  $G^c$  is an L-pre-closed (L-semi-p-closed) with  $x \notin G^c$ . Therefore  $x \notin \mathfrak{S}$  which is a contradiction hence every L-pre-neighbourhood (L-semi-p-neighbourhood) of  $x$  must intersect  $A$ .

Conversely

Assume that every L-pre-neighbourhood (L-semi-p-neighbourhood) of  $x$  intersects  $A$ , and suppose that  $x$  is not L-pre-closure (L-semi-p-closure) point of  $A$ , then  $x \notin \mathfrak{S}$ , that is, there exists an L-pre-closed (L-semi-p-closed) subset  $F$  of  $X$  with  $A \subseteq F$  such that  $x \notin F$ , it follows that  $x \in F^c$  which is an L-pre-open(L-semi-p-open) set. Now there is an L-pre-neighbourhood (L-semi-p-neighbourhood)  $F^c$  of  $x$  with  $A \cap F^c = \emptyset$ . that implies to contradiction with our assumption. Hence  $x$  must be an L-pre-(L-semi-p-) closure point of  $A$

**Theorem (2.24):**

Let  $(X, \tau_1, \tau_2)$  be a bitopological space. A subset  $A$  of  $X$  is an L-pre-(L-semi-p-) closed if and only if  $A = L-Pcl(A)(L-SPcl(A))$ .

Proof:

Suppose that  $A \in L-PC(X)(L-SPC(X))$

and  $A \neq L-Pcl(A)(L-SPcl(A))$ .

Since  $A \subseteq L-Pcl(A)(L-SPcl(A))$ , so

$L-Pcl(A)(L-SPcl(A)) \not\subseteq A$ , that is, there exists an element  $r \in L-Pcl(A)(L-SPcl(A))$  and

$r \notin A$ , it follows that  $r \in A^c$  which is an L-pre-(L-semi-p-) open set. Then by theorem (2.23)

$A \cap A^c \neq \emptyset$  which is a contradiction with the fact  $A \cap A^c = \emptyset$ . Hence  $A = L-Pcl(A)(L-SPcl(A))$

Conversly

Assume that  $A = L-Pcl(A)(L-SPcl(A))$ , but  $L-Pcl(A)(L-SPcl(A))$  is an L-pre-(L-semi-p-) closed subset of  $X$  by definition of L-pre-(L-semi-p-) closure of a set  $A$  which is the intersection of all L-pre-(L-semi-p-) closed subsets of  $X$  containing  $A$ . So  $A$  is an L-pre-(L-semi-p-) closed set.

**Definition (2.25):**

Let  $(X, \tau_1, \tau_2)$  be a bitopological space and let  $(f, X, A, \geq)$  be a net in  $X$ . Then  $f$  is said to be:

- 1- "L-pre-convergent" to a point  $x_o$  in  $X$  if and only if for each L-pre-nhd.  $M$  of  $x_o$  there exists an element  $a_o \in A$  such that  $f_a \in M$  for each  $a \geq a_o$ .
- 2- "L-semi-p-convergent" to a point  $x_o$  in  $X$  if and only if for each L-semi-p-nhd.  $M$  of  $x_o$  there exists an element  $a_o \in A$  such that  $f_a \in M$  for each  $a \geq a_o$ .

**Definition (2.26):**

Let  $(X, \tau_1, \tau_2)$  be a bitopological space and let  $(f, X, A, \geq)$  be a net in  $X$ . A point  $x_o$  in  $X$  is called:

- 1- "L-pre-cluster point" of  $f$  if and only if for each  $a \in A$  and for each L-



pre-nhd.  $M$  of  $\mathcal{X}_o$  there exists an element  $b \geq a$  in  $A$  such that  $f_b \in M$ .

2- “L-semi-p-cluster point” of  $f$  if and only if for each  $a \in A$  and for each L-semi-p-nhd.  $M$  of  $\mathcal{X}_o$  there exists an element  $b \geq a$  in  $A$  such that  $f_b \in M$ .

**Theorem (2.27):**

Let  $(X, \tau_1, \tau_2)$  be a bitopological space and let  $(f, X, A, \geq)$  be a net in  $X$ . for each  $a \in A$ , let  $K_a = \{f_x : x \geq a \text{ in } A\}$ , then a point  $p$  of  $X$  is an L-pre-cluster(L-semi-p-cluster) point of  $f$  if and only if  $p \in L-Pcl(K_a) \setminus (L-SPcl(K_a))$ .

**Proof:**

Assum that  $p$  is an L-pre-(L-semi-p-) cluster point of  $f$  and let  $M$  be an L-pre-(L-semi-p-) nhd. of  $p$ , then for each  $a \in A$ , there exists an element  $x \geq a$  in  $A$  such that  $f_x \in M$ . hence  $K_a \cap M \neq \emptyset$  for each  $a \in A$ . So by theorem (2.23)

$$p \in L-Pcl(K_a) \setminus (L-SPcl(K_a)) \text{ for each } a \in A.$$

Conversely

Assum that  $p \in L-Pcl(K_a) \setminus (L-SPcl(K_a))$  for each  $a \in A$ , and suppose, if possible,  $p$  is not an L-pre-(L-semi-p-) cluster point of  $f$ , then there exists an L-pre-(L-semi-p-) nhd.  $M$  of  $p$  and an element  $a \in A$  such that  $f_x \notin M$  for every  $x \geq a$  in  $A$ . this implies that  $K_a \cap M = \emptyset$ , it follows that  $p \notin L-Pcl(K_a) \setminus (L-SPcl(K_a))$  for this  $a$  which is a contradiction. Hence  $p$  must be an L-pre-(L-semi-p-) cluster point of the net  $f$ .

**Definition (2.28):**

Let  $(X, \tau_1, \tau_2)$  be a bitopological space and let  $\mathbf{F}$  be a filter on  $X$ . A point  $x$  in  $X$  is called

1- An” L-pre-cluster” point of  $\mathbf{F}$  if and only if each L-pre-nhd.  $M$  of  $x$  intersects every member of  $\mathbf{F}$ .

2- An” L-semi-p-cluster” point of  $\mathbf{F}$  if and only if each L-semi-p-nhd.  $M$  of  $x$  intersects every member of  $\mathbf{F}$ .

**Theorem (2.29):**

Let  $(X, \tau_1, \tau_2)$  be a bitopological space and let  $\mathbf{F}$  be a filter on  $X$ . A point  $p$  in  $X$  is an L-pre-(L-semi-p-) cluster point of  $\mathbf{F}$  if and only if  $p \in L-Pcl(F) \setminus (L-SPcl(F))$ , for each  $F \in \mathbf{F}$ .

**Proof:**

Suppose that  $p$  is an L-pre-(L-semi-p-) cluster point of  $\mathbf{F}$ , then each L-pre-(L-semi-p-) nhd.  $M$  of  $p$ ,  $F \cap M \neq \emptyset$  for every  $F \in \mathbf{F}$ . It follows by theorem (2.23) that  $p \in L-Pcl(F) \setminus (L-SPcl(F))$  for each  $F \in \mathbf{F}$ .

Conversly

Assum that  $p \in L-Pcl(F) \setminus (L-SPcl(F))$  for each  $F \in \mathbf{F}$ , then by theorem (2.23) every L-pre-(L-semi-p-) nhd. of  $p$  intersects  $F$  for each  $F \in \mathbf{F}$ . Hence  $p$  is an L-pre-(L-semi-p-) cluster point of  $\mathbf{F}$ .

**Theorem (2.30):[6]**

Let  $\mathcal{A}$  be a non empty collection of subsets of a set  $X$  such that  $\mathcal{A}$  has the FIP. Then there exists an ultrafilter  $\mathbf{F}$  containing  $\mathcal{A}$ .

**Remark (2.31): [6]**

Every filter in a non- empty set  $X$  has the FIP.

**Theorem (2.32):**

Let  $(X, \tau_1, \tau_2)$  be a bitopological space. Then the following statements are equivalent

1-  $X$  an L-pre-(L-semi-p-) compact space,

2- Every collection of an L-pre-(L-semi-p-) closed subsets of X with FIP has a non empty intersection ,and

3- Every filter on X has an L-pre-(L-semi-p-) cluster point

**Proof:**

1→2

Let  $\{F_\alpha : \alpha \in \Lambda\}$  be a collection of L-pre-(L-semi-p-) closed subset of X with the FIP . suppose that  $\bigcap_{\alpha \in \Lambda} F_\alpha = \phi$ , it follows by De-Morgen

Laws that  $\bigcup_{\alpha \in \Lambda} F_\alpha^c = X$  where  $F_\alpha^c$  is an

L-pre-(L-semi-p-) open set for each  $\alpha \in \Lambda$  .therefore  $\{F_\alpha^c : \alpha \in \Lambda\}$  forms an

L-pre-(L-semi-p-) open cover for X which is an L-pre-(L-semi-p-) compact space, then there exist finitely many elements  $\alpha_1, \alpha_2, \dots, \alpha_n$  such that

$\bigcup_{i=1}^n F_{\alpha_i}^c = X$  . Again by De-Morgen

Laws we have that  $\bigcap_{i=1}^n F_{\alpha_i} = \phi$  which

is a contradiction since  $\{F_\alpha : \alpha \in \Lambda\}$

has the FIP. Hence  $\bigcap_{\alpha \in \Lambda} F_\alpha = \phi$

2→3

Let  $\mathbf{F}$  be a filter on X, then by remark(2.31)  $\mathbf{F}$  has the FIP, it follows that the collection  $\{L-Pcl(F)(L-SPcl(F)) : F \in \mathbf{F}\}$  of L-pre-(L-semi-p-) closed subsets of X also has the FIP, so by (2) there exists at least one point  $x \in \bigcap \{L-Pcl(F)(L-SPcl(F)) : F \in \mathbf{F}\}$  then by theorem (2.29) x is an L-pre-(L-semi-p-) cluster point of  $\mathbf{F}$  . thus every filter on X has an L-pre-(L-semi-p-) cluster point.

3→1

Assume that every filter on X has an L-pre-(L-semi-p-) -cluster point. To show that X is an L-pre-(L-semi-p-) compact space and let  $\mathfrak{S}$  be an L-pre-

(L-semi-p-) open cover of X and suppose ,if possible,  $\mathfrak{S}$  has no finite sub cover the collection  $A = \{X - G : G \in \mathfrak{S}\}$  has the FIP, for if there is a finite sub collection  $\{X - G_i : 1 \leq i \leq n\}$  of A such that  $\bigcap \{X - G_i : 1 \leq i \leq n\} = \phi$  this implies that  $\bigcup \{G_i : 1 \leq i \leq n\} = X$  which is

contradicts our supposition that  $\mathfrak{S}$  has no finite sub cover, thus A must have the FIP, it follows by theorem (2.30) that there exists an ultrafilter  $\mathbf{F}$  on X containing A .by (3)  $\mathbf{F}$  has an L-pre-(L-semi-p-) cluster point  $x \in X$  , then by theorem (3.39)

$x \in L-Pcl(\mathbf{F})(L-SPcl(\mathbf{F}))$  for each  $F \in \mathbf{F}$ , in particular  $x \in L-Pcl(X - G)(L-SPcl(X - G))$

for each  $G \in \mathfrak{S}$ . But X-G is an L-pre-(L-semi-p-) closed subset of X for each  $G \in \mathfrak{S}$ , therefore by proposition (2.24)

$L-Pcl(X - G)(L-SPcl(X - G)) = X - G$  for every  $G \in \mathfrak{S}$ . This implies  $x \in \bigcap \{X - G : G \in \mathfrak{S}\}$ , so by De-Morgen Laws  $x \in X - \bigcup \{G : G \in \mathfrak{S}\}$ , that is,  $x \notin \bigcup \{G : G \in \mathfrak{S}\}$ , which is a contradiction with the fact that  $\mathfrak{S}$  is an L-pre-(L-semi-p-) open cover of X ,hence  $\mathfrak{S}$  must have a finite sub cover and consequently X is an L-pre-(L-semi-p-) compact space.

**Theorem (2.33):**

Let  $(X, \tau_1, \tau_2)$  be a bitopological space, if X is an L-pre-(L-semi-p-) compact space, then every net in X has an L-pre-(L-semi-p-) cluster point

**Proof:**

let  $(f, X, A, \geq)$  be a net in X. for each  $a \in A$  let  $K_a = \{f_x : x \geq a \text{ in } A\}$ . Since A is directed by  $\geq$ , so the collection  $\{K_a : a \in A\}$  has the FIP.

Hence

$\{L-Pcl(K_a)(L-SPcl(K_a)) : a \in A\}$  also has FIP , it follows by theorem (2.32)



that

$$\bigcap \{L-Pcl(K_a)(L-SPcl(K_a)): a \in A\} \neq \emptyset.$$

Let

$$p \in \bigcap \{L-Pcl(K_a)(L-SPcl(K_a)): a \in A\} \neq \emptyset$$

$$, \text{ then } p \in L-Pcl(K_a)(L-SPcl(K_a))$$

for each  $a \in A$ , so by theorem (3.37)  $p$  is an L-pre-(L-semi-p-) cluster point of  $f$ .

### References:

- [1] Kelly, J.C., 1963, "Bitopological spaces", Proc. London Math. Soc. 13, p.p.71-89.
- [2] Navalagi, G.B., 2000, "Definition Bank in General Topology", which is available at Topology Atlas – Survey Articles Section.
- [3] Nasir, A.I., 2005, "some Kinds of strongly Compact and Pair- wise

compact Spaces" M.Sc. Thesis, College of Education Ibn Al-Haitham, University of Baghdad.

- [4] Al-Khazaraji, R.B., 2004, "On Semi-p-open Sets", M.Sc. Thesis, College of Education Ibn Al-Haitham, University of Baghdad.
- [5] AL-Talkhany, Y.K., 2001, "Separation Axioms in Bitopological spaces", Research submitted to college of Education Babylon University as a partial Fulfillment of the Requirement for Degree of master of science in Math..
- [6] Sharma, L.J.N., 2000, "Topology", Krishna Prakashan Media (P) Ltd, India, Twenty Fifth Edition.

## فضاءات الرص من النوع (L-pre- and L-semi-P-)

سعاد جدعان جاسم\*

نرجس عبد الجبار داود\*

\*قسم الرياضيات - كلية التربية- ابن الهيثم - جامعة بغداد

### الخلاصة:

الغرض من هذا البحث هو دراسة انواع جديدة من التراص في الفضاءات التبولوجية الثنائية . اذ سنقدم التراص من النوع (L-pre- and L-semi-P-)