# The effect of depression on the outcome of infertile females underwent the first In Vitro Fertilization procedure

Ali Abbas Ibrahim\*, Abeer Abdulhadi Rashid\*

\*Clinical Pharmacy department, College of Pharmacy, Mustansiriyah University, Baghdad, Iraq.

Article Info:

Received Sept 2023 Revised Oct 2023 Accepted Nov 2023 Corresponding Author email: aliabas200954@mustansiriyah.edu.iq Orcid: https://orcid.org/0009-0000-1632-0226 DOI:<u>https://doi.org/10.32947/ajps.v24i4.1086</u> Abstract:

**Background:** Infertility affects married adults, and In Vitro Fertilization (IVF) is an Assisted Reproductive Technology (ART) that can treat it. Women undergoing IVF are more likely to experience depression. There is a need to reduce depression by supporting and advising sufferers. The aim of this study is to evaluate the effect of depression on pregnancy rates among infertile females underwent IVF treatment.

#### **Materials and Methods:**

This observational study analyzed 149 patients attending the infertility center at Kamal Al-Samaraie hospital seeking a chance of a conception. The participants received a traditional hospital protocol. The study assessed depression status using the CES-D (Center for Epidemiologic Studies Depression) scale at the beginning of the IVF cycle.

# **Results:**

The mean age of women included in the study was  $30.30 \pm 5.65$  years with a mean body mass index of  $27.03 \pm 3.4$  kg/m<sup>2</sup>. The clinical pregnancy rate was 38.9%. The current study showed that there was no significant relationship between the risk of depression and pregnancy rate (P-value = 0.304).

#### **Conclusion:**

Our findings show that the pregnancy rate was not related to baseline depression of women underwent the first IVF procedure.

**Keywords:** Center for Epidemiologic Studies Depression (CES-D), In Vitro Fertilization (IVF), depression, Assisted Reproductive Technology (ART).

تأثير الاكتئاب على نتائج النساء المصابات بالعقم اللاتي يخضعنَّ لأول عملية تلقيح صناعي علي عباس ابراهيم , عبير عبد الهادي رشيد \* حُليةُ الصيدلة، الجامعة المستنصرية، فرع الصيدلة السريرية، العراق.

الخلاصة:

 $\odot$ 

**خلفية:** يؤثر العقم على البالغين المتزوجين، والتخصيب داخل المختبر هو تقنية مساعدة على الانجاب يمكنها علاجه. النساء اللاتي يخضعن لعملية التلقيح الاصطناعي أكثر عرضة للإصابة بالأكتئاب. هناك حاجة للحد من الاكتئاب من خلال دعم وتقديم المشورة للمرضى. الهدف من هذه الدراسة هو تقييم تأثير الاكتئاب على معدلات الحمل لدى الاناث المصابات بالعقم اثناء عملية اطفال الانابيب.

الادوات والطرق: قامت هذه الدراسة الرصدية بتحليل 149 مريضة يترددن على مركز العقم في مستشفى كمال السامرائي الباحثين عن فرصة للأنجاب. تلقى المشاركات البروتوكول التقليدي للمستشفى. قامت الدراسة بتقييم حالة الاكتئاب باستخدام مقياس مركز الدراسات الوبائية للأكتئاب (CES-D) في بداية دورة التلقيح الاصطناعي.

AJPS (2024)

388

AJPS is licensed under a Creative Commons Attribution 4.0 International License

النتيجة: كان متوسط اعمار النساء المشمولات في الدراسة (30,30 ± 5,65 سنة) مع متوسط مؤشر كتلة الجسم (27,03 ± 3,4 ي وبلغ معدل الحمل السريري (38,9%). اظهرت الدراسة الحالية لا توجد علاقة ذات دلالة إحصائية بين خطر الأكتئاب ومعدل الحمل (0.304 = P-value). الاستنتاج: تظهر النتائج التي توصلنا إليها أن معدل الحمل لم يكن مرتبطا بالأكتئاب الاساسي لدى النساء اللاتي خضعن لإجراء التلقيح الصناعي الأول.

الكلمات المفتاحية: عملية التلقيح الاصطناعي، مركز الدراسات الوبائية للأكتئاب, الأكتئاب, تكنلوجية الانجاب المساعد.

# Introduction

The inability to conceive after one year of regular, unprotected sexual activity may indicate male or female infertility<sup>[1][2]</sup>. Couples often view infertility as a major life experience. Infertility may be seen as a cause of anxiety, worry, grief, and disappointment<sup>[3,4]</sup>. While it is estimated that 15% of couples worldwide experience infertility, this number rises to over 30% in some underdeveloped countries and to between 17 and 28% in industrialized nations<sup>[5]</sup>.

There are numerous factors that can contribute to the occurrence of infertility. For about a quarter of all couples, the etiology remains unknown. Some factors, including ovulatory disorders (25%), endometriosis (15%), pelvic adhesions (12%), tubal blockage (11%), other tubal abnormalities (11%), and hyperprolactinemia (7%), together accounted for 81% of all cases of female infertility<sup>[6,7]</sup>.

Age has a negative impact on the fertility of women, and the prevalence of age-related infertility rises as their reproductive years decrease. Fertility declines with age because of a decline in sexual activity<sup>[8]</sup>.

Assisted Reproductive Technology (ART) is a fertility treatment that involves combining eggs and sperm to create embryos, which are then returned to the mother. In Vitro Fertilization (IVF) is widely recognized as the predominant and highly effective method of ART<sup>[9]</sup>.

Anxiety and depression are more common in infertility patients with psychiatric illness, miscarriage, a longer infertility timeline, and a definitive medical diagnosis<sup>[10]</sup>. IVF patients suffer considerable anxiety and depression throughout pre-treatment, treatment, and the two-week wait before the pregnancy test. IVF impact on emotional distress is uncertain. A previous study showed that depression lowers IVF pregnancy rates<sup>[11]</sup>.

#### **Measurement of depression**

According to the Centre for Epidemiological Studies Depression Scale (CES-D), Nelson et al. (2008) showed that a significant percentage of infertile women experienced depression in varying degrees. Specifically, 19% of the participants reported low levels of depression, while 13% reported major depression<sup>[12]</sup>.

The CES-D scale is a self-report tool used to assess the frequency, severity, and duration of depressive symptoms across various demographics, including race, gender, and age<sup>[13]</sup>.

The depression of infertile females is a major problem which might affect the pregnancy outcome; therefore, the aim of the current study was to examine the clinical risk of depression among infertile women who underwent IVF procedures, and their possible associations with pregnancy rate.

## Patients and methods Ethical Consideration

The research proposal was fully discussed with and approved by the ethical and scientific committee in the College of Pharmacy/ Mustansiriyah University.

The agreement of health authority in Kamal Al-Samarrai hospital was approved before starting data collection.

#### AJPS (2024)



The written consent was taken from each included patient after a full explanation of the aims of the study and ensuring the patient's confidentiality.

#### Study design

This study was longitudinal observational study carried out in the period from (1<sup>st</sup> January 2022-1<sup>st</sup> July 2022), designed to examine the effect of depressive state of infertile women on outcome of IVF.

Among the many married women who were attending the infertility center at Kamal Al-Samaraie hospital seeking conception, 206 infertile ladies referred to IVF technology, of which 33 were excluded as they did not meet the inclusion criteria.

Of the one hundred seventy-three infertile women who were included in this observational study, 24 were refused and 149 agreed to participate in this study.

#### **Inclusion criteria**

The study requires females to meet the following criteria:

- 1- To be infertile means to have failed to conceive for at least one year after marriage.
- 2- To be within the reproductive age range (18–45).
- 3- To be capable of reading and comprehension.

#### **Exclusion criteria**

The patients were not eligible if they met any of the following conditions:

- 1- Patients who have had an IVF attempt fail in the past.
- 2- Very obese  $(BMI > 40)^{[14]}$ .
- 3- Patient with psychiatric issues (such as a history of psychiatric hospitalization, addiction, neurological or other progressive disease, or current use of psychiatric medications).
- 4- Individual with a condition that makes it difficult to communicate (e.g., language or hearing impairment).

The participants were evaluated and assessed by CES-D score at the beginning of admission of IVF cycle. All patients in this study received frozen embryo transfers (fertilized eggs), not fresh embryos.

### **Data collection**

The patient's data sheet included information about age, residence, duration of marriage, weight, height, and history of IVF and chronic mental disease.

The data collection for each patient took about 15–20 minutes.

# Methods and materials

#### Assessment of depression status

Using CES-D, first published by Radloff in 1977, the depressive state of each patient was determined<sup>[15]</sup>. The CES-D questionnaire is a 20-item assessment that asks patients to rate how frequently they encounter depressive symptoms, such as disturbed sleep, poor appetite, and feelings of isolation, during the previous week. Answer options vary from 0 to 3 for each question (0 = Rarely or Never, 1 = Some or Little, 2 = Moderately or Often, and 3 = Frequently or Nearly Always)<sup>[16]</sup>.

Scores can range from 0 to 60. As is typical for this instrument in clinical investigations<sup>[17,18]</sup>, high scores indicate more severely depressed symptoms. With excellent sensitivity, specificity. and internal consistency, the CES-D also gives cutoff values (16 or higher) that help identify people at risk of clinical depression. The CES-D has been utilized groups<sup>[19]</sup>. well across broad age Depression levels were measured at the start of IVF cycle only.

## Statistical analysis

The collected data was introduced into Microsoft Excel 2016 and data was loaded into SPSS statistical software (version 26). The descriptive statistics were presented through tables and graphs. Chi-square test and independent T-test were used for statistical analysis. The 95% confidence

#### AJPS (2024)





interval was established. P-value < 0.05 was considered a discrimination point for significance.

## Result

The rate of pregnancy in all included cases was 38.9%.

## Patient's demographic distribution

The demographic data of 149 infertile females exposed to IVF procedures are shown in table 1.

Table 1. Distribution of patients	, including age,	duration	of marriage,	and BMI	of total
	cases 14	9			

	age in years	duration of marriage in years	BMI		
Mean	30.3087	3.7383	27.03		
Std. Deviation	5.65912	2.18223	3.400		
Minimum	19.00	1.00	21		
Maximum	45.00	12.00	39		
Data presented as Mean, Standard deviation, Number of patients (N), Minimum and					
Maximum. BMI= body mass index					

## Assessment of Depression status

Table 2 shows that the mean of CES-D of women before IVF procedure. There were

no significant differences between pregnant and non-pregnant women in their baseline CES-D score.

Table 2. Association betweer	pregnancy outcome and	baseline CES-D
------------------------------	-----------------------	----------------

		pregnant			Non-pregnant			
		Ν	Mean	Standard	Ν	Mean	Standard	P-value*
				Deviation			Deviation	
pre-	CES-D	91	21.95	5.01	58	21.71	6.42	0.799 <sup>NS</sup>
score								
Data presented as Mean, Standard deviation, Number of patients (N) and percentage %, *								
independent T-test Statistics was used in comparison; NS (P-value >0.05) is considered not								

significant difference.

# Assessment of the risk of clinical depression

Table 3 shows that there is no significant

difference between pregnant and nonpregnant women according to the risk of clinical depression (P-value = 0.304).



	Females depression	with no risk of before IVF treatment	Females depression treatment	with risk before	of IVF	P- value <sup>*</sup>
	Ν	%	N	%		
Non-	9	9.9%	82	90.1%		0.304 <sup>NS</sup>
pregnant						
pregnant	9	15.5%	49	84.5%		
Data presented as Mean, Standard deviation, Number of patients (N) and percentage						
Chi-square test was used for statistical analysis; NS (P-value >0.05) is considered not						

 Table 3. Comparison between pregnant and non-pregnant women according to the risk of clinical depression

Discussion

significant difference.

Infertility is a significant public health issue with medical, psychological, and social implications<sup>[20]</sup>.

In Islamic nations such as Iraq, family status, especially having children, is highly valuable and essential; having a baby will stabilize the family and enhance marital happiness. In today's society, stigmas around infertility are growing. Divorce or even remarriage can occur if children are not present.

Depression is one of the most common psychological problems<sup>[21]</sup>. A person's ideas, actions, feelings, and overall sense of well-being can all be impacted by depression<sup>[22]</sup>.

All participants in the current study are between the ages of 19 and 45, which is consistent within the reproductive age range<sup>[23]</sup>.

The survey found that the mean age is  $(30.3087\pm5.659 \text{ years})$ . This may be due to the reproductive age of the Iraqi community as well as the desire to have a baby<sup>[24]</sup>.

This study shows that the mean duration of marriage was  $3.73\pm2.18$  years, with a maximum period of 12 years. The result of a previous study conducted in the Erbil Kurdistan Region of Iraq shows that about 55% of infertile females seeking infertility treatment had a marriage duration of 3-7 years<sup>[25]</sup>.

In different countries and regions, the mean duration of marriage among women seeking IVF varies. In India, the data suggests a mean of about 8 years<sup>[26]</sup>; in Korea, it is about 4 years<sup>[27]</sup>; and in Italy, more than 50% of couples would seek ART following an average duration of marriage less than 3 years<sup>[28]</sup>. Those differences are due to the social nature of each country.

Infertility patients experience depression rates comparable to cancer patients, with global symptoms similar<sup>[29]</sup>. It has been found that infertility-related stress and severe depressive symptoms have a strong connection with social worry, which may be the individual's reaction to social demand having kids<sup>[30,31]</sup>. However, for the association between IVF treatment outcome and psychological distress is vet unclear<sup>[32,33]</sup>

The result of this study showed that there is no significant difference in the depressive state between pregnant and non-pregnant women (P-value = 0.799), and also there is no significant difference in the risk of depression between them (P-value = 0.304). The findings of this study were consistent other researches showing with that depression symptoms did not affect the success of IVF treatment<sup>[34,35]</sup>. However according to the recent meta-analysis of elven studies shows that the women who did get pregnant had considerably lower depression scores than women who did not get pregnant at baseline<sup>[36]</sup>.

According to the results of this study, the success rate of a pregnancy rate of around 38.9%. This result is higher than previous

#### AJPS (2024)

392



study conducted in Iran, where 142 women underwent IVF who participated in the survey and found the clinical pregnancy rate to be 26.8%<sup>[37]</sup>.

## Limitations

When evaluating the results, it is important to take into account the limitations of the current study. First, the results may not be as broadly applicable as they may be because the study was limited to a single center. Secondly, there was a relatively small study sample. Third, we are not able to investigate the connection between depression and delivery or live birth. Fourth, psychological distress encompasses a wide range of symptoms, but the current study examined one type of psychological distress (depression). Moreover, the study only analyzed IVF, so the findings cannot be generalized to other ARTs.

# Conclusion

In our study, we concluded that the rate of pregnancy for women who undergo IVF procedures was not impacted by the depression and quality of life experienced by the patients.

# References

- 1- Neyra O. Reproductive Ethics and Family : An Argument to Cover Access to ART for the LGBTQ Community. 2021;7.
- 2- View of The correlation between the sociodemographic characteristics and some hormones with the infertility of women and men [Internet]. [cited 2023 Sep 17];Available from: https://ajps.uomustansiriyah.edu.iq/ind ex.php/AJPS/article/view/452/444
- 3- Maleki-Saghooni N, Amirian M, Sadeghi R, Latifnejad Roudsari R. Effectiveness of infertility counseling on pregnancy rate in infertile patients undergoing assisted reproductive technologies: A systematic review and meta-analysis. Int J Reprod Biomed [Internet] 2017;15(7):391–402.

Available from: https://pubmed.ncbi.nlm.nih.gov/29177 241

- 4- Ali MM. The prevalence of Depression among Infertile Women in Iraq. Karbala
  J Med [Internet] 2021 [cited 2023 Sep 17];14(2). Available from: https://www.iasj.net/iasj/article/230513
- 5- Al-Asadi JN, Hussein ZB. Depression among infertile women in Basrah, Iraq: Prevalence and risk factors. J Chinese Med Assoc 2015;78(11):673–7.
- 6- Causes of female infertility UpToDate [Internet]. [cited 2022 Apr 22];Available from: https://www.uptodate.com/contents/cau ses-of-female-infertility
- 7- Ali MH, Kareem A, Al-Kazaz A, Faisal AJ. Identification of a Methylation Pattern in the SNRPN Gene Promoter and its Association with Semen Abnormality Among Iraqi Males. Al-Mustansiriyah J Sci [Internet] 2023 [cited 2023 Nov 5];33(5):17–22.
- 8- Female age-related fertility decline. Committee Opinion No. 589. Fertil Steril [Internet] 2014 [cited 2022 Apr 22];101(3):633–4.
- 9- Assisted Reproductive Technology | IVF | MedlinePlus [Internet]. [cited 2022 Apr 22];Available from: https://medlineplus.gov/assistedreprod uctivetechnology.html
- 10-Psychological stress and infertility -UpToDate [Internet]. [cited 2022 Jan 4];Available from: https://www.uptodate.com/contents/psy chological-stress-andinfertility?search=infertility stress ivf&source=search\_result&selectedTitl e=1~150&usage\_type=default&display \_rank=1
- 11-Nct. Impact of Mindfulness a Intervention on Infertile Women Undergoing Assisted Reproductive Technique Treatments. http://libdb.csu.edu.cn:80/rwt/TCL/http s/MNXGT5UJMNRXZ7DTNFRXZ63 PM7YYG/show/NCT04419740 2020;

#### AJPS (2024)

393



- 12- Nelson CJ, Shindel AW, Naughton CK, Ohebshalom M, Mulhall JP. Prevalence and predictors of sexual problems, relationship stress, and depression in female partners of infertile couples. J Sex Med 2008;5(8):1907–14.
- 13-Macaskill P, Gatsonis C, Deeks J, Harbord R, Takwoingi Y. Cochrane handbook for systematic reviews of diagnostic test accuracy. 2010;
- 14- About Obesity & Morbid Obesity | Northside Hospital [Internet]. [cited 2023 Jun 7];Available from: https://www.northside.com/services/we ight-management-services/aboutobesity-morbid-obesity
- 15-Center for Epidemiological Studies Depression Scale (CES-D and CES-D-10) - SCIRE Professional [Internet]. [cited 2023 Jul 9];Available from: https://scireproject.com/outcome/center -for-epidemiological-studiesdepression-scale-ces-d-and-ces-d-10/
- 16- Radloff LS. The CES-D Scale: A Self-Report Depression Scale for Research in the General Population. Appl Psychol Meas [Internet] 1977;1(3):385–401. Available from: https://doi.org/10.1177/014662167700 100306
- 17-Sutter-Dallay AL, Cosnefroy O, Glatigny-Dallay E, Verdoux H, Rascle N. Evolution of perinatal depressive symptoms from pregnancy to two years postpartum in a low-risk sample: the MATQUID cohort. J Affect Disord 2012;139(1):23–9.
- 18-Vahratian A, Smith YR, Dorman M, Flynn HA. Longitudinal depressive symptoms and state anxiety among women using assisted reproductive technology. Fertil Steril 2011;95(3):1192–4.
- 19- Lewinsohn PM, Seeley JR, Roberts RE, Allen NB. Center for epidemiologic studies depression scale (CES-D) as a screening instrument for depression among community-residing older adults. Psychol Aging [Internet] 1997

[cited 2022 Nov 25];12(2):277–87. Available from: /doiLanding?doi=10.1037%2F0882-7974.12.2.277

- 20- Moeenizadeh M, Zarif H. The Efficacy of Well-Being Therapy for Depression in Infertile Women. Int J Fertil Steril [Internet] 2017 [cited 2023 Jun 6];10(4):363–70. Available from: https://pubmed.ncbi.nlm.nih.gov/28042 417/
- 21-Lorenzo EC, Kuchel GA, Kuo CL, Moffitt TE, Diniz BS. Major depression and the biological hallmarks of aging. Ageing Res Rev [Internet] 2023 [cited 2023 Jun 6];83. 22. De Zwart PL. Jeronimus BF, De Jonge P. Empirical evidence for definitions of episode, remission, recovery, relapse and recurrence in depression: a systematic Epidemiol Psychiatr review. Sci [Internet] 2019 [cited 2023 Jun 6];28(5):544-62.
- 22-Women of reproductive age (15-49 years) population (thousands) [Internet]. [cited 2023 Jun 5];Available from: https://www.who.int/data/maternalnewborn-child-adolescentageing/indicator-explorernew/mca/women-of-reproductive-age-(15-49-years)-population-(thousands)
- 23- Abdel-Hussein NH, Mohamed S. Anxiety and Depression among Couples Attending The Infertility Clinic Center in AL-Hilla. kufa J Nurs Sci [Internet] 2015 [cited 2023 Jun 6];5(3). Available from: https://www.iasj.net/iasj/article/107690
- 24- Mohammed SH, Gardi AHH. Assessment of Psychological status of Infertile Women in Erbil Kurdistan Region. kufa J Nurs Sci [Internet] 2013 [cited 2023 Jun 7];3(1). Available from: https://www.iasj.net/iasj/article/74946
- 25-Malhotra N, Bahadur A, Singh N, Kalaivani M, Mittal S. Does obesity compromise ovarian reserve markers? A clinician's perspective. Arch Gynecol

#### AJPS (2024)

394



Obstet [Internet] 2013 [cited 2023 Jun 7];287(1):161–6.

- 26- Yoo JH, Cha SH, Park CW, Kim JY, Yang KM, Song IO, et al. Comparison of mild ovarian stimulation with conventional ovarian stimulation in poor responders. Clin Exp Reprod Med 2011;38(3):159.
- 27- Chiaffarino F, Baldini MP, Scarduelli C, Bommarito F, Ambrosio S, D'Orsi C, et al. Prevalence and incidence of depressive and anxious symptoms in couples undergoing assisted reproductive treatment in an Italian infertility department. Eur J Obstet Gynecol Reprod Biol 2011;158(2):235–41.
- 28- The psychological impact of infertility: a comparison with patients with other medical conditions - PubMed [Internet]. [cited 2023 Jul 4]
- 29- Miles LM, Keitel M, Jackson M, Harris A, Licciardi F. Predictors of distress in women being treated for infertility. http://dx.doi.org/101080/02646830802 350880 [Internet] 2009 [cited 2023 Jun 9];27(3):238–57.
- 30-Lund R, Sejbaek CS, Christensen U, Schmidt L. The impact of social relations on the incidence of severe depressive symptoms among infertile women and men. Hum Reprod [Internet] 2009 [cited 2023 Jun 9];24(11):2810–20.
- 31-Cesta CE, Viktorin A, Olsson H, Johansson V, Sjölander A, Bergh C, et al. Depression, anxiety, and

antidepressant treatment in women: association with in vitro fertilization outcome. Fertil Steril [Internet] 2016 [cited 2023 Oct 21];105(6):1594-1602.e3.

- 32-Lynch CD, Sundaram R, Buck Louis GM. Biomarkers of preconception stress and the incidence of pregnancy loss. Hum Reprod [Internet] 2018 [cited 2023 Nov 1];33(4):728–35.
- 33- Anderheim L, Holter H, Bergh C, Möller A. Does psychological stress affect the outcome of in vitro fertilization? Hum Reprod [Internet] 2005 [cited 2023 Oct 21];20(10):2969– 75.
- 34- Lintsen AME, Verhaak CM, Eijkemans MJC, Smeenk JMJ, Braat DDM. Anxiety and depression have no influence on the cancellation and pregnancy rates of a first IVF or ICSI treatment. Hum Reprod [Internet] 2009 [cited 2023 Oct 21];24(5):1092–8.
- 35-Purewal S, Chapman SCE, van den Akker OBA. Depression and state anxiety scores during assisted reproductive treatment are associated with outcome: a meta-analysis. Reprod Biomed Online [Internet] 2018 [cited 2023 Nov 1];36(6):646–57.
- 36- Maroufizadeh S, Navid B, Omani-Samani R, Amini P. The effects of depression, anxiety and stress symptoms on the clinical pregnancy rate in women undergoing IVF treatment. BMC Res Notes [Internet] 2019 [cited 2023 Sep 17];12(1).

