

ISSN:1813-1646 (Print); 2664-0597 (Online) Tikrit Journal for Agricultural Sciences

Journal Homepage: http://tujas.tu.edu.iq



Mohammed A. Amean *

IRAQI

Taher A. Shujaa

Animal Production Dept., College of Agriculture, University of Tikrit, Tikrit, Iraq

KEY WORDS:

Awassi lambs, biochar, Growth performance, Characteristics carcass.

ARTICLE HISTORY:

Received: 21/10/2019 Accepted: 22/12/2019 Available online: 01/01/2020

Effect of adding biochar to ration on productive performance and some carcass characteristics of Iraqi Awassi lambs

راق

ملات الأكساد

ABSTRACT

Tikrit Journal for Agricultural Sciences

Tikrit Journal for Agricultural Sciences (TJAS)

This study was conducted at Department of Animal production farm College of Agriculture University of Tikrit from 6/12/2017 until 21/2/2018 (77days). Sixteen Awassi lambs were used, aged 4-5 months and average weight 23.38±0.14 Kg. Lambs were divided in to four groups (four lambs at each group) according to their weight, treatments distributed randomly into the groups.

Individual feeding was used which included four concentrate diet with different levels of biochar (0%, 1%, 2% and 3%).

Statistical analysis results showed that there was a significant increase (P < 0.05) in the final weight for second group (1% biochar) compared with fourth group, also result showed a significant increase (P<0.05) for second group in the average daily gain and final body gain and feed conversion ratio compared with third and fourth groups. Also significant increase (P > 0.05) in the second group in hot and cold carcass weight and in the percentage Dressing based on hot and cold weight, relative to live weight as well as empty body weight and in Rib-eye muscle area compared with the other groups.

© 2019 TJAS. College of Agriculture, Tikrit University

INTRODUCTION

Animals production is an important agricultural sector in Iraq and has its own place in the national economy. Sheep are one of the most important animals farm and the main sources of meat production in Iraq, which is increasing the nutritional awareness of the consumer accompanied by the increase in population(Al-Saigh &Al-kass, 1992) .. Researchers since a long time study the problems affected animal performance and production, they found most important is nutrition, which requires a formula suited to the needs of the animal for rapid growth and increase production especially ruminants (ARC, 1984). The use of some nutritives additives and non-nutritives additive improves growth, increases the efficiency of feed utilization and improves the properties of carcasses (Hassan, 2009).

Biochar is a non-feed additive in ruminant feed (Gerlach et al., 2013), it's refers to the carbon compounds prepared from the remains of some plants and some animal waste (Yalcin and Arol, 2002). The second step is to activate carbonated substances by exposing them to oxidizing agents such as carbon dioxide (CO2) or water vapor, which helps to burn the carbonate materials that block the pores. This helps to increase the surface area of this compound. The process of adsorption is one of the most important properties of biochar (Girgis et al., 2002).

Biochar are currently used to feed ruminants like additive because of their adsorption of gases and to provide a favorable environment for microorganisms in the rumen by increase growth of bacteria (Gerlach et al., 2013) .Biochar today are considered as non-food additives for animal

^{*} Corresponding author: E-mail: <u>Mohamad.abdulkader@yahoo.com</u>

nutrition to improve bowel function by eliminating toxins and impurities in the digestive system, increasing feed intake and improving animal growth (Strahsaker *et al.*, 1997; Villalba *et al.*, 2002). In view of the above, biochar was used in this study as non-feed additives for the Awassi lambs feed to study its effect on the productive performance and the characteristics of its carcasses.

MATERIALS AND METHODS

This study was conducted at the college of Agriculture-University of Tikrit from 6/12/2017 until 21/2/2018 (77days). Sixteen Awassi lambs were used, they were bought from local farms aged 4-5 months and average weight of 23.38 ± 0.14 Kg. Lambs divided in to four groups according to their weight, each group had four lambs, then distributed the treatments randomly on to the groups.

The Lambs feed individual with different levels of biochar (0, 1, 2 and 3) % respectively. The lambs were numbered with plastic figures then placed in cages of $1 \times 2 \text{ m}^2$ in a semi - enclosed. Each cage was equipped with plastic containers for concentrate feed, roughage and water . Straw was used as a source of roughage, and soybean meal corn with barley used as sources of protein and energy, as well as a mixture of minerals, vitamins and salt .The lambs were fed on experimental diets for two weeks as adaptation period before starting the experiment and then weighed for two days at 8 am after fasting 12 hours for the purpose of fixing the primary weight. The weighting process continued on a weekly basis before feeding in the morning using an balance weight with Iron box . Until the end of the experiment to calculate the average daily gain and the amount of feed provided to animals was adjusted according to the new weight per week, diet was provided once a day at 8:00 am by 3% of the body weight throughout the trial . 350 g straw / animal / day as a dry matter were added . The animals were in good health, with no disease or digestive disorders observed until the end of the experiment.

Slaughtering lamb

The lambs were weighed at the end of the experiment, which they were fasted for 12 hours before slaughter, while the water was left. Twelve lambs were slaughtered (three lambs from each treatment) and emptying the contents of the gut and then weighing the head, skin, gastrointestinal tract, empty gut, heart, liver, lungs, spleen, testicles, and the abdominal fat. Then recorded hot carcass weight . The carcasses were remained in the refrigerator for 24 hours at 4 ° C, then recorded the cold carcass weight (Field *et al.*,1963).

Dressing percentage

Dressing percentage(1)= weight of hot carcass to live body weight x 100

Dressing percentage (2)= weight of hot carcass to the weight of the empty animal \times 100

After cooling for 24 hours at a temperature of 4 $^{\circ}$ C to extract the percentage of recovery based on the weight of the cold carcass as follows:

Dressing percentage (3)=weight of cold carcass to live body weight x 100

Dressing percentage (4)= weight of cold carcass to empty animal weight \times 100

As reorted by (Bowman *et al.*,1968)

Longissimus dorsi Area

A long cross-section of the long dorsi (Longissimus dorsi) was drawn between the twelve and thirteenth ribs on Trace paper based on the way (Riley et al., 1966) measured by a Placom-digital Planimeter, three readings for each sample were taken to get the most accurate reading. As mentioned by (Riley *et al.*, 1966)

Thickness Fat on rib eye muscles

The thickness of the fat layer on the 12th rib of the left part of each carcass was measured by Digital Vernier and two readings were taken to obtain the most accurate reading for each sample. As mentioned by (Riley *et al.*, 1966)

Table (1): The proportions and components of the feed used in the experiment (%) and the cher	nical
composition of the feed based on $g / kg dry matter$	

Feed materials	Feed Groups					
	First 0 biochar%	Second 1 biochar%	Third 2biochar %	Fourthbioch a 3%		
Barly		30				
Soybean meal		10				
yellow corn	30					
Wheat bran	28					
Vitamins and salts	2					
	Gram\kg					
Crud protein (CP)	140.0					
Organic matter (OM)	964.40					
Ether Extract (EE)	30.10					
Crud Fibers (CF)	63.90					
Ash	35.60					
NFE Extract *	730.40					
Energy metabolism **	13.15					
NFE=Nitrogen Free Extract						
*NFE=OM-(CP+CF+EE)						
Maaf 1977 **Calculated metabolic energy by						
ME(MJ\KgDM)=0.012*CP+0.031*EE+0.005*CF+0.014*NFE						

The data were analyzed using complete random design (CRD) to study the effect of the coefficients in the studied traits. Significant differences were compared between the averages using the Duncan Multidisciplinary Test (Duncan, 1955). By using statistical program SAS (2001) according to statistical equation to the model.

 $Yij = \mu + Ti + eij$

Yij = the transaction value of j for transaction i

 μ = The general mean of the studied character

Ti = the effect of transaction i since i = 4,3,2,1

 $eij = random error distributed by natural and independent distribution with an average of zero and an equal variation of <math>\delta e^2$.

RESULTS AND DISCUSSION

Effect of adding biochar on performance of awassi lambs :

The second treatment with (1% biochar) showed significant effect (P < 0.05) in the final weight compared with the forth treatment (3% biochar), also showed significant (P < 0.05) effect in the total weight and average daily gain (16.68 kg and 216 g/day) compared with Third and forth treatment(14.35, 13.92 kg,185.75 and 180.50 g / day) respectively (Table, 2). The reason for the positive performance of The second treatment (1% biochar) was an increased on bacterial population , led to formation of several colonies called biofilm (Leng *et al*, 2012) which supported increased digestion of dry matter and organic matter cause the benefit of the diet. the significant lowered performance on the 2% and 3% of biochar treatment may be lower efficient of microorganisms and increased adsorption urea or nitrogen in the pores of the biochar make these substances unavailable for the microorganism this effected microble protein production.

Further explanation was that no significant differences between adding 1 % biochar and 0% (control), although 9 % improvement in average daily gain with adding biochar 1%, it may be due to diets contain energy and protein suitable for lambs .It may be conclude that biochar more effected with poor diets (Erickson *et al.*, 2011) and level of biochar added to the feed (Chu *et al.*, 2013) .The results

of this study were consistent with studies by Van *et al.* (2006) ; leng *et al.* (2012) ; Silivong and Preston ,(2015) ; Vongkhamch *et al.* (2018) ; Khoa and Thang. (2018) and Saroen and Preston, (2018) indicating improvement in growth rate and feed conversion by adding biochar for cattle and goats feed . There was little defferences with feed intake and feed conversion ratio between treatments.

	amaara e men)	•				
Traits Treatment	initial weight (kg)	Final weight gain (kg)	Total weight (kg)	Average daily gain (gm)	Total feed intake)Straw + center) Gm	Total Feed conversion ratio dry matter / gm increase in weight
T1	0.19 ±23.8	0.54±39.0 ab	0.51±15.20 ab	6.69±197.0 ab	14.73±1454.7	0.18±7.39 ab
T2	0.29±23.02	0.90±39.70 a	0.67±16.68 a	8.67±217.0 a	20.17±1436.37 ab	0.18±6.66 b
Т3	0.16±23.72	0.63±38.07 ab	0.61±14.35 b	7.92±185.75 b	11.05±1440.95 a	0.40±7.53 ab
T4	0.29±23.0	0.41±36.92 b	0.65±13.92 b	8.50±180.50 b	10.93±1392.92 b	0.36±7.76 a
General average ± standard error	0.14±23.38	0.39±38.42	0.38±15.04	5.01±194.81	8.88±1431.23	0.17±7.34
Significant level	N.S	*	*	*	*	*

Table (2) Effect of the use of different percentages	s of biochar on performance of awassi lambs (me	an
\pm standard error).		

*The vertically different numbers within the one character indicate that there is a significant difference below the probability level (P < 0.05).

T1 = ingredients of the diet + Zero biochar, T2 = ingredients of the diet + 1% biochar, = T3 ingredients of the diet + 2% biochar T4 = ingredients of the diet + 3% biochar.

The Effect of Adding Biochar on carcass Characteristics of Awassi lambs:

The results of the statistical analysis in table (3) showed that the second treatment was significantly higher (P <0.05) in the weight of the hot and cold carcass (20,306 and 19,793) kg respectively compared the first treatment (18.426 and 17.713 kg) respectively and the third treatment (18.280 and 17.890) Kg respectively and the fourth (18.133 and 17.756 kg), respectively. Also the results founded significant (P <0.05) differences in each of the percentage of reflux on the basis of the weight of warm relative to the live weight and the percentage of reflux on the basis of cold weight in relation to the live weight and the percentage of reflux on the basis of cold weight empty body and the percentage of recovery on the basis of cold weight compared to the weight of the empty body (52.16, 50.85, 59.68 and 58.18) % respectively compared to the first treatment (47.67, 45.81, 55.05 and 52.92) % respectively and the third treatment (48.27, 47.24, 55.61 and 54.42) % Respectively and the fourth (48.78, 47.77, 55.28 and 54.14) % respectively. Table (3) shows a significant increase (P <0.05) in the area of the ocular muscle in favor of the second treatment 10.80 cm² compared to the first treatment 9.63 cm².

The results of the statistical analysis in Table (3) indicate that there were no significant differences between the different treatments in subcutaneous fat thickness (2.43, 2.03, 2.53 and 2.43) Mm for the first to fourth treatments respectively. The increase in the weight of the carcasses for the 1% biochar may be due to the higher final weights of the lambs compared to the other treatment . These results were consistent with Al-Mahdawi (2002) which observed that the increase in the weight of the live animal caused an increase in the weight of the carcass. The reason for the increase in the

percentage of reflux for the second treatment was probably due to the increase in the final weight of the second treatment.

The superiority of the second treatment in the area of the ocular muscle was somewhat consistent with the observed increase in the weight of the warm and cold carcass (Table 3). This is an inevitable result of the high and negative correlation between 0.95 and the cold carcass weight (Al-Mahdawi, 2002). The increase in the amount of microbial protein, the increase in animal protein availability due to biofuels, and the increase in microorganisms, improve fermentation and increase the efficiency of feed intake, which is reflected in production (LongThuy *et al.*, 2018).

 Table (3) Effect of adding different level of biochar on some characteristics carcass of Awassi lambs (values represent averages ± standard error)

Traits Treatment	Final weight (kg)	Hot carcass weight (kg)	Cold carcass weight (kg)	Dressing percentage (1)	Dressing percentage (2)	Dressing percentage (3)	Dressing percentage (4)	area Longissimu Dorsi Cm²	Fat thicknes Mm
Т1	0.54±39.0	0.08 ± 18.42	0.28±17.71	0.54±47.67	0.41±45.81	0.60 ± 55.05	0.96 ± 52.92	0.38±9.63	0 20+2 43
11	ab	b	b	b	с	b	b	b	0.20=2.13
Т2	0.90±39.70	0.49 ± 20.30	0.47±19.79	0.66 ± 52.16	0.75 ± 50.85	0.40 ± 59.68	0.59 ± 58.18	0.20 ± 10.80	0 08+2 03
12	а	а	а	а	а	а	а	а	0.00=2.05
тз	0.63±38.07	0.37±18.28	0.39±17.89	0.30 ± 48.27	0.27 ± 47.24	0.55 ± 55.61	0.44 ± 54.42	0.15±9.90	0 08+2 53
15	ab	b	b	b	bc	b	b	ab	0.08±2.55
Т4	0.41±36.92	0.29±18.13	0.24±17.75	0.13±48.78	0.09±47.77	0.22±55.28	0.39±54.14	0.37±9.80	0 21+2 43
14	b	b	b	b	b	b	b	ab	0.21±2.+3
General									
average ±	0.39 ± 38.42	0.30±18.78	0.30±18.28	0.56±49.22	0.58±47.91	0.60 ± 56.41	0.65 ± 54.91	0.18±10.03	0.08 ± 2.35
standard	0.09 -0 01.2	0.000-10070	0.000-10.20	0.000-19.22	0.000-17091	0.00-0011	0.00-0.001	0110-10100	0100-2100
error Significant									
level	*	*	*	*	*	*	*	*	NS

*The vertically different numbers within the one character indicate that there is a significant difference below the probability level (P <0.05). T1 = ingredients of the diet + Zero biochar, T2 = ingredients of the bush + 1% biochar, = T3 ingredients of the diet + biochar (2%, T4 = ingredients of the diet + 3% biochar.

REFERENCES

- AL-Mahdawi,M.K.K.(2002)Effect of energy and protein source at ration on growth fattening locale lamp. Ph.d. thesis–College of Agric.and forestry Mosul Univ.(In Arabic).
- Al-Saigh, M.N. and Al-Kass, J.E. (1992). Production of sheep and goats-Dar Al-Hakima- Basrh.(In Arabic)
- ARC, (1984). The nutrient requirements of ruminant Livestock. Commonwealth Agricultural Bureau, Slough. U.K.
- Bowman, J. G.; Marshall, J. F. and Broddbent, J. B.(1968). Gentic parameters of carcass quality in Down cross sheep . Anim . Prod. 10: 183-191
- Chu, G. M.; Kim, J. H. and Kim, H. Y.(2013). Effects of bamboo charcoal on the growth performance ,blood characteristics and noxious gas emission in fattening pigs. Journal of Applied Animal Research, 41(1):48-55.
- Duncan, D. (1955). Multiple range and multiple F-test.Biometrics.11:1-24
- Erickson, P. S; Whitehouse, N. L. and Dunn, M. L.(2011). Activated carbon supplementation of dairy cow diets: effects on apparent nutrient digestibility and taste preference. J. Anim. Sci. 27: 428–434.
- Field,R.A.;Kamp,J.D. ;and Varney,W.Y.(1963).Carcass evaluation of lambs from selected sires .J. Animal Sci.,2:364-367.
- Gerlach, H.; Gerlach, A.; Schrödl, W.; Schottdorf, B. and Haufe, S. (2013). Oral application of charcoal and humic acids to dairy Cows influences *Clostridium botulinum*, blood serum antibody level and glyph sate excretion in urine. J.Clin.Toxicol.4:186.

- Girgis,B.S.; Samya, S.Y. and Ashraf, M.S.(2002). Characteristics of activated carbon from peanut hulls in relation to conditions of preparation. Materials Letters, 57: 164-172.
- Hassan, E. H. S. (2009). Utilization of growth promoters and bentonite in sheep ration. Ph. D. Thesis, Fac. of Agric. Al- Azhar Univ. Egypt .
- Khoa, N.H. and Thang, T.V. (2018). Effect of tannin in green tea by-product in combination with bio-char supplemented into basal beef cattle diet on nutrient digestibility, methane production and animal performance. Livestock Research for Rural Development.volume 27, page 58.
- Leng, R. A.; Preston, T. R. and Inthapanya, S. (2012). Biochar reduces enteric methane and improves growth and feed conversion in local "Yellow" cattle fed cassava root chips and fresh cassava foliage. Livestock Research for Rural Development. Volume 24, page24.
- LongThuy, L. T.; Preston, T. R.; Leng, R. A. and Ba,N. X. (2018). Effect of biochar and water spinach on feed intake, digestibility and N-retention in goats fed urea-treated cassava stems. Livestock Research for Rural Development. Volume 30, page 93.(http://www.lrrd.org/lrrd30/5/thuyh30093.html)
- Riley, M. L., Field, R. A. and Neims A.H. .(1966). Comparison of two methods of measuring the area of longissimus dorsi muscle . J. Anim. Sci. 25 : 587-592.
- Saroen, P. and Preston, T. R. (2018) Rice distillers byproduct and molasses-urea blocks containing biochar improved the growth performance of local Yellow cattle fed ensiled cassava roots, cassava foliage and rice straw .Livestock Research for Rural Development.Volume 27, Article #58. <u>http://www.lrrd.org/lrrd27/3/sili27058.html</u>
- SAS, (2001) SAS system under P.C. Dos . SAS institute . Ine . Cary . NC.
- Sengsouly, P. and Preston, T. R. (2016) Effect of rice-wine distillers' byproduct and biochar on growth performance and methane emissions in local "Yellow" cattle fed ensiled cassava root, urea, cassava foliage and rice straw. Livestock Research for Rural Development. Volume 28, Article #178.(<u>http://www.lrrd.org/lrrd28/10/seng28178.html</u>)
- Silivong, P. and Preston, T. R. (2015) Growth performance of goats was improved when a basal diet of foliage of *Bauhinia acuminata* was supplemented with water spinach and biochar.Livestock Research for Rural Development. Volume 27, Article #58.. (http://www.lrrd.org/lrrd27/3/sili27058.html).
- Struhsaker, T.T.; Cooney, D.O. and Siex, K.S.(1997) Adsorptive capacity of charcoals eaten by Zanzibar red colobus monkeys: Its function and its ecological and demographic consequences. International Journal of Primatology. 18: 61-72.
- Van, D. T. T.; Mui, N. T. and Ledin I. (2006) Effect of method of processing foliage of Acacia mangium and inclusion of bamboo charcoal in the diet on performance of growing goats. Anim. Feed Sci. Tech. 130:242–256.
- Villalba, J.J.; Provenza, F.D. and Banner, R.E. (2002). The influence of macronutrients and activated charcoal on the intake of sagebrush by sheep and goats. J. Anim. Sci. 80:2099-2109.
- Vongkhamch, B. : Preston, T R. ; Leng, R. A. An, L. V. and Hai, D. T. (2018). Effect of biochar on growth performance of local "Yellow" cattle fed ensiled cassava roots, fresh brewers' grains and rice straw. Livestock Research for Rural Development. Volume 30, Article #168.(<u>http://www.lrrd.org/lrrd30/9/bobby30168.html</u>)
- Yalcin, M. and Arol, A.I., (2002). Gold cyanide adsorption characteristics of activated carbon of non-coconut shell origin. Hydrometallurgy, 63: 201-206.

تأثير اضافة الفحم الحيوي الى العلائق في الاداء الانتاجي للحملان العواسية وبعض صفات ذبائحها

محمد عبدالقادر محمد امين وطاهر عبداللطيف شجاع قسم الإنتاج الحيواني - كلية الزراعة - جامعة تكريت

المستخلص

أجريت هذه الدراسة في حقول الأنتاج الحيواني في كلية الزراعة – جامعة تكريت للمدة من 2017/12/6 ولغاية 2018/2/21 ورز 2018/2/21 ورز 2018/2/22 ورز 2018/2/22 ورز 2018/2/22 ورز 2018/2/22 ورز 2018/2) ورز 2019 (70 ورز 2019) . ورز 2019 ورز 2018 ورز 2018 ورز 2018 الحملان إلى أربعة مجاميع تغذوية حسب الوزن وضمت كل مجموعة 4 حملان ثم وزعت المعاملات عشوائياً على المجاميع. واتبع في هذه التجربة نظام التغذية الفردية لعليقة واحدة مع مستويات مختلفة من الفحم الحيوي (0% , 1% , 2% , 3%) . أشارت في هذه التجربة نظام التغذية الفردية لعليقة واحدة مع مستويات مختلفة من الفحم الحيوي (0% , 1% , 2% , 3%) . أشارت نتائج التحليل الإحصائي إلى ظهور زيادة معنوية في معدل الوزن النهائي لصالح المعاملة الثانية (30.07) كغم مقارنة بالمعاملة التائية والزيادة الوزنية اليومية وكفاءة الرابعة (30.05) كغم ، كما أظهرت تحسناً معنوياً (0.05 < P) في كل من معدل الزيادة الوزنية الكلية والزيادة الوزنية اليومية وكفاءة الرابعة (30.05) كغم ، كما أظهرت تحسناً معنوياً (0.05 < P) في كل من معدل الزيادة الوزنية المعاملة الثانية (30.05) كغم مقارنة بالمعاملة الرابعة (2018) كغم ، كما أظهرت تحسناً معنوياً (0.05 < P) في كل من معدل الزيادة الوزنية الكلية والزيادة الوزنية اليومية وكفاءة الرابعة (2008) كغم ، كما أظهرت تحسناً معنوياً (0.05 < P) في كل من معدل الزيادة الوزنية الكلية والزيادة الوزنية اليومية وكفاءة الرابعة (2005) كغم ، كما أظهرت تحسناً معنوياً (20.05 < P) في كل من معدل الزيادة الوزنية الكلية والزيادة الوزنية اليومية وكفاءة والرابعة (20.05 < P) في كل من معدل الزيادة الوزنية الكلية والزيادة الوزنية معنوياً وكماءة التحويل الغذائي لصالح المعاملة الثانية معارياً والرابعة .كما أسارت النتائج الى تفوق المعاملة الثانية معنوياً والحويل الغذائي لصالح المعاملة الثانية مقارنة بالمعاملات الثالثة والرابعة .كما أسارت النتائج الى تحو والبارد الى الحي كذلك (20.05 P) في كل من وزن الذبيحة الحار وفي نسبة التصافي على أساس الوزن الدبيحة الحار الى الفارغ والبارد الى الفارغ وفي مساحة العضلة العينية معارية المعاملة الثانية مقارنة بالمعاملة الثانية مامالات .

الكلمات المفتاحية : الحملان العواسية ,الفحم الحيوي ، الأداء الإنتاجي ، صفات الذبيحة