

Causes And Management Of Neonatal Jaundice In Almosawi Pediatric Hospital In Nasiryiah City A: Retrospective Study

Dr.Abdulreda Abed Hatam (M.B.CH.B. D.C.M. Almosawi pediatric hospital)
abdulreda_ahatam@yahoo.com

Dr. Mishaal Zoori Jabbar (M.B.CH.B. D.H.A. Almosawi pediatric hospital)
mzla74@yahoo.com

Dr. Hussein Ali Hamad (MBChB, DS, CABMS)
gg99ffr@mailru

Abstract

- Background

Neonatal jaundice is the most common reason for admission in the neonatal period. The yellow coloration of the skin and sclera in newborn with jaundice is the result of accumulation of unconjugated bilirubin. Neonatal physiological jaundice results from simultaneous occurrence of two phenomena, bilirubin production is elevated and hepatic excretory capacity is low (1).The aim of this study is to determine the etiology,severity,risk factors and management of neonatal jaundice.

- Methods

A retrospective study was conducted on 100 case_sheets (random sample) to neonates who had jaundice in Almosawi paediatric hospital in Nasiryiah city.The obtained data were analysed and the results were tabulated.

- Results

100 case_sheets of neonates who have jaundice were studied.ABO blood group incompatibility were the most common cause of jaundice.Phototherapy is the mainstay of hyperbilirubinemia treatment in neonates.It has resulted in a marked reduction in the need to perform exchange transfusion.

- Conclusion

Blood group incompatibility,and unknown causes,are the most common cause of neonatal jaundice .The less common cause is G6pd deficiency.Most of neonates who have jaundice were breast feeding,lowbirth weight and preterm are risk factors.

Keyword: (jaundice; neonatal; Causes)

Introduction

Hyperbilirubinemia is common and, in most cases, a benign problem in neonates. Jaundice is observed during the 1st wk of life in approximately 60% of term infants and 80% of preterm infants. The yellow color usually results from the accumulation of unconjugated, nonpolar, lipid-soluble bilirubin pigment in the skin (2). Unconjugated free bilirubin is neurotoxic and can cause kernicterus (3). Jaundice is usually detectable clinically when the plasma bilirubin exceeds 3mg/dl (4). Severe neonatal jaundice may cause permanent brain injury and hearing loss in infants worldwide, including those born in developed countries. Bilirubin neurotoxicity is due to the effects of unbound unconjugated bilirubin in the central nervous system. The risk of kernicterus depends not only on the total serum bilirubin, but also factors such as the bilirubin-albumin binding capacity, prematurity and other neonatal illnesses (5). The amount and duration of hyperbilirubinemia and the neurodevelopmental age (preterm neonates) at the time of insult exposure is supposed to influence the location of selective brain damage as well as the severity of consequences (6). The history of jaundice is very long and described as a sign of (causeless hatred) in the Babylonian Talmud. There are various ancient references related to jaundice which are presented in Babylonian Talmud, Sumerian Tablets, Ebers Papyrus and in Ancient Ayurveda (the Indian traditional system of medicine). Moreover, the work of Hippocrates (460-370 B.C) also provides reference to jaundice (7).

Background:

Jaundice is still a leading cause of preventable brain damage, physical and mental handicap, and early death among newborns in many communities. Neonatal morbidity and mortality remain very high in the developing countries (8).

The causes of neonatal jaundice are: breastfeeding and underlying health conditions which include hypothyroidism, blood group incompatibility, rhesus factor disease, urinary tract infection, Grigler Najjar syndrome, G6pd deficiency, and sepsis. There is controversy as to whether breastfeeding increases the incidence of jaundice in the first few days of life (9).

Male sex, high birth weight, breastfeeding, warm air temperature, primiparity, skill birth attendance, place of delivery, prolonged labour, oil massage, paternal education, and ethnicity were significant risk factors. Among infants with difficulty feeding, exclusive breastfeeding was a risk factor for neonatal jaundice, whereas exclusive breastfeeding was protective among infants with no report of difficulty feeding (10).

Methods:

This retrospective study was conducted on 100 neonates with diagnosed jaundice at Almosawi hospital in Nasirya city, between January and December 2018. They were taken as a random sample. Data was collected. The

neonates were predominantly treated with phototherapy and small minority required exchange transfusion. History include the birth weight, breastfeeding, formula feeding, and family history.

Results of investigations including blood group and rh type of mother and neonate, serum bilirubin levels, c-reactive protein, blood culture, and G6pd enzyme.

Results:

In our study ABO incompatibility 43% was the most common cause. In nearly 30% the cause was not known. 13% of the treated neonates were diagnosed with sepsis. 10% had rh incompatibility, (this is severe neonatal jaundice) (11), and 4% G6pd deficiency. Two patients have kernicterus, as in table No.1. The majority of cases 89% treated by phototherapy, 11% by blood exchange, table No.2.

Phototherapy is a simple and effective way to reduce bilirubin level. Most term babies have physiological jaundice which responds to a short period of phototherapy, and require no other treatment. A few babies have rapidly rising bilirubin levels which place them at risk of kernicterus (12). It is unclear whether aggressive phototherapy to prevent neurotoxic effects of bilirubin benefits or harm infants with extremely low birth weight (1000g or less) (13).

Aggressive use of phototherapy has decreased the use of the invasive exchange transfusion method, relegating exchange transfusion to only severely emergent cases. Lack of adequate phototherapy and facilities, severity of hyperbilirubinemia complicated by genetic factors, and system deficit make exchange transfusion a more common treatment in developing countries (14).

An Iranian study in Fars province revealed that the most common causes of severe hyperbilirubinemia were sepsis, blood group incompatibility, G6pd deficiency and unknown (15).

21% of the neonates who required treatment in our study were preterm. Table No.3. Croatian study showed that the neonatal jaundice was associated with low birth weight, maternal infection, gestational age and premature rupture of membranes (16).

A study conducted in Asia documented ABO incompatibility and G6pd deficiency as leading causes of neonatal jaundice. G6pd abnormality was found in 12% of the neonates with jaundice in south Iran and 4.2% in Taiwan (17).

The prevalence of neonatal jaundice and risk factors in healthy term neonates at national district hospital in Bloemfontein. A total of 96 mother-infant pairs included in the study. The prevalence of neonatal jaundice was 55.2%, however only 10% of black babies who were diagnosed with jaundice appeared clinically jaundiced. Normal vaginal delivery was only associated with neonatal jaundice. Black races and maternal smoking were not protective against neonatal jaundice as in some other studies (18).

TABLE NO.1 Causes of neonatal jaundice

Cause	No.of cases	Kernicterus
ASO incomp	43	1
Others	30	
Sepsis	13	
Rh incomp.	10	1
G6pd def.	4	
Total	100	2

TABLE NO.2 Causes and type of treatment of neonatal jaundice

Cause	Type of treatment	
	Phototherapy	blood exchange(severe)
ABOincomp.	42	1
Others	25	5
Sepsis	12	1
Rh incomp.	8	2
G6pd def.	2	2
Total	89	11

Table No.3 Causes and No.of preterms

Cause	No.of preterms
ABOincom.	10
Others	8
Sepsis	2
Rh incomp.	1
G6pd	-
Total	21

Discussion

This study sought to identify the possible factors associated with neonatal jaundice. Majority of neonates developed jaundice within 1-3 days with 10% having it at birth. Birth weight and prolonged duration of labour were associated with neonatal jaundice. Mothers had inadequate knowledge of neonatal jaundice. (17) They usually put garlic cloves in beds and use sugar water which do not work to get rid of bilirubin (19). Most of neonates with jaundice had low birth weight, compared to those without jaundice (20). Preterm neonates who have concurrent illnesses and physiologic derangement are more vulnerable to bilirubin neurotoxicity. Bilirubin related neurotoxicity can result in neonatal death or multisystem acute manifestation and long term impairment including irreversible ethetoid palsy and speech, visumotor, auditory, and other sensory processing disability (21).

Conclusion

Neonatal jaundice is easily diagnosable, however require quick and on spot treatment. If not treated properly, it leads to kernicterus. Currently the treatment options for neonatal jaundice include phototherapy and exchange transfusions. Low birth weight and preterm labour were associated with neonatal jaundice. A more serious form of neonatal was due to Rh incompatibility.

This study conclude that the ABO incompatibility is the most common cause of neonatal jaundice in this hospital. This is followed by other causes, sepsis, Rh

incompatibility and G6pd deficiency. Understanding the aetiology and risk factors for neonatal jaundice in our setting helps in prioritizing the group of neonates who require more intensive monitoring for early identification and timely management of this condition.

The majority of the neonates admitted in the hospital were improved at discharge. Therefore early recognition of neonatal hyperbilirubinemia is an important public health concern.

Clinical experience has revealed that mothers have an inadequate understanding of the jaundice and perceive it to be far more serious than it is.

Recommendation

1- It is recommended that the bilirubin level of all babies should be checked before discharge from the hospital, because sometime the healthy term babies develop neonatal jaundice.

2- As phototherapy is a safe effective method for decreasing or preventing the rise of serum unconjugated bilirubin level and reduce the need for exchange transfusion in neonates, we suggest to provide the hospitals with conservative and aggressive phototherapy devices..

3- Rh incompatibility can be prevented by receiving preventive treatment with immunoglobulin which will prevent serious effects.

4- Education on the condition and its causes should be intensified especially by healthcare workers during regular antenatal visit. Causes of neonatal jaundice need to be examined in the routine management of neonates.

References

- 1-Devesh Tewari, Ande Journal of neonatal biology .
- 2-Namasivagon Ambalavnan and walderar cario. Nelson Textbook of pediatric, 2011, United States of America. 6-3.
- 3-Oxford handbook of paediatrics .2016.130. United states of America.
- 4-Davidson's principles and practice of medicine, 930, 2010. London.
- 5-Raye-Ann, Dergnier. MD. The journal of paediatrics April 2017, volume 183, page 2-3.
- 6-Matteo Dal Ben, Silvia Gazzin and Claudia Tiribelli. Italian journal of paediatrics 2014 volume 40 supplement 2, A10
- 7-Devesh Tewari, Andre Mocan, Emil D Parvanov, Archano N Ssab, Seyed M Nabavi, Lukasz Huminiecki, Zheny Feeima, Yeong Yeh Lee, Jaroslaw O. Horba and Atanas G Aanasov. Front Pharmacol Journal volume 8. 2017. 518.
- 8-Zupan. J. perinatal mortality in developing countries. New Engl. J. med. 2005. 352.
- 9-Maisles MJ. Gifford K. neonatal jaundice in full term infants 1983. 137.
- 10-Carolyn G. Scraftford, Luke C. Mullany, Joanne Katz, Sabarna K. Khatry, Steven C. Leclercq, Gary L. Darmstad and James M. Tielseh. Trop Med Int Health. 2013 nov, 18(11). 1317-1328.-
- 11-Davutoglu M. Garipardic M. Guler E. Karabibe H. Erhan D. The etiology of severe neonatal hyperbilirubinemia and complication of exchange transfusion. Turk J Pediatr 2010. 52.
- 12-Dr. Sabhabrata Mitra, Dr. Janet Rennie. British journal of hospital medicine, volume 78, No. 12. page 699-704.
- 13-Brenda H. Morris, M.D., William Oh, M.D., Jon E. Tyson, M.D., M.P.H., David K. Stenson, M.D., Dale L. M.D., T. Michael O Shea, M.D., M.P.H., Georgia E. McDavid, R.N., Rebecca L. Perritt, M.S., Kriss P. Van Meurs, M.D., Betty R. Vohr, M.D., Cathy Grisby, B.S.N., Qing YAD, Ph.D., Etal., for the NICHD. The New England journal of medicine. October 30, 2008, 359, 1885-1869.

- 14-Well,Lourtenage MS,FNP(BC),RN,Ahmed,Azza DNSC,CPNP,RN,IBCLC,Masser,Anne MS,PNP(BC),RN.The American journal of maternal/child nursing.november/December 2013,volume38,No.6.page 377-382.
- 15-Najih Ks,Saki F,Hemmat F,Imloo S,incidence,risk factor and cause of sever hyperbilirubinemia in the south of Iran,Fars province,Iran red crescent med! 2013 15.
- 16-Mesic I,Milas V,Medimurec M,Rimar Z.unconjugated pathological jaundice in newborns .Call Antrpol 2014.38,
- 17-Prince Adoba,Rivhard K.D.Ephraim,Kate Adoma kowcah Kontor,Joseph-Josiah,Bentail,Patrick Adu,Moxwell Anderson,Samuel Asamoah Sakyi,and Paul Neiah.International journal of paediatric volume 2018.
- 18-Henneke Birts,Jeanic Adendorff,Dyanti Huisman,Dohne Benkes,Kristian Botha,Harve Herbst,and Gina Jourbet.African journal of primary health care and family medicine.2018,10(1),1582.
- 19-Hassan Boskabadi,MD,Gholemal Maamouri,MD,and Shahin Mafinjad,MD.Iranian Journal of pediatric 2011 sep.21(3) 325-330
- 20-D.E.Omekwe,M.Duke Georg B.T.Kennis etal.Survey and mangment outcome of neonatal jaundice from developing tertiary health center,southern Nigeria.
- 21-V.K Bhutani,R.J.Wong.Bilirubin neurotoxicity inpreterm infants:risk and prevention .Journal of clinical neurology vol.2no.2 pp 61

اسباب ومعالجة اليرقان حديثي الولادة في مستشفى محمد الموسوي للأطفال في مدينة الناصرية : دراسة ارتدادية

د. مشعل زوري جبار

د.حسين علي حمد

د. عبدالرضا عبد حاتم

الخلاصة

يرقان حديثي الولادة اكثر الاسباب لرقود حديثي الولادة.اصفرار الجلد والصلب الابيض للعين بسبب تجمع البليروبين الغير مقترن.ويرقان حديثي الولادة الفسلجي بسبب زيدة انتاج البليروبين وهبوط قابلية الكبد للافراز.والهدف من هذه الدراسة لتحديد الاسباب والشدة وعوامل الخطورة ومعالجة يرقان حديثي الولادة.

دراسة استيعادية لمئة طلبة (نموذج عشوائي) لحديثي الولادة مصابين باليرقان في مستشفى محمد الموسوي للأطفال في الناصرية. جمعت البيانات وحللت ونظمت بجدول. عدم مطابقة مجاميع الدم هي اكثر الاسباب لهذا اليرقان.العلاج الضوئي له دور كبير بمعالجة الحالات وقلل بشكل كبير الحاجة لتبديل الدم. حالات كثيرة من اليرقان الولادي غير معروفة السبب،واقول الاسباب هو داء الباقلاء.واكثر المصابين هم رضاعة طبيعية.نقص الوزن والولادة المبكرة هي عوامل خطورة.