

PREPARATION AND CHARACTERIZATION OF CR (III), MO (V) AND W (VI) COMPLEXES USING SOME SCHIFF BASES

ABD ALHADY R.H. ALHITI

OMAR H. S. AL- OBAIDI,

CHEMISTRY DEPT. COLL. OF EDUCATION FOR WOMEN, AL-ANBAR UNIV



ARTICLE INFO

Received: 19 / 6 /2006
Accepted: 26 / 11 /2006
Available online: 14/6/2012
DOI: 10.37652/juaps.2007.15434

Keywords:

Preparation .
characterization .
Cr (III),
Mo (V) .
W (VI) .
Schiff bases.

ABSTRACT

This research consists of the synthesis of two Schiff base ligands whose name are as follow: phenyl -3- methyl -5- pyrazolone thiosemicarbazone = (L1) and Phenyl -3- methyl -5- pyrazolone-4- phenyl thiosemicarbazone = (L2) The complexation of the above ligands with some transition metal ions (Cr (III), Mo(V), W(VI)) afforded metal complexes which have been characterized on the basis of their (C.H.N) elemental analysis, IR, UV-VIS, and atomic absorption. The molar conductivity measurement and melting point of prepared complexes were measured to investigate the proposed structure as an octahedral.


Introduction:

The complexes of Schiff base and their derivatives have useful and importance applications, as well as, biological activities have taken a keen interest in medicine and a significant correlation between the chelating properties and anti tumor activity or various disease because it is known that some drugs have increase activity when administered as metal complexes and a number of metal chelates inhibit tumor growth⁽¹⁾.

The thiosemicarbazide and thiosemicarbazones have attached special attention due to their activity against viruses prato zoa, small pox and certain kinds of tumor^(2,3), it is known that some drugs have increased activity when administrated as metal complexes (4,5) and a number of metal chelates inhibit tumor growth⁽⁶⁾. For many years thiosemicarbazide has been known to form complexes with certain transition metal ions and most of this work has been reported by Jensen⁽⁷⁾.


Extended studies of the investigation of metal complexes of thiosemicarbazone have been given in table⁽⁸⁾. Complexes of Schiff bases with metals have not been prepared except in Ziad⁽⁹⁾ was prepared some of this complexes by reaction these ligands with metal chloride, acetate, sulphate and nitrite some transition elements.

We have, therefore, undertaken a synthetic and structural study of the above new type of complexes metal (Cr (III), Mo (V), W (VI); bidentate ligands these:



1-phenyl -3- methyl -5- pyrazolone thiosemicarbazone
= (L1)

* Corresponding author at: Chemistry dept. Coll. of Education for Women, Al-Anbar Univ
, Iraq.E-mail address:



1-Phenyl -3- methyl -5- pyrazolone-4-phenyl thiosemicarbazone = (L2)

EXPERIMENTAL:

INSTRUMENTATION:

Apye–Unicam sp3-100 infrared spectrophotometer was used to record the ir spectra as KBr disc and CsI disc , UV/VIS spectra were measured by a HITACHI U-2000 spectrophotometer, Elemental Analysis were done by atomic absorption AA-680G (Shimadzu). The elemental analysis (C.H.N) founded on (Carlo Erba micro analyzer type 1106).Electrical conductance was measured on conductivity CDC304 (Jenway4070) in DMSO solvent at (10-3M), Melting points were determined by an electric heated block apparatus (Gallen Kamp), and were uncorrected.

MATERIALS:

The hydrated metal chlorides [CrCl₃.6H₂O] , [MoCl₅.6H₂O] , [WCl₆.6H₂O] were supplied by BDH chemicals, ethanol absolute , diethyl ether, dimethyl sulfoxide , 4-phenyl thiosemicarbazide, 1-phenyl-3-methyl-5-pyrazolone and thiosemicarbazide supplied by Aldrich .

A- Preparation of (L₁), (L₂) was prepared according to the literature ⁽¹⁰⁾ as shown bellow:

Preparation of (1-phenyl-3-methyl-5-pyrazolone thiosemicarbazide) (L₁): To a hot solution of (1-phenyl-3-methyl-5-pyrazolone) (1.91 gm, 0.011

mole) in absolute ethanol (15 ml) a hot solution of (Thiosemicarbazide) (1 gm, 0.011 mole) in absolute ethanol (15 ml) was added. The resulting mixture was then refluxed for 6 hours, during which time a yellow precipitate formed then was cooled, lifted overnight at room temperature. This was collected by filtration in vacuo, washed and recrystallized from absolute ethanol/ether to give a yellow needle shaped crystalline solid.

Preparation of (1-phenyl-3-methyl-5-pyrazolone 4- phenyl thiosemicarbazide) (L₂):

To a hot solution of (1-phenyl-3-methyl-5-pyrazolone) (2.083 gm, 0.012mole) in absolute ethanol (10 ml) a hot solution of (4- phenyl thiosemicarbazide) (2 gm, 0.012 mole) in absolute ethanol (10 ml) was added. The resulting mixture was then refluxed for 6 hours, during which time the solution became bright yellow in colour, then was cooled, lifted overnight at room temperature, after this time a yellow precipitate formed. This was collected by filtration in vacuo, washed and recrystallized from absolute ethanol/ether to give a white crystalline solid.

The full name of the Schiff base will be replaced by the symbols (L₁, L₂) respectively as shown in table (1) for the rest of this paper. The physical properties of these compounds (L₁, L₂) are listed in table (1). The characters ir bands and uv/vis spectrum in DMSO as shown in table (2), (3).

B- General procedure for preparation of metal complexes :

To a hot solution of ligands (L₁ or L₂) (2 mmole) in absolute ethanol (5 ml), a hot solution of metal chloride (1 m mole) in absolute ethanol (5 ml) (dissolved in dilute HCl)⁽¹¹⁾ was added and the resultant mixture was stirred and refluxed for 2 hours, the color of the solution changed immediately, the reaction mixture was cooled, and the solution was evaporated in vacuum, or left over night at room

temperature, then precipitate formed, collected by filtration in vacuo, washed and recrystallized from absolute ethanol/ether.

The physical properties of prepared complexes are listed in table (4). The analogous complexes were prepared in a similar manner to that described above by adding a hot solution of (1 mmole) in absolute ethanol (5 ml) to a hot solution of metal chloride (1 mmole) in absolute ethanol (5ml). The molar ratio of the complexes was determined according to the methods (12).

RESULTS AND DISCUSSION:

The structures of Schiff base complexes were confirmed by spectroscopic techniques like IR and UV/visible. The infrared spectra of the two ligands (L1), (L2) show the usual broad bands in the region around (3400-3350 cm^{-1}) and (3360-3475 cm^{-1}) respectively due to the NH and NH₂ stretching frequency⁽¹³⁾ of the amide groups of the ligand, practically no effect on these frequencies after complexation precludes the possibility of complexation at this group⁽¹⁰⁾.

The band at 1640 cm^{-1} in the spectrum of (L1) due to symmetrical C=N stretching decreased in the complexes, while the band at 1620 cm^{-1} was assigned to asymmetrical C=N frequency, which shifted to the lower frequencies in the complexes (14) (table 4), the C=N stretching frequency of the (L2) observed at 1625 cm^{-1} shifted to 1600-1620 cm^{-1} in the complexes (15) (table 4). The negative shift generally in C=N further suggested the coordination to metal ions through nitrogen atom of (-C=N-) Schiff's base^(16,17) of the ligand and on complexation indicates involvement of azomethine nitrogen (6, 11, 13, 14, 18) with metal ions.

The band at 825 cm^{-1} in the spectrum of (L1) due to C=S stretching vibrations. In the metal

complexes this band is weakened and lowered⁽¹⁹⁾ (table 4). The C=S stretching vibrations of the (L2) contributed two bands at 1280 cm^{-1} and 895 cm^{-1} ⁽²⁰⁾. Also in the metal complexes these bands are weakened and lowered (table 4), the observations indicate the coordination of the ligands through sulphur atom.

New bands were appeared at low frequencies region in the spectra of the prepared complexes were probably due to (metal- nitrogen), (metal- sulphur), and (metal- chloride), vibration frequencies (table 4). The complexes give different colour from the transition metal salts and the ligands, then this was important indication to coordinate occurrence^(21,22), therefore these colour complexes show different characteristic absorption band, intensity or together when compared with the bands of ligand and this was another indication for the coordination occurrence^(23,24). The UV/visible spectra of the two prepared ligands (L1, L2) at (10-3M) in ethanol were showed two absorption bands⁽²⁵⁾. The first band between (245-259) nm represented (π - π^*) while the second band (308-310) nm represented (π - π^*) transition and called (B-band) for phenyl group^(24, 25). Generally in the new prepared complexes these bands are shifted to short or long wavelength compared with free ligands and high intensity of the bands is indicate for complexes formation (23,24) are shown in table (4).

The measurements of the molar electrical conductivity of the complexes at (10-3M) at (25°C) in di methyl sulphoxide are presented in table (4). These results show the high values of the molar conductivity, these complexes are electrolyte and low values refer to the complexes are non-electrolyte, are in agreement with the proposed structures of the complexes (26).

The continuous variation method to evaluate the mole ratio (M: L) were employed in this work, the molar ratio (1:2) metal to ligand for (1-6) complexes and (1:1) metal to ligand for (7-12) complexes.

According to the results obtained from (ir, uv/vis, molar ratio, molar conductivity, atomic absorption) measurements for the prepared complexes, the proposed molecular structure of the complexes has octahedral structure as shown below:

References:

[1].R.S. Verna, K. C. Gupta, Amarnath and V. S. Mishra, Indian. J. Microbiol., 64, 13124e (1966).
[2]. D. West, A. Liberta, K. Rajendran and I. Hall, Anticancer Drugs, 4,241(1993).
[3]. I.Cory, A. Cory, Grappa, A.Lorico, M.Liu, T.Lio and.Sartorelli, Adv.Enz.Regul. 35, 55(1995).
[4]. M.M.Mostava, A.M.Shallaby and A.A.El.Asmy, J.Inorg.Nucl.Chem, 43, 2992 (1981).
[5]. A.Papageorgeion, Z.Iakovidou, D.Mourelators, E.Mioglouand et.al, Anticancer.Res. 7, 175 (1997).
[6]. M. Ferrari, G. Fava and et.al. J.Inorg.Biochem. 70, 145 (1998).
[7]. R.K. Agarwal, H. Agarwal and I. Chakraborti, Qatar Univ. Sci.J. 14(c), 92 (1994).
[8]. J. Lim, C. Mathias and M. Green, J. Med. Chem. 40, 132(1997).
[9]. Ziad kalay, Ph.D Thesis, Almustansiryra University, (2000).
[10]. A.Quiroga, J.Perez, E.Montero and et.al. , J. Inorg. Biochem. 70, 117 (1998).
[11]. C.B.Mahto, J.Indian.Chem.Soc. 58, 935 (1981).
[12]. V.J. Baber, D.V. Khasnin and V. M. Shinde, J. Indian. Chem. Soc. 58, 970 (1981).
[13]. G.B. Mahto, J. Indian. Chem. Soc. 57, 481(1980).
[14]. D.X. West, S.A. Van. Rockel and Buntiry, Trasion Met.Chem., 13, 53 (1988).
[15]. B.Pradham and D.V.R.Rao, J.Indian.Chem.Soc. LIV, 136 (1977).
[16]. U.K.Mauthy, N.D.Jaghik and M.GPrjpe, J.Indian.Chem.Soc. 53, 419 (1976).

[17]. P.L.Maurya, B.V.Agarwala and K.Dey Arun, J.Indian.Chem.Soc. 55, 418 (1978).
[18]. P.L.Maurya, B.V.Agarwala and K.Dey Arun, Polym, Bull., 1,631 (1979).
[19]. M.M. Mostava, A.M. Shallabyand A.A. El. Asmy, J. Inorg. Nucl. Chem. 43, 2992 (1981).
[20]. D.X. West, S.A. Van. Rokel and Buntiry, Transition Met. Chem., 13, 53 (1988).
[21]. E. Russeva, V. Kuban and L. Sommer, Coll, Czech. Chem. Commun, 44,374 (1979).
[22]. B.G. Saha, R.P. Shatnagan and K. Banerji, J. Indian. Chem. Soc, LIV, 927(1982).
[23] R.G. Pearson, J. Am.Chem.Soc. 85, 3533(1963).
[24]. A.E. Gillam and E.S. Strm, "An Introduction to the Electronic Absorption Spectroscopy"2nd ed.Edward Arnold Ltd, London, P.115, 150(1957).
[25]. R.M. Silverstein and G.C. Bassler," Spectrometric Identification of Organic Compound," 2nd Ed. John Wiely and Sons Ltd .New York (1969) and 4th Ed. (1981).
[26]. B.G. Saha, R.P. Shatnaganand K. Banerji, J.Indian. Chem. Soc., LIV, 927 (1982).

Table (1): physical properties of the Schiff base ligands

No.	Yield %	M.P. °C	Elemental analysis (% cal.) % found			
			C	H	N	
L ₁	85%	174-176	52.98 (53.42)	5.37 (5.30)	28.24 (28.32)	Yellow
L ₂	87%	134-136	62.87 (63.14)	5.40 (5.30)	21.45 (21.65)	white

Table (2): The characteristic ir bands of the Schiff base ligands

No.	C=N sy. Str. cm ⁻¹	C=N asy. Str. cm ⁻¹	C=S sy. Str. cm ⁻¹

	L ₂	L ₁
	1625 vs.	1640 vs.
	-	1620 vs.
	1280 m, 895 m	825 bm

Sy=symmetrical, asy= asymmetrical, vs. = very sharp, medium, bm=broad medium,

Table (3): UV-VISIBAL absorption of the Schiff base ligands

	L ₂	L ₁	No.
	246 nm	249 nm	$\pi-\pi^*$
	310 nm	308 nm	$\pi-\pi^*$

Table (4): some physical and chemical properties of the prepared complexes

3	2	1	No. complexes	Colour	ΔM#	M.P °C	UV/VIS nm		Elemental analysis		IR SPECTRA cm ⁻¹
							M %	Cl %	(% found) % cal		
[W(L ₁) ₂ Cl ₂ Cl ₄] DG 88	[Mo(L ₁) ₂ Cl ₂ Cl ₃] B 80	[Cr(L ₁) ₂ Cl ₂ Cl] DG 65					211-213	185-187	294,310,595		
284,308,530	252,310,495	294,310,595					21.15	8.3			
24.97	24.25	17.25									
1625m	1615s	1620 bs									
1600	1590	1595									
790 bm	795 bw	800 bm									
465 bs	475 bm	460 bm									
365 m	385 w	390 bw									
310 bw	290 m	270 w									

	12	11	10	9	8	7	6	5	4
	[W(L ₂)Cl ₂ Cl ₂] DG 60	[Mo(L ₂)Cl ₄ Cl] B 56	[Cr(L ₂)Cl ₃ H ₂ O] DG 15	[W(L ₁)Cl ₄ Cl ₂] DG 65	[Mo(L ₁)Cl ₄ Cl] B 58	[Cr(L ₁)Cl ₃ H ₂ O] DG 15	[W(L ₂) ₂ Cl ₂ Cl ₄] DG 90	[Mo(L ₂) ₂ Cl ₂ Cl ₃] B 85	[Cr(L ₂) ₂ Cl ₂ Cl] DG 67
	230-232	215-217	195-197	218-220	209-211	188-190	210-213	208-210	190-193
	250,312,510	249,311,485	247,312,587	249,310,520	251,311,490	250,309,590	249,313,545	250,315,501	248,314,605
	25.95	16.41	10.67	29.06	18.86	12.65	18.07	10.73	6.68
	30.46	30.76	22.17	34.12	35.36	26.27	21.21	20.13	13.88
	-	-	-	1615	1630 bm	1625 ms	-	-	-
	1600	1615	1605	1595	1585	1590	1605	1600	1610
	1275,875	1265,855	1255,865	795	802 bm	790 m	1270,860	1260,850	1250,870
	455	460	475	460	470 m	450 bs	450 bm	470 bm	480 bw
	380	365	390	370	375 m	380 m	380 w	373 bs	385 bm
	230	225	242	300	285	260	215 w	235 m	250 bm

تحضير وتوصيف معقدات Cr (III), Mo (V) and W (VI) باستعمال بعض قواعد شف

عبد الهادي رجب حبيب الهيتي عمر حمد شهاب العبيدي

الخلاصة:

يتضمن البحث تحضير معقدات اثنين من قواعد شف (كليكندات) التي لها الاسمين الآتيين:

١- فنيل -٣-مثيل-٥- بايروزولون ثايوسيميكاريزون = (L1) و فنيل -٣-مثيل-٥- بايروزولون -٤- فنيل ثايوسيميكاريزون = (L2) حيث تم تحضير معقدات لهذين الليكندين مع بعض أملاح العناصر الانتقالية التي تشمل (Cr(III), Mo(V) W(VI)) وقد تم تشخيص ودراسة تراكيب الليكندات والمعقدات المحضرة منهما باستخدام تقنية التحليل الدقيق للعناصر (C.H.N) ومطيافية الأشعة تحت الحمراء والأشعة فوق البنفسجية وتقنية الامتصاص الذري إضافة إلى قياس الموصلية الكهربائية المولارية ودرجات الانصهار للمعقدات المحضرة وعلى ضوء النتائج تم التحقق من تراكيب هذه المعقدات بشكل ثماني السطح.