

Role of Polyethylene Glycol in production of Anticancer Alkaloids Vincristine, Vinblastine, and Vindoline in *Catharanthus roseus* via Callus Culture

Zainab Abdulwahid, Abdulminam Hussien Ali and Abdullah Hamad Lafta

Faculty of Science, University of Basrah, Basrah, Republic of Iraq

Corresponding author Email: abdul_minam.ali@uobasrah.edu.iq

DOI: <https://doi.org/10.36077/kjas/2024/v16i3.11847>

Received date: 7 / 4 /2023

Accepted date: 15/ 5 /2023

Abstract

Periwinkle *Catharanthus roseus* is an ornamental plant that produces over 200 alkaloids, including vincristine, vinblastine, and vindoline, which are used to treat various types of cancer. However, the low yield of terpenoid indole alkaloids (TIAs) from periwinkle and the high cost of extraction has led researchers to explore alternative methods for their production. The study aims to investigate the influence of polyethylene glycol (PEG) as an elicitor to enhance alkaloid production in periwinkle callus cultures while examining the efficacy and safety of using PEG. Callus was induced and abiotic elicitation experiments were conducted to study callus growth enhancement. The dry weight of the callus was calculated, and vincristine, vinblastine and vindoline were extracted and analyzed using high-performance liquid chromatography. The results showed that low to moderate concentrations of PEG₄₀₀₀ can enhance callus growth, while higher concentrations can reduce it. The addition of PEG₄₀₀₀ also increased the production of important alkaloids, including vindoline, vincristine, and vinblastine, by up to 2.65-fold. The findings suggest that PEG₄₀₀₀ can be used to optimize the production of alkaloids in *C. roseus* callus cultures.

Keywords: Periwinkle, *Catharanthus roseus*, polyethylene glycol (PEG), vincristine, vinblastine, vindoline



Introduction

Plants produce a wide range of secondary metabolites with diverse applications, including pharmaceuticals, dyes, and insecticides (24). Among these, periwinkle (*Catharanthus roseus*) stands out as an essential source of more than 130 different terpenoid indole alkaloids (TIAs), which have pharmaceutical uses (20, 26). Vincristine and vinblastine are bisindole alkaloids that are widely antineoplastic drugs and cancer chemotherapies (4, 23). Vindoline is also of great significance in pharmacological function. In their research, Eltayeb *et al.*, (6) demonstrated that the leaf extract of *C. roseus* has the ability to inhibit the invasive properties of two types of breast cancer cells by regulating the activity of matrix metalloproteinases. Similarly, Goboza *et al.*, (8) found that vindoline can promote insulin action, leading to a reduction in blood sugar levels in rats. However, the natural yield of these alkaloids in periwinkle is meager, making it necessary to find alternative methods for their production.

Callus culture is an unconventional technology that can be used to produce secondary metabolites and other natural products (18,12). By providing a continuous and reliable source of natural products in the long term, this technology is becoming increasingly popular among researchers. One of the key advantages of callus culture is the ability to use various elicitation factors to trigger and increase the yield of cell secondary metabolites (16, 21). Elicitors are compounds that stimulate the production of secondary metabolites in plants, and among the abiotic elicitors, polyethylene glycol (PEG) has been shown to be effective in enhancing the production of alkaloids in various plants, including

periwinkle (13, 3, 1). However, some studies have raised concerns about the safety and effectiveness of using PEG as an elicitor (17, 10, 5).

The main aim of this study is to investigate the influence of PEG on biomass production and the accumulation of dimeric alkaloids, including vincristine, vinblastine, and vindoline, in callus cultures of periwinkle. Specifically, the study aims to examine the efficacy and safety of using PEG as an elicitor to enhance alkaloid production in periwinkle callus cultures. The hypothesis is that PEG will increase the yield of alkaloids in periwinkle callus cultures without causing any adverse effects on cell growth or viability. The present study will aid in the creation of a productive and long-lasting technique for manufacturing TIAs from periwinkle, which may be utilized for cancer treatment.

Material and Methods

Plant Materials

Catharanthus roseus (L.) Don was selected as the plant species for this study, and pinkish-purple blooms were obtained from a reputable local nursery. The plants used in the experiment were identified and certified by the Department of Biology, College of Science, University of Basrah. Young and healthy leaves were carefully collected from the plants and washed with liquid dish soap and tap water for 15 minutes to remove dust and dirt. To sterilize the leaves, 0.1% mercuric chloride solution containing 2-3 drops of Tween 20 was used for 4-5 minutes. To eliminate any remaining HgCl₂ residue, the explants were washed five times with autoclaved

distilled water. The leaves were then cut into small pieces measuring 0.4-0.5 cm and kept in wet sterile Petri dishes until inoculated into the culture vessels. All sterilization procedures were conducted under aseptic conditions to ensure the purity of the samples.

Callus induction

Sterilized leaf explants (0.4-0.5 cm) were inoculated into 250 ml jars containing 50 ml of MS medium Murashige and Skoog (1962) basal medium supplemented with myo-inositol (100 mg L^{-1}), $\text{NaH}_2\text{PO}_4 \cdot 2\text{H}_2\text{O}$ (150 mg L^{-1}), and 5% sucrose, as well as a combination of phytohormones, including naphthalene acetic acid (NAA) at 5 mg L^{-1} , benzyl aminopurine (BA) at 2 mg L^{-1} , and kinetin (KN) at 2 mg L^{-1} . The pH of the medium was adjusted to 5.8 using 0.1 M NaOH or 0.1 M HCl, followed by the addition of 7 g L^{-1} agar. The medium was then autoclaved at 121°C for 20 minutes. All cultures were incubated in the dark at 27°C in a plant growth room for two months. All chemicals utilized in the tissue culture experiment were ordered from Duchefa Company, Holland.

Callus Treatments

Abiotic elicitation experiments were conducted using the callus culture obtained in Step 1. To study the enhancement of callus growth, vincristine, vinblastine, and vindoline, one gram of callus was inoculated into a medium containing different levels of polyethylene glycol (PEG). Cultures that were not treated with abiotic elicitor (0 g) were used as controls.

Callus Fresh and Dry Weight

After a two-months treatment with an abiotic elicitor, calli were harvested and

carefully rinsed with distilled water to remove any residual medium. The calli were then placed on filter paper to blot away excess water before being weighed to determine their fresh weight using a calibrated balance. To determine the dry weight of the calli, fresh callus samples were placed on filter paper and left to air-dry for 72 hours at 25°C . The dry weight of each callus was then calculated using an electronic balance.

Extraction of Vincristine, Vinblastine, and Vindoline from Callus

To prepare the samples for analysis, 5 g of dried callus was ground into a fine powder and extracted three times with 90% ethanol for 12 hours at room temperature. The resulting alcohol extract was then passed through a Millipore filter and concentrated to 10 mL using a rotary evaporator. For further extraction, a 10 mL water extract was acidified with 3% HCl (10 mL) and then washed three times with 30 mL hexane. The aqueous fraction was then basified to pH 8.5 with ammonia and extracted three times with 30 mL of chloroform. The resulting chloroform extracts were washed with water, dried over sodium sulfate, and vacuum-concentrated to obtain the final sample. Finally, 10 mL of methanol was used to dissolve the sample (9).

Determination of Vincristine, Vinblastine and Vindoline by HPLC

The callus extracts were subjected to high-performance liquid chromatography (HPLC) analysis following the method described by Gupta *et al.* (9). The analysis was performed using an HPLC system (SYKAMN, Germany) equipped with a C18-ODS column and detector. The

mobile phase used was 1.2 mL/min isocratic acetonitrile-0.1 M phosphate buffer containing 0.5% glacial acetic acid (30:70). The system was injected with diluted standards to analyze vincristine, vinblastine, and vindoline. The results were recorded and analyzed using appropriate software.

Data recording and statistical analysis

A series of three *in vitro* experiments were conducted to investigate the response of calli to abiotic stress. Five vessels were used for each treatment. Data on callus induction and proliferation were collected after two months under both stress-free and stressful conditions. The experimental design followed a completely randomized design (C.R.D.) and the Genstats 0.7 program was utilized to determine the least significant difference (LSD) between the means of the treatments. The correlation between stress factor and the production of vincristine, vinblastine, and vindoline was analyzed at a significance level of 0.05 using SPSS 24.

Results and Discussion

Influence of PEG on Callus Biomass and Alkaloid Production

The results of the experiment are presented in Table 1, which shows the callus weight for each PEG treatment. The callus fresh weight was highest in the 0.5 g PEG treatment with a mean of 4.44 ± 0.4 g, while the lowest fresh weight was observed in the 2.0 g PEG treatment with a mean of 4.00 ± 0.5 g. The mean callus fresh weight for all treatments was 4.20 ± 0.6 g. The callus dry weight followed a decreasing trend with increasing PEG levels. The mean dry weight was 0.23 ± 0.08 g, with the lowest dry weight observed in the 2.0 g

PEG treatment (0.22 ± 0.06 g). The LSD values for callus fresh and dry weight were 0.42 and 0.05, respectively. The data suggest that 0.5 g PEG treatment can enhance callus growth, whereas higher levels of PEG can reduce callus growth.

The results of our study are consistent with previous findings reported by Sarmadi *et al.*, (22), who observed a decrease in fresh and dry weights of *Taxus baccata* L. cultures with increasing levels of PEG. The decline in fresh and dry weight levels can be attributed to the reduction in water content, which causes a decrease in cell turgor pressure and oxidative stress levels, resulting in brown calli. However, low to moderate concentrations of PEG was found to reduce the accumulation and oxidation of phenolic compounds in calli, which was attributed to the absorption of phenolic compounds and a reduction in the activity of polyphenol oxidase. Mahmood *et al.*, (14) also confirmed the inhibitory effect of PEG on callus growth and development of wheat, where growth and survival of callus were significantly reduced when transferred to a medium fortified with high levels of PEG. AL-Taha (2) reported similar results, where a reduction in callus growth was observed due to the decrease in cell turgor pressure caused by the reduction in water content. Our HPLC analysis (Table 2, Fig. 1) revealed significant differences in the production of alkaloids in callus culture due to the addition of PEG. The highest amount of vindoline, vincristine, and vinblastine was obtained on a medium containing 2 g PEG (15.90, 29.36, and $41.49 \mu\text{g ml}^{-1}$), followed by 1.5 g PEG (14.79, 28.22, and $40.19 \mu\text{g ml}^{-1}$), respectively. On the other hand, the lowest amount of vindoline,

Table 1 Impact of various different concentrations of PEG₄₀₀₀ on the fresh and dry weight of *Catharanthus roseus* L. callus. *Values followed by the different letters within the same group indicate statistically significant difference.

PEG gm L ⁻¹	SE=	Standard error
	Callus Wight (gm)	
	Fresh Weight+ SE	Dry Weight +SE
Control	4.34±0.5 ^a	0.33±0.04 ^a
0.5	4.44±0.4 ^a	0.29±0.09 ^{ab}
1	4.11±1.09 ^a	0.24±0.1 ^b
1.5	4.09±0.5 ^a	0.23±0.07 ^{ab}
2.0	4.00±0.5 ^a	0.22±0.06 ^{ab}
Mean	4.20± 0.6	0. 23± 0.08
LSD	0.42	0.05

Values followed by the different letters within the same group indicate statistically significant difference

Table 2 Effect of various concentrations of PEG₄₀₀₀ on dimeric alkaloids content of *Catharanthus roseus* callus.

Samples	Vincristine			Vinblastine			Vindoline		
	RT/Min.	Area	Con. µg ml ⁻¹	RT/Min.	Area	Con. µg ml ⁻¹	RT/Min.	Area	Con. µg ml ⁻¹
Callus grew on MS medium w/o PEG ₄₀₀₀	5.96	11253.98	26.90	7.93	15248.99	38.99	4.07	5521.08	13.45
Callus grew on MS medium enriched w/ 0.5 g PEG ₄₀₀₀	5.94	8642.58	27.12	7.96	9652.14	39.20	4.03	3256.58	13.69
Callus grew on MS medium enriched w/ 1 g PEG ₄₀₀₀	5.99	8965.28	27.50	7.93	10240.22	39.69	4.08	3521.49	13.98
Callus grew on MS medium enriched w/ 1.5 g PEG ₄₀₀₀	5.96	9242.58	28.22	7.92	11253.08	40.19	4.05	3652.18	14.79
Callus grew on MS medium enriched w/ 2 g PEG ₄₀₀₀	5.90	9521.28	29.36	7.92	12465.80	41.49	4.03	3854.08	15.90

w= with w/o= without, RT=Retention time, Con.= concentrations




Figure 1 The predictive curve for analyzing alkaloids from callus treated with different concentrations of PEG shows an increase in alkaloid production with an increase in the concentration of PEG's osmotic effect. w= with w/o= without

vincristine and vinblastine were recorded in callus grown on a medium free of PEG and a medium containing 0.5 g PEG. Interestingly, the production of these alkaloids increased 2.65, 2.61, and 2.5-fold, respectively, with the addition of PEG. These findings suggest that PEG can be used to enhance the production of important alkaloids in *C. roseus* callus cultures.

PEG is one of the most commonly used materials in water stress experiments and for creating drying conditions similar to the natural environment. Where, it creates a solution with a lower water potential than the plant cells, which causes water to move out of the cells by osmosis. However, PEG is widely used in various fields, including biochemical and industrial applications, due to its non-toxicity (15). Polyethylene glycol (PEG) has also been used in *in vitro* plant culture to simulate water stress conditions, which can result in increased

production of secondary metabolites (1). Taha *et al.* (24) found that adding a combination of PEG and chitosan to the growth medium of *C. roseus* increased the production of several alkaloids, including vincristine and vinblastine. Amirjani, (3) confirmed the stimulatory effect of PEG on the production of vincristine and vinblastine in *C. roseus* treated with 12% PEG for 72. Iskandar and Iriawati (10), demonstrated that exposing plants to drought using PEG₄₀₀₀, up to a concentration of 12% (w/v), did not have a significant impact on the production of vinblastine. While high vincristine concentration was obtained on a medium containing 0% of PEG. According to the study findings, there is a distinct pattern in the concentration of alkaloids that appears to be linked to the level of drought stress. This is likely due to the fact that the administered PEG concentration was not sufficiently high to reduce alkaloid

biosynthesis while staying within the cells' Plant growth, morphology, and metabolic processes are all impacted by drought and salt stresses. To cope with these stresses, plants undergo adaptations that involve altering their metabolic processes. This includes producing and storing primary and specialized metabolites that aid in drought and salt resistance (27). Arabidopsis plants respond to drought stress by accumulating glucosinolates (11). In *C. roseus* plants, both drought and salt stress lead to an increase in the accumulation of terpenoid indole alkaloids (TIAs), such as ajmalicine, catharanthine, vindoline, vinblastine, and vincristine (12).

Theoretically, an increase in the concentration of PEG administered to the culture would result in a corresponding increase in the level of drought stress experienced by the plant cells. This, in turn, would trigger the plant cells to produce higher levels of abscisic acid as an initial response to drought stress (7). Absciscic acid (ABA) is a critical hormone in plant responses to drought stress. It regulates various physiological processes such as stomatal closure, which helps the plant to conserve water. The production of alkaloids is also regulated by ABA, and previous studies have shown that an increase in ABA levels can lead to promoting catharanthine production in *C. roseus* suspension cells (28).

Therefore, it is plausible to assume that the increase in abscisic acid levels triggered by PEG-induced drought stress may have contributed to the observed increase in alkaloid production in the present study. However, further research is needed to confirm this hypothesis and to elucidate the underlying mechanisms involved.

It should also be noted that the effects of PEG-induced drought stress on alkaloid

tolerance limits.

production may vary depending on the plant species and the type of alkaloids produced. The correlation analysis provides information on the stress factor of polyethylene glycol (PEG) and the TIAs compounds' production. The correlation between the compound vincristine and other alkaloids compounds was a strong inverse correlation (+), with R value of (0.975), followed by the compound vindoline with a value of (0.971), and then the vinblastine with a value of (0.966) Plate 1. The significant increase in the quantity of secondary compounds under the influence of the osmotic stress factor is attributed to the fact that growing cells under stress conditions can produce more polyamines, which are key components in the production of many secondary metabolites (25). Additionally, PEG increases the negative osmotic pressure of the cell, thereby increasing the exchange of solutes in the cell and consequently the production of secondary compounds. Therefore, caution should be exercised when extrapolating the findings of this study to other plant species or alkaloids.


Conclusion

Results of the present study suggest that PEG₄₀₀₀ in low concentrations with co-stress factor (sucrose 5%) induced drought stress can enhance alkaloid production in plants, possibly by increasing abscisic acid levels. This finding has potential implications for the pharmaceutical industry, as alkaloids are an important source of drugs and drug precursors. Further research is needed to explore the potential of using drought stress as a tool to enhance alkaloid production in plants.

Conflict of interest

The authors have no conflict of interest.





References

- Ahmad, M.A., Javed, R., Adeel, M., Rizwan, M., Yang, Y. 2020. PEG 6000-Stimulated drought stress improves the attributes of *In Vitro* growth, steviol glycosides production, and antioxidant activities in *Stevia rebaudiana* Bertoni. Plants, (Basel). 9(11): 1552.<http://doi.org/10.3390/plants9111552>. PMID:33198205 ; PMCID: PMC7696522.
- AL-Taha, H.A.K. 2013. Effect of shock and gradual drought by PEG on callus growth and proline accumulation in sour orange (*Citrus aurantium*). Advances in Agriculture and Botany, 5(2): 77-83.<http://www.aab.bioflux.com.ro/docs/2013.77-83.pdf>
- Amirjani, MR. 2013. Effects of drought stress on the alkaloid contents and growth parameters of *Catharanthus roseus*. Journal of Agriculture and Biological Sciences, 8(11):745-750. http://www.arpnjournals.com/jabs/research_papers/rp_2013/jabs1113_617.pdf
- Birat, K., Binsuwaidan, R., Siddiqi, T. O., Mir, S. R., Alshammari, N.,

- Adnan, M., Nazir, R., Ejaz, B., Malik, M.Q., Dewangan, R.P., Ashraf, S.A. and Panda, B.P. 2022. Report on Vincristine-Producing endophytic fungus *Nigrospora zimmermanii* from leaves of *Catharanthus roseus*. *Metabolites*, 12(11): 1-13.<http://dx.doi.org/10.3390/metabo1211119>
5. Elham A., Ziba G. H., Amin E., Hojatollah B. 2022. Expression of CrMPK3 and alkaloid synthesis genes with antioxidants in callus of *Catharanthus roseus* in response to polyethylene glycol. *Industrial Crops and Products*, 178: 114634 <http://doi.org/10.1016/j.indc.rop.2022.114634>
 6. Eltayeb, N. M., Ng, S. Y., Ismail, Z., and Salhimi, S. M. 2016. Anti-invasive effect of *Catharanthus roseus* extract on highly metastatic human cancer MDA-MB-231 cells. *Jurnal Teknologi*, 78(11-3): 35-40 <https://doi.org/10.11113/jt.v78.9870>
 7. Farooq, M., Hussain, M., Wahid, A. and Siddique, K.H.M. 2012. Drought Stress in Plants: An Overview. In: *Plant Responses to Drought Stress From Morphological to Molecular Features* ed. Aroca R. pp. 1-33. Springer Verlag. Berlin. Heidelberg, http://doi.org/10.1007/978-3-642-32653-0_1.
 8. Goboza, M., Aboua, Y. G., Chegou, N., and Oguntibeju, O. O. 2019. Vindoline effectively ameliorated diabetes-induced hepatotoxicity by docking oxidative stress, inflammation and hypertriglyceridemia in type 2 diabetes-induced male Wistar rats. *Biomedicine and pharmacotherapy*, 112:1-11.<https://doi.org/10.1016/j.biopha.2019.108638>
 9. Gupta, M. M., Singh, D. V., Tripathi, A. K., Pandey, R., Verma, R. K., Singh, S., Shasany, A. K., and Khanuja, S. P. 2005. Simultaneous determination of vincristine, vinblastine, catharanthine, and vindoline in leaves of *Catharanthus roseus* by high-performance liquid chromatography. *Journal of Chromatographic Science*, 43(9): 450-453.<https://doi.org/10.1093/chromsci/43.9.450>
 10. Iskandar, N. N and Iriawati, 2016. Vinblastine and vincristine production on Madagascar periwinkle (*Catharanthus roseus* (L.) G. Don) callus culture treated with polyethylene glycol. *Makara Journal of Science*, 20(1): 7-16. <https://doi.org/10.7454/mss.v20i1.5656>
 11. Li, B., Fan, R., Guo, S., Wang, P., Zhu, X., Fan, Y., Chen, Y., He, K., Kumar, A. and Shi, J. 2019. The Arabidopsis MYB transcription factor, MYB111 modulates salt responses by regulating flavonoid biosynthesis. *Environmental and Experimental Botany*, 166: 103807 <https://doi.org/10.1016/j.envexpbot.2019.103807>.
 12. Liu, Y., Meng, Q., Duan, X., Zhang, Z. and Li, D. 2017. Effects of PEG induced drought stress on regulation of indole alkaloid biosynthesis in *Catharanthus roseus*. *Journal of Plant Interactions*, 12(1):87-91 : <https://doi.org/10.1080/17429145.2017.1293852>
 13. Lucas, D.M., Still, P.C., Pérez, L.B., Grever, M.R. and Kinghorn, A.D. 2010. Potential of plant-derived natural products in the treatment of leukemia and lymphoma. *Current Drug Targets*, (7): 812-822. <https://doi.org/10.2174/138945010791320809>
 14. Mahmood, I., Razzaq, A., Hafiz, I.A., Kaleem, S., Khan, A.A. and Ahmad, M. 2012. Interaction of callus selection media and stress duration



- for *in vitro* selection of drought tolerant callus of wheat. African Journal of Biotechnology, 11(17): 4000-4006. <http://doi.org/10.1080/14786419.2019.1689500>
15. Meher, Shivakrishna P, Ashok, Reddy K and Manohar, Rao D. 2018. Effect of PEG-6000 imposed drought stress on RNA content, relative water content (RWC), and chlorophyll content in peanut leaves and roots. Saudi Journal of Biological Sciences, 25(2):285-289. <http://doi.org/10.1016/j.sjbs.2017.04.008>
 16. Miclea, I., Suhani, A., Zahan, M. and Bunea, A. 2020. Effect of jasmonic acid and salicylic acid on growth and biochemical composition of in-vitro-propagated *Lavandula angustifolia* Mill. Agronomy, 10: 1722-1736. <https://doi.org/10.3390/agronomy10111722>
 17. Morgan, J. A., and Shanks, J. V. 2000. Enhancement of indole alkaloid production by *Catharanthus roseus* cell cultures using the alkaloid precursor, tryptophan. Biotechnology and Bioengineering, 67(2): 246-251.
 18. Mujib, A., Ilah, A., Aslam, J., Fatima, S., Siddiqui, Z.H., and Maqsood, M. 2012. *Catharanthus roseus* alkaloids: Application of biotechnology for improving yield. Plant Growth Regulation, 68: 111–127. <https://doi.org/10.1007/s10725-012-9704-4>
 19. Murashige, T. and Skoog, F. 1962. A revised medium for rapid growth and bioassays with tobacco tissue culture. Physiologia Plantarum, 15:473-497. <https://doi.org/10.1111/j.1399-3054.1962.tb08052.x>
 20. Nejat, N., Valdiani, A., Cahill, D., Tan, Y.H., Maziah, M. and Abiri, R. 2015. Ornamental exterior versus therapeutic interior of Madagascar periwinkle (*Catharanthus roseus*): the two faces of a versatile herb. The Scientific World Journal, 2015: 1-19. <https://doi.org/10.1155/2015/982412>
 21. Sák, M., Dokupilová, I., Kaňuková Š., Mrkvová, M., Mihálik, D., Hauptvogel, P. and Kraic, J. 2021. Biotic and abiotic elicitors of stilbenes production in *Vitis vinifera* L. cell culture. Plants, 10: 1-14. <https://doi.org/10.3390/plants10030490>
 22. Sarmadi, M., Karimi, N., Palazón, J., Ghassempour, A. and Mirjalili, M. H. 2019. Improved effects of polyethylene glycol on the growth, antioxidative enzymes activity and taxanes production in a *Taxus baccata* L. callus culture. Plant Cell, Tissue and Organ Culture, 137: 319–328. <https://doi.org/10.1007/s11240-019-01573-y>
 23. Tang, W., Liu, X., He, Y. and Yang, F. 2022. Enhancement of vindoline and catharanthine accumulation, antioxidant enzymes activities, and gene expression levels in *Catharanthus roseus* leaves by Chitoooligosaccharides elicitation. (2022). Marine Drugs, 20: 188-201 <https://doi.org/10.3390/md20030188>
 24. Taha, H.S., El-Bahr, M.K. and Seif-El-Nasr, M.M. 2009. *In vitro* studies on Egyptian *Catharanthus roseus* (L.). Effect of biotic and abiotic stress on indole alkaloids production. Journal of Applied Sciences Research, 5: 1826-1831. <http://www.ansiweb.com/old/jasr/jasr/2009/1826-1831.pdf>
 25. Tun, N. N., Santa-Catarina, C., Begum, T., Silveira, V., Handro, W., Floh, E. I., and Scherer, G. F. 2006. Polyamines induce rapid biosynthesis of nitric oxide (NO) in Arabidopsis thaliana seedlings. Plant Cell Physiology, 47(3): 346–354. <https://doi.org/10.1093/pcp/pci252>
 26. Vu, P. T. B., Cao, D. M., Bui, A. L., Nguyen, N. N., Bui, L. V. and Quach, P. N. D. 2022. *In vitro* growth and



- content of vincristine and vinblastine of *Catharanthus roseus* L. hairy roots in response to precursors and elicitors. *Plant Science Today*, 9(1): 21–28. <https://doi.org/10.14719/pst.1337>
27. Zahedi, S.M., Karimi, M. and Venditti, A. 2019. Plants adapted to arid areas: specialized metabolites. *Natural Product Research*, 35(9): 3314–3331. <https://doi.org/10.1080/14786419.2019.1689500>
28. Zhou, P., Yang, J., Zhu, J., He, S., Zhang, W., Yu, R., Zi, J., Song, L. and Huang, X. 2015. Effects of β -cyclodextrin and methyl jasmonate on the production of vindoline, catharanthine, and ajmalicine in *Catharanthus roseus* cambial meristematic cell cultures. *Applied Microbiology Biotechnology*, 99: 7035–7045. <http://doi.org/10.1007/s00253-015-6651-9>

