

Sahar Abd Department of Biology, College
of Science for Women,
University of Baghdad**Soolaf A. Kathier**Department of Biology, College
of Science for Women,
University of Baghdad.**Emad Ahmed Mahmmod**Department of Biology, College
of Science for Women,
University of Baghdad.

Received on: 10/11/2018

Accepted on: 07/04/2019

Published online: 25/04/2019

Effect of Gamma Radiation on *Callosobruchus maculatus* (F.) (Coleoptera: Bruchidae) and Seed Germination

Abstract: The grain legumes are the second most important economic crops attacked by many of storage pests, one of these pests is Cowpea seed beetle (*Callosobruchus maculatus*). Gamma radiation appears to be a potential alternative to chemicals control in stored products. Therefore, this research aims to study the effect of gamma radiation (0.03, 0.06, 0.09, 0.12, 0.15, 0.18 and 0.21 (kGy) on eggs, larvae, pupae and seed germination. The results of this study showed that the irradiated eggs did not hatch. While the last larval stage was very sensitive to all doses of gamma rays, no adults emerged from larvae were exposed to 0.03 (kGy) or higher doses. In addition, the highest death percentages of pupae were 88% at 0.21(kGy). Finally, the results showed the same doses used in this study did not affect the rate of seed germination compared to control. This technique would help to improve using for other agriculture pests.

Keywords: Gamma Radiation, *Callosobruchus maculatus*, larval stage, Coleoptera

How to cite this article: S. Abd, S.A. Kathier and E.A. Mahmmod, "Effect of Gamma Radiation on *Callosobruchus maculatus* (F.) (Coleoptera: Bruchidae) and Seed Germination," *Engineering and Technology Journal*, Vol. 37, Part C, No. 1, pp. 90-92, 2019.

1. Introduction

Grain legumes are the second most important group of crops worldwide. Globally, 840 million people are undernourished mainly on account of inadequate intake of proteins, vitamins and minerals in their diets, pulses are excellent sources of proteins (20-40%), carbohydrates (50-60%) and are fairly good sources of thiamin, niacin, calcium and iron [1]. One of the major constraints in production is insect pests, which inflict severe losses in both the field and storage. Cowpea seed beetle (*Callosobruchus maculatus*) is a major pest that causes serious damage [2,3]. There are various methods for control of the cowpea weevils and other storage pests including, insecticides which are widely used to protect stored commodities from insect infestations and contamination but their use lead to problem of undesirable [4,5]. Ionizing radiation used successfully to control insect species and may prove effective in controlling many others. Gamma radiation appears to be a potential alternative to chemicals for insect control in stored products, One of the modern techniques in controlling stored food pests is the use of nuclear technology [6,7,8]. Therefore, this work aims to study the effect of gamma radiation on these pests at some stages and the effect of radiation on seed germination.

2. Materials and Methods

I. Stock culture

All experimental procedures were carried out at Entomology Laboratory, University of Baghdad and under the same environmental conditions. Beans were collected from various stores, and were brought to the Laboratory; grains were kept in jars and covered with muslin cloth at $29 \pm 1^{\circ}\text{C}$ and relative humidity 60 ± 5 . The culture was observed regularly for the presence of *Callosobruchus maculatus* [9].

II. Irradiation of egg, larval, and pupal stages:

The beans had larvae of same age were isolated and collected from the culture. The one day old eggs, nine day old larvae, and four day old pupae were exposed to 0.03, 0.06, 0.09, 0.12, 0.15, 0.18 and 0.21 (kGy). 220-GammaCell, a radiation source of Co^{60} . The development of all stages were recorded after gamma ray treatments. The experiment was designed in Complete Randomized Design with three replications and twenty-five for each.

III. Seed germination

50 seeds of beans in three replications were exposed to gamma rays (cobalt-60 source) at doses 0.03, 0.06, 0.09, 0.12, 0.15, 0.18 and 0.21 (kGy). Also the healthy seeds were exposed to these doses and subjected to germination test.

IV. Statistical analysis

Experimental results were analyzed by Complete Randomized Design. Duncan- test was used to separate means when significant overall differences occurred at 5%.

3. Results and Discussion

Table 1 shows that eggs did not hatch at all doses, which means that gamma radiation has influence on the embryo development. Eggs of *Callosobruchus maculatus* are very sensitive and killed when exposed to gamma radiation, especially when the one - day old eggs [10]. Table 2 shows that larval stage has different radiosensitive to gamma radiation. The dose 0.15 (kGy) gave 100 killing, because the larvae stop feeding due to the breakdown of lining of the digestive system [11]. The percentage of normal pupae from irradiated larvae decreased from 93% in control to 3% at treatment with 0.12 (kGy), the percentage of abnormal pupa increased from zero in control group to 7% at dose 0.21 (kGy), and showed reduction percentage of normal adults from 87% in control to 0% at dose (0.03kGy). These results are consistent with the results of Al-Badry and Ahmed [12] where it was found that a

dose of 3 (kGy) lead to the death of all the larvae of the *Callosobruchus maculatus* before they reach to adults. Treated *Callosobruchus maculatus* with dose (20 kGy) induces semi-sterility [13]. Table 3 shows that the mortality of pupa was increased significantly with increasing dose radiation. However, the pupa stage was less sensitive than larval stage. Significant differences were found among tested doses compare to control in percent pupal mortality [14].

Table 1. Effect of radiation doses on egg hatching of *Callosobruchus maculatus*

Radiation Dose (kGy)	Egg hatching %
0	83.5 a
0.03	0 b
0.06	0 b
0.09	0 b
0.12	0 b
0.15	0 b
0.18	0 b
0.21	0 b

Similar letters within the columns indicate no significant differences between the different doses at 5%

Table 2. Effect of radiation doses on the mortality rate of the last larval stage of *Callosobruchus maculatus*

Radiation Dose (kGy)	Mortality larva %	Abnormal pupa %	Normal pupa %	Abnormal adults %	Normal adults %
0	70 d	0 c	93 a	0 b	87a
0.03	78 c	40 b	18 b	7 a	0 b
0.06	84 bc	10 a	6 c	0 b	0 b
0.09	89 b	9 a	2 c	0 b	0 b
0.12	90 b	7 a	3 c	0 b	0 b
0.15	100 a	-	-	-	-
0.18	100 a	-	-	-	-
0.21	100 a	-	-	-	-

Similar letters within the columns indicate no significant differences between the different doses at 5%.

Table 3. Effect of radiation doses on the mortality rate of the pupal stage of *Callosobruchus maculatus*

Radiation Dose (kGy)	Mortality pupa %	Abnormal adults %	Normal adults %
0	17.4 e	0 c	82.6 a
0.03	22.6 de	0 c	77.3 ab
0.06	27.7 d	4 b	68.3 b
0.09	56 c	4 b	40 c
0.12	49.3 c	10.7 a	40 c
0.15	72 b	8 a	20 d
0.18	80 ab	6.7 ab	13.3 d
0.21	88 a	8 a	4 e

Similar letters within the columns indicate no significant differences between the different doses at 5%

Table 4 shows no significant differences in seed germination rate after irradiated with a different doses of gamma rays compared to control. However, only the seeds of Barley (*Hordeum vulgare*) treated with 0.4 kGy had a significant lower germination rate than the other three treatments (control, 0,1 and 0.2 kGy).

Table 4. Effect of radiation doses in the germination of beans seeds

Radiation dose	Germination rate
0	63 +5.05 a
0.03	70.5+ 5.09 a

0.06	56+ 3.26	a
0.09	62+ 6.93	a
0.12	62+ 5.16	a
0.15	56+ 3.27	a
0.18	60+ 4.60	a
0.21	62+ 5.16	a

Similar letters within the columns indicate no significant differences between the different doses at 5%

4. Conclusions

Eggs of *Callosobruchus maculatus* are very sensitive when exposed to gamma radiation, while the larval stage has different radiosensitive and not effect in seed germination rate after irradiated with a different doses of gamma rays .

References

- [1] CAB International. Crop Protection Compendium. Wallingford, UK: CAB International. 2007.
- [2] The Arab Organization for Agricultural Development. Study of the possibilities of using irradiation technology in the conservation and storage of food products in the Arab world, 281 p. 1995.
- [3] C. Fair, T. Harness, O. Mall. Reducing Loss of Crops after Harvest, by Dr. Fares Amin Mohamed, Zagazig University, 284 p. 1994.
- [4] R .Champ, F. Dyte. FAO global survey of pesticides susceptibility of stored grain pests. FAO plant protection bull ; 25(2);67-82. 1977.
- [5] E. Rechcigl , A. Rechcigl . Insect pest management techniques for environmental production. Lewis Publishers , Boca Roton, New York, Washington, D.C., 392. 2000
- [6] A. El-Kady, M.Hekal. Irradiation disinfestations of pulse and resistance of packing film to insect penetration. Proc. Of the final research coordination meeting on insect disinfestations of food and agricultural products by irradiation , Beijing , China , 25-29, IAEA, 59-68. 1991
- [7] IAEA. Insect disinfestations of food and agricultural products by irradiation, Beijing , China , 25-29, IAEA, Vienna .174pp. 1991.
- [8] E. Kovacs, "Irradiation disinfestations of wheat, dried wheat products by irradiation," Proc. of the final research coordination meeting on insect disinfestations of food and agricultural products by irradiation, Beijing, China, 25-29, IAEA, Vienna pp.69-88, 1991.
- [9] S. Fatima, A. Usman, M.Afzaal. Rearing and identification of *Callosobruchus maculatus* (Bruchidae: Coleoptera) in Chickpea. *Journal of Entomology and Zoology Studies*, Vol. 4, No. 2, pp. 264-266, 2016.
- [10] H. Belal, "Effect of gamma radiation on Eggs of callosobruchus maculatus (fabricius), (Coleoptera: bruchidae)," Damascus University *Journal for the Agricultural Sciences*, Vol. 22, No. 1, pp. 131-146, 2006.
- [11] M. Diop, D.Ba, C.Hasselmann. Radiation disinfestations of cowpea seeds contaminated by *Callosobruchus maculatus*. *Jfood process. Preserv Mar*, Vol..21, No. 1, PP. 69-81, 1997.
- [12] A. Al-Badry , Y.Ahmed.effect of gamma radiation on the egg stage of southern cowpea weevil *Callosobruchus maculatus* *Zeitschrift fur Angewandte Entomology*, Vol. 79, No. 3, pp. 323-328, 1975.
- [13] A. Ibrahim, S. Fawki , M. Abd El-Baret. Inherited influence of low dose gamma radiation on the reproductive potential and spermiogenesis of the cowpea weevil, *Callosobruchus maculatus* (F) (Coleoptera: Chrysomelidae). *Journal of Radiation Research and Applied Sciences*, Vol. 10 pp. 338-347, 2017.
- [14] J .Supawan , P.Hormchan, M.Sutantawong. Effects of Gramma Radiation on Azuki Bean Weevil, *Callosobruchus chinensis* (L.). *Kasetsart J. (Nat. Sci.)* Vol.39, pp. 206 – 215, 2005.
- [15] L. Rozman. The effect of gamma radiation on seed germination of barley (*Hordeum vulgare* L.). *Acta agriculturae Slovenica*, Vol.103, No. 2, pp. 307 – 311, 2014.