



## Study of photolysis on the active material Phenylbutazone in veterinarian drug Isophen

Hanan H. Flaeh      Wajeeh Y. Mohammed      Authman S. Ibrahim      Sattar S. Ibrahim

University of Anbar - College of Science.

### ARTICLE INFO

Received: 5 / 1 /2012  
Accepted: 9 / 5 /2012  
Available online: 29/8/2013  
DOI: 10.37652/juaps.2012.80690

#### Keywords:

Isophen,  
Phenylbutazone,  
photolysis.

### ABSTRACT

The veterinary medicine isophen was used in this study. Isophen contains phenylbutazone as active material. Numbers of samples of isophen were prepared and were determined absorbance were determined and comparison subjected to radiation for different periods (1, 2, 3, 4, 5) hrs. the Maximum absorbance were determined and compression Maximum absorbance's Study are performed. And ensure that the impact of irradiation on the active ingredient in the medication. The results reveal that has been reached that there is significant impact of irradiation as well as the time of irradiation on the decomposition of the active ingredient in the medication was found that with the continuation of the time of irradiation increases decomposition.

### Introduction:

Phenylbutazone is an effective non-steroidal anti-inflammatory drug (NSAID) with antipyretic and analgesic activity, used in veterinary medicine for more than 50 years to treat bone and joint inflammations laminitis and inflammation of soft tissues [1].

It is widely used in dogs and horses but because of toxicity and the lack of established maximum residue limit, is not approved for use in food producing animals. The most serious adverse reaction of phenylbutazone observed in human and animals are: gastric and intestinal ulceration and bleeding, disturbances in platelet function.

The prolongation of gestation or spontaneous labour and changes in renal function. In Finland phenylbutazone has been one of the most widely used non-steroidal anti-inflammatory drug after acetylsalicylic acid. There is still no maximum residue level for phenylbutazone at the 1998. [2]

phenylbutazone effects by preventing the synthesis of prostaglandins. As a medicine phenylbutazone has antipyretic, analgesic and anti-inflammatory effects oxyphenbutazone a metabolic of phenylbutazone has one fifth of the medical activity of phenylbutazone [1, 3].

### Mechanism of action.

Inhibition of the arachidonic acid cascade at the level of prostaglandin H synthetases and prostaglandin synthetases results in decrease production of prostaglandins and thromboxane. also Inhibits urate crystal phagocytosis by synoviocytosis [4].

### Phenylbutazone

**Distribution:** phenylbutazone is distributed mainly in to plasma and extra cellular fluid, as indicated by the relatively small volume of distribution, this is low volume of distribution is also indicative of only minimal tissue binding [5].

**Pharmacokinetics:** phenylbutazone is absorbed from both the stomach and small intestine. the drug is distributed throughout the body with highest levels attained in the liver, heart, lungs, kidneys, and blood. Both phenylbutazone and oxyphenbutazone cross the placenta and are excreted into milk [5].

### Adverse effect / warnings.

The primary concern with phenylbutazone therapy in humans include its bone marrow effects (agranulocytosis, aplastic anemia) renal and cardiovascular effects (fluid retention to acute renal failure)[5].

**Dosage and administration:** phenylbutazone may be administered orally (via paste, powder, or feed-in) or intravenously. It should not be given intramuscularly or injected in any place other than a vein, as it can cause tissue damage.

Side effect and disadvantages of phenylbutazone :

\* Corresponding author at: University of Anbar - College of Science. E-mail address:

Side effect of phenylbutazon are similar to that of the Non-steroidal anti-inflammatory drugs overdos or prolonged use Can cause gastrointestinal ulcers, blood dyscrasia ,kidney damage.Phenylbutazon Is abtained in straight forward manner by Condensation of diethyl-n-butylmalonate with hydrazobenzen in the presence of base.In effect ,this represent the formation of heterocyclic system by simple Lactamization [6] phenylbutazon should be used cautiously in pregnant or nursing mares ,as it may be toxic to the embryo and can be transferred via the umbilical cord and milk.

High dose of phenylbutazon may be considered arules violation under some equestrian organization as the drug may remain in the blood stream four to five days after administration . In human phenylbutazon is very dangerous as it can cause aplastic anemia. The medicine should be give in apast from to avoid contact with the medicin .Never breathe powder from crushing tablets.

**Methods:**

The drug used in this investigation was Isophen and the active material in this medicine is Phenylebutazone which is applied as Injected solution of 200ml samples was prepared according to the following steps 1-5ml of drug was dissolved in ethanol as solvent:

- 2- No. of samples were prepared and subjected to radiation for different Periods (1,2,3,4,5) hours.
- 3-Absorbances was measured using spectrophotometer and  $\lambda_{max}$  was termine.
- 4- differences between the values of  $\lambda_{max}$  for all samples was recorded .

**Result and Discussion:**


Many authors [ 3,6,7] observed degradation of Penyle butazone During analysis , if samples were exposed to acidic condition ,left Dry and open to the atmosphere ,or when containing oxygen ,diethyl ether was used as elution solvent in solid – phase extraction . our experiences indicate that addition of ascorbic acid solution as stabilizer to the extract . Aroom temperature, photochemical spectrophotometric method has been developed for the assay of Phenyle butazone and its degradation products are reported as well as irradiation times (1-5) hrs. which correspond to maximum ultra violet signals of photo products. Our study investigate that the results of samples radiation for different Periods (1,2,3,4,5) hours showed a

lowering in  $\lambda_{max}$  values with Increasing of time radiation also absorbance increased , as shown In table (1) and figures. (1-6).

**Table ( 1 ):  $\lambda_{max}$  ( nm ) for isophen at different time of radiation .**

No.	Radiation time (hrs .)	$\lambda_{max}$ ( nm )
1	Pure	280
2	1	278
3	2	277
4	3	275
5	4	273
6	5	272

Studies of oxygenation of phenyle butazone have established that phenyle butazone is true reducing cofactor for peroxidase activity of prostaglandin H synthase [ 6,8,9] oxidized Phenyle butazone incorporates molecular oxygen to yield 4-hydroperoxy - Phenyle butazone which then reduced to 4- hydroxy- Phenyle butazone [10] , these observation have been used to support the mechanism [ 8 ] . shown in scheme ( 1 ) , table (2).



schem (1) Mechanism of photolysis for phenylbutazone

No	R	Products
1	R=NHPh	2-butyl-N1,N3-diphenyl-2-(phenylamino)malonamide
2	R=OH	2-butyl-2-hydroxy-N1,N3-diphenylmalonamide
3	R= N(CH <sub>2</sub> CH <sub>3</sub> ) <sub>2</sub>	2-butyl-2-(diethylamino)-N1,N3-diphenylmalonamide
4	R= OCH <sub>3</sub>	2-butyl-2-methoxy-N1,N3-diphenylmalonamide

**Table (2): products of mechanism of photolysis for phenylbutazone**

The malonamide in no. 1 when ( R= NHph ) and No.2 when ( R= OH ) And the 2-oxohexanamide in No.5 . when the solutions was basified With diethyl amine . The amino diamid in no. 3 when ( R= N(Et)2)

was Produced in addition to No.2 when ( R= OH ) and No. 2 . In methanol Solution the malonamide in No. 1 when ( R= NHph ) and No. 4 when ( R= OMe ) where obtained .

### References:

- 1-Less, P., Ayliffe, T. Maitho, T.E., 1988. Pharmacokinetics, metabolism and excretion of phenylbutazon in cattle following intravenous, intramuscular and oral administration .Research in veterinary science in veterinary science :44:57-67 .
- 2- Gonzalez Martin M.I., Sanchez Gonzalez C.I., Jimenez Hernandez A., Garcia Cachan M.D., Castro de Cabo M.J., Garzon Cuadrado A.L.,2002, Determination by highperformance liquid chromatography of phenylbutazone in samples of plasma from fighting bulls. J Chromatography, 769, 119-126.
- 3-Less, P. and May, S.A. 1992, chapter 56 : inflammation and anti-inflammatory drugs P: 843-851,in bovine medicine :diseases and husbandry of Cattle. Ed.A.H.Andrews . Black well Scientific publications. London.
- 4-Katzung BG, 1995 editor . Basic and clinical Pharmacology 6th . Norwalk Connecticut : Apleton and Lang.
- 5-Williams RJ, Smith JA ,Boudinot F,1990, pharmacokinetics of phenyl butazon in mature Holstein bulls: steady-state kinetics After multiple oral and dosing , Am J.Vet.: 51 (3) : 367- 370 .
- 6-Grippa E. Santini L ., Castellano G., Gatto M. T. , Leon M.G., Saso L.,2000 Simultaneous ,determination of hydro cortisone dexamethason indomethacin; phenylbutazon and oxyphenbutazon in equine serum by high performance liquid chromatography. J. Chromatography B (2000) ,738 ,pp 17 – 25 .
- 7- Gowik P., Jülicher B., Uchlig S.,1998, Multi-residue method for non-steroidal anti-inflammatory drugs in plasma using high performance liquid chromatography–photodiode array detection. Method description and comprehensive in-house validation. J.Chromatography B,(1998)716,221-232.
- 8-Marnet,L.J., Dix.T.A., Sachs, R.J.,and Siedlik , P. H., (1983), in advances in prostaglandin. Thromboxane, and Leukotriene. Research (Samuelsson ,B paoletti,R and, Ramwell ,p )Vol. 11, pp: 79-86, Raven press , New York

- 9- Siedlik.P.H. and Marnet, L.J.,(1984), Methods Enzymol, in press
- 10-Marnet,L.J., Siedlik, P. H.,and Fung,L.W.M., (1982 ), J. Biol.chem., 257, pp: 6975-6964.
- 11-Marnet,L.J., Dix.T.A., Sachs, R.J.,and Siedlik , P. H., (1983), advances prostaglandins. Thromboxane, and Leukotriene. Vol. 6, pp: 149-151, Raven press , New York.




Fig.(1): Absorbance of drug before radiation( pure )




Fig.(2): Absorbance of drug after one hour of radiation




Fig.(3): Absorbance of drug after two hour of radiation




Fig.(4): Absorbance of drug after three hour of radiation




Fig.(5): Absorbance of drug after four hour of radiation




Fig.(6): Absorbance of drug after five hour of radiation

## دراسة التحلل الضوئي للمادة الفعالة Phenylbutazone في الدواء البيطري Isophen

حنان حسن فليح      وجيه يونس محمد      عثمان سالم إبراهيم      ستار سالم إبراهيم

### الخلاصة:

أستخدم في هذا البحث الدواء البيطري آيزوفين Isophen الحاوي على المادة الفعالة فنيل بيوتازون Phenylbutazone إذ تم تحضير عدد من النماذج للدواء البيطري المذكور ثم عرضت النماذج المحضرة للتشعيع ولفترات زمنية مختلفة (1, 2, 3, 4, 5) ساعة ثم حسبت الامتصاصية العظمى ( $\lambda_{max}$ ) لكل نموذج بعد التشعيع لغرض دراسة المقارنة بين القيم العظمى للامتصاصية لجميع النماذج المحضرة من الدواء آيزوفين والتأكد من تأثير فترة التشعيع على المادة الفعالة في الدواء . حيث بينت النتائج التي تم التوصل إليها انه يوجد تأثير كبير للتشعيع وكذلك الفترة الزمنية للتشعيع على تحلل المادة الفعالة في الدواء إذ وجد انه مع استمرار فترة التشعيع تزداد نسبة التحلل.