

A STUDY OF *AEGYPTIANELLA SPP* IN SOME SPECIES OF BIRDS IN MOSUL CITY-IRAQ

E.G.Suleiman

Department of Microbiology, College of Veterinary Medicine, University of Mosul,
Mosul, Iraq

(Received 3 July 2012, Accepted 23 July 2012)

Keywords; *Aegyptianella spp*, *Anaplasma*, turkeys.

ABSTRACT

During this study examination of 205 blood samples collected from four species of birds , chickens(50), ducks(75), turkeys(30), and pigeon(50) in different regions of Mosul city for the detection of *Aegyptianella spp*. The total percentage of infection was 28.29% and infection was (33.33%), (30%), (26.66%), and (20%) in ducks ,pigeons, turkeys and chickens respectively . An *Anaplasma* –like organism form ,rounded and piriform shaped with varying sizes 0.83-4.15 μ . Parasitemia in chickens, pigeons, turkeys, and ducks respectively were 1.7%, 2.5%, 2.6%, 2.8%.

INTRODUCTION

Poultry products are important sources of protein for human throughout the world. Poultry diseases can be divided into five groups namely, bacterial, viral, fungal, parasitic and nutritional (1). Parasitic diseases in particular haemoparasites have been identified as the major impediment to chicken health world wide (2). Parasites affecting both wild and domestic birds (3), one of these a micro-organisms belonging to the genus *Aegyptianella* are rickettsial pathogens, seriously affecting majority of birds (4,5).

Aegyptianella was first described by carpano (1929) in both chickens and geese in Egypt (6). *Aegyptianella spp* occur within the erythrocytes and the morphology was described by (7). *Aegyptianella spp* are transmitted by the fowl tick *Argas persicus* (8,9).

The presence of those rickettsial organisms in the red cells of birds may result in conditions varying from a healthy carrier to a highly pathogenic and often fatal disease (6,10). The main clinical signs of acute *Aegyptianellosis* are fever, anemia, anorexia, diarrhea, pale discoloration of legs, staggering gait and jaundice (7,10,11). The aim of this study was to determine the prevalence of *Aegyptianella spp* affecting chickens, ducks, turkeys and pigeons in Mosul due to scarcity of publication in this area .

MATERIALS AND METHODS

A total of 205 blood samples were collected from 4 species of birds representing chickens (50), Ducks (75), Turkeys (30) and pigeons (50) from different regions in Mosul city.

Blood samples were taken from the brachial wing vein of each bird using disposable sterile syringes and needle (7). Thin blood smears were prepared and stained with Giemsa's stain to identify the *Aegyptianella* (12) .Parasitemia was determined according to the method described by (13).

Identification of this species was made according to morphological features (5,7,11) and ocular micrometer was used for measurements .

RESULTS

The total percentage of infection with *Aegyptianella spp* was 28.29% and the highest percentages of infection appeared in ducks and pigeon were 33.33% ,30% respectively . The percentage of infection in chickens and turkeys were 20 %, 26.66 respectively (Table 1). Examination of thin blood smears of four birds species revealed the presence of initial bodies in the cytoplasm of red blood cells .These organisms appeared in different forms such as Anaplasma –like bodies, ,round and piriform shaped resembling those of babesia, violet or dark colored and lacking pigmented granules (Fig1,2,3) .The measurements of the organisms in different birds species was presented in Table (2).

Table (2) also shows the range and mean of the parasitemia in different birds ,the highest was 2.8%in ducks and the lower was 1.7% in chickens.

Table 1: Percentage of infection with *Aegyptianella spp.* in different species of birds examined

Species of birds	Number of birds examined	Number of infected birds	Percentage of infection
Chickens	50	10	20
Ducks	75	25	33.33
Turkeys	30	8	26.66
Pigeons	50	15	30
Total	205	58	28.29

Table 2 : The morphology and parasitemia from *Aegyptianella spp* in blood smears of different species of birds examined

Species of birds	Measurements Range(mean) μ	The percentage of parasitemia range (mean)
Chickens	0.83-4.15(2.11)	0.5-4.5(1.7)
Ducks	0.83-3.99(1.86)	0.55-4.5(2.8)
Turkeys	0.83- 4.15(1.56)	0.85-3.5(2.6)
Pigeons	0.83-3.32(1.84)	0.85-6.2(2.5)




Fig 1: *Aegyptianella spp* (piriform shape) in blood smear of chickens(A) and ducks(B) ,1000X




Fig 2: *Aegyptianella spp* (round form) in blood smears of turkeys (A) and chickens(B) by using digital camera, 1000X




Fig 3: *Aegyptianella spp* in blood smears of chickens as small-anaplasma organisms by using digital camera, 1000X

DISCUSSION

In this study, the total percentage of infection with *Aegyptianella spp* was 28.29% and the percentage of infection in chickens, ducks, turkeys and pigeons were 20%, 33.33%, 26.66%, 30% respectively.

In Iraq, there are no studies on *Aegyptianella spp* prevalence with which to compare the results of this study, Al-Alousi *etal* (14) reported 2% infection rate with *Aegyptianella spp*. among this study of endoparasites in turkeys in Mosul city .

In northern, southern and central regions of Lorestan province in Iran , Desfoulan *etal* (15) recorded that the percentage of infection in chickens, ducks, geese and turkeys was 33.3%, 9.5%, 33.3%, 23.9% respectively with the total percentage of infection was 7.6% .Data from a study in wild turkeys from southern Texas revealed the presence a small, intraerythrocytic rickettsia *Aegyptianella pullorum* in 24 of 300 blood samples (6).

Other reports which were performed in different parts of the world, *Aegyptianella* is considered to be highly pathogenic to birds especially chickens with mortality risks in the range of 30-80% amongst young birds and the *Aegyptianella* is described as one of important hemoparasites in poultry production(2). Results from a study in west and southern Africa indicate that *Aegyptianella pullorum* infection can be quite widespread among free range birds raised in the tropics, for example, in the blood smear investigations, prevalence of 9% and 6% have been reported in Ghana and Zimbabwe respectively (16,17). *Aegyptianella pullorum* is common and is widely distributed in the Tansania and the percentage of infection which were recorded in chickens was 15.3% (2). The variability in the prevalence of *A. pullorum* between studies could be attributed to differences in management system of the birds, ecoclimates or level of infestation of *Argas persicus*. Swai *etal* (2) revealed that the high prevalence of *Aegyptianella* occurs in birds if they had history of contact with ticks and ectoparasites infestation considerably with geographical location of a varying scale.

In this study, during examination thin blood smear by light microscope. *Aegyptianella* appeared in different forms they are : small-anaplasma like organisms, round, oval to piriform and clover-like shape violet to dark coloured organisms with

different sizes ranged between 0.83-4.15 μ m in all species of examined birds .According to the morphological features and microscopical measurements the species of Aegyptianella may be related *A.pullorum*, these results were in agreement with (2, 4,6, 10, 15). Euzeby (18) and Rikihisa *etal* (19) referred that the genus Aegyptianella contains a single species namely *A. pullorum* and a large variety of birds species appear to be susceptible to infection with *A.pullorum*. The species of Aegyptianella which were found in wild turkeys (*Meleagris gallopavo*) in the USA, diagnosed as *A.pullorum* (6).Soulsby (11) revealed that there are two species of Aegyptianella occurs in birds they are : *A. pullorum* and *A. moshkovskii* Al-alousi *etal* (14) diagnosed the species *A. moshkovskii* in turkeys in Mosul city. however are different reports on the ultrastructural studies concerning Aegyptianellosis in birds in the world(15).

The mean percentage of parasitemia in chickens, ducks, turkeys and pigeons was 1.7%, 2.8%, 2.6%, 2.5% respectively and the infection could be regarded as subacute and chronic, these results were agreement with (10, 20, 21) and (11) referred that the Aegyptianellosis may be acute, subacute or chronic. Indigenous poultry rarely suffer the acute disease, but freshly introduced stock may die within a few days of the onset of the clinical entity, (20, 21) showed that identification of species of hemosporidians was possible when the intensity of infection was > 0.01% and when fully grown parasites were present in blood films.

ACKNOWLEDGEMENTS

This study was supported by the College of Veterinary Medicine, University of Mosul.

دراسة لطيفيلي *Aegyptianella spp* في بعض أنواع الطيور في مدينة الموصل /العراق

أيمن غانم سليمان

فرع الأحياء المجهرية، كلية الطب البيطري، جامعة الموصل، الموصل، العراق

الخلاصة

خلال هذه الدراسة تم فحص 205 عينة دم جمعت من أربعة أنواع مختلفة من الطيور وهي الدجاج (50) عينة والبط (75) عينة والديك الرومي (30) عينة والحمم (50) عينة من مناطق مختلفة في مدينة الموصل وذلك للكشف عن طفيلي *Aegyptianella spp*، بلغت نسبة الإصابة الكلية 28.29% وان الخمج قد شُخص في البط والحمم والديك الرومي والدجاج (33.33%) (30%) (26.66%) (20%) على التوالي ولقد ظهر الطفيلي على شكل أجسام صغيرة تشبه الانابلازما وأجسام دائرية وبيضوية وكمثرية الشكل و بإحجام مختلفة تراوحت بين 0.83-4.15 مايكرون وان معدل نسبة التطفل بلغ في كل من الدجاج والحمم والديك الرومي والبط 1,7% و 2,5% و 2,6% و 2,8% على التوالي .

REFERENCES

1. Calnek BW, Barnes HJ, Beard CW, Mc Dougals LR, Saif YM. Diseases of poultry. Low state University Press. Ames. Iowa ; 1997 : 1-1080.
2. Swai ES, Kessy M, Sanka P, Bwanga S, Kaaya JE. A survey on ectoparasites and heamoparasites of free-range indigenous chickens of Northern Tanzania. Livestock Research for Rural Development.2010; 22(9):<http://www.ird.org/17/6/conr 17070.htm>.
3. Peirce MA, Bevan BJ. Blood parasites of imported psittacine birds. Vet Rec. 1977 ; 2: 282-3.

4. Tarello W, Riccieri N, Aegyptianella like inclusion bodies in two birds of prey from central Italy. *Revue Med Vet.* 2003 ; 154 (11): 715-717.
5. Urquhart GM, Armour J, Duncan JL, Dunn AM, Jennings FW. *Veterinary parasitology.* 2nded. Blackwell Science. Oxford;2003.252p.
6. Castle MD, Christensen BM. Isolation and identification of *Aegyptianella pullorum* (Rickettsiales, Anaplasmataceae) in wild turkeys from North America. *Avian Dis.* 1985 ; 29(2): 437-45.
7. Thrall MA, Baker DS, Campbell TW, Denieola D, Fettman M, Lassen ED, Rebar A, Weiser G. *Veterinary Hematology and clinical chemistry.* Lippincott Williams and Wilkins. United State of America; 2004. p. 248-249.
8. Gothe R, Hartmann S. The viability of Cryopreserved *Aegyptianella pullorum* carpano. 1928 in the Vector *Argas* (Persicargas) Walker and Kaiser and Hoogstraal. *Parasitology Research.* 1979 ; 58(2): 189-190.
9. Leeftang P, Ilemobade AA. Tick-borne diseases of domestic animals in northern Nigeria. *Tropical Animal Health and Production.* 1977;9 (4): 211-218.
10. Tarello W. Aegyptianella-like organisms and microfilariae in a severely diseased Bittern (*Botaurus Stellaris Stellaris*). *Revue Med Vet.* 2001 ; 152(2): 189-193.
11. Soulsby E. L. *Helminths, arthropods and protozoa of domesticated animals.* 7th ed. Philadelphia, Bailliere Tindall, London ; 1986. p.756-757.
12. Hendrix CM. *Diagnostic Veterinary Parasitology.* 2nd ed. Mosby Inc. United States of America ; 1998. p.265-268.
13. Adam KMG, Paul J. Zaman V. *Medical and veterinary protozoology.* Churchill Livingstone. Edinburgh and London ; 1971.p. 126-133.
14. Al-Alousi TI, Daoud MS, Al-Bayati MM. A study of endoparasites of turkeys in Mosul – Iraq. *Iraq J Vet Sci.* 1994 ; 7(3): 123-128.
15. Dezfoulan O, Zibaei M, Nayeabzabeh H, Zakian N, Haghgoo M. Prevalence and ultrastructural study of *Aegyptianella spp.* in domestic birds from Southwestern area, Iran. *Iranian Journal of Veterinary Research, Shiraz University.* 2011 ; 12(2): 167-171.
16. Poulsen AB, Permin AO, Yelifari HL, Nasen P, Block P. Prevalence and distribution of gastro-intestinal helminthes and haemoparasites in young scavenging chickens in upper eastern region of Ghana. West Africa. *Preventive Veterinary Medicine.* 2000 ; 45 (3-4): 237-245.
17. Permin A, Esmann B, Hoj CH, Hove S. Ectogendo- and haemoparasites in free-range chickens in the Goromonzi district in Zimbabwe. *Preventive Veterinary Medicine.* 2002 ; 54(3): 213-224.
18. Euzeby JP. List of Bacterial names with standing in nomenclature : a folder available on the internet. *Int J Syst Bacteriol.* 1997; 47(2):590-2.
19. Rikihisa Y, Zhang C, Christensen BM. Molecular Characterization of *Aegyptianella pullorum* (Anaplasmataceae). *J Clin Microbiol.* 2003 ; 41(11): 5294-5297.
20. Valkianas G, Iezhova TA, Brooks D, Hanelt B, Brant SV, Sutherlin ME, Causey D. Additional observations on blood parasites of birds in Costa Rica. *Journal of Wildlife diseases.* 2004 ; 40(3): 555-561.
21. Valkianas G, Ravinder N, Sehgal M, Tatjana A, Iezhovos A, Smith TB. Further observations on the blood parasites of birds in Uganda. *Journal of Wildlife Diseases.* 2005 ; 41(3): 580-587.