



<http://doi.org/10.36582/j.Alkuno.2024.09.07>

Al-Kunooze University College

Journal homepage: <http://journals.kunoozu.edu.iq/1/archive> &
<https://www.iasj.net/iasj/journal/340>



Evaluation of Anti -Cancer Potential of luteolin 7 -o- glucoside extracted from *Cuminum cyminum* and *Ocimum basilicum* in colon Cancer Cell Line

Faten Essam Hussain^a, Falah Hasan Shari^b Ula M. Noor AlMousawi^d, Hiba Abdul-Razaq^e

a,c:Community Health Techniques Department, Health and Medical Techniques College in Basra, Southern Technical University, Iraq

mmuhsen@stu.edu.iq

b: Public Health Department, Basra Health Directorate, Iraqi ministry of health, Basra, Iraq

Corresponding author: Amlalrubbye737@gmail.com

Abstract

This study involved two plant species of different plant families, cumin (*Cuminum cyminum*) family Apiaceae and basil (*Ocimum basilicum*) family Lamiaceae of medical and economic importance. Aim: study luteolin o -7- glucoside in both species and evaluate its potential effect on colon cancer.

Method: extraction by (70% ethanol) using (Soxhlet and reflux), high-performance liquid chromatography (HPLC) for detection of compound , and *in vitro* evaluation of the potential biological activity of crude plant extracts and luteolin O-7 glycoside with anticancer activity of Colon HCT Cell lines were tested using (MTT) assay.

Results: of *in vitro* anticancer activity study conducted against colon cancer cell line showed moderate cytotoxicity of these plant extracts and luteolin 0 7 glucoside compounds against colon cancer. In which inhibition in viability of colon cancer cell by *Cuminum cyminum* (17.48% at concentration 250µg , 18.39 at concentration 500µg and 34 at 1000µg) , *Ocimum basilicum* showed inhibition in the viability of cancer cell (17.66 at 250 µg , 26.8 500µg and 40.6 at 1000µg) While luteolin 0 - 7-glucoside stander showed decrease in the viability of cancer cell by (29.14 at 250 µg 11.11 500µg and 39 at 1000µg

keywords: luteolin, colon cancer, cumin , basil

Introduction

Colorectal cancer is the third and second most common cancer in men and women. It is closely related to environmental and genetic risk.



<http://doi.org/10.36582/j.Alkuno.2024.09.07>

Al-Kunooze University College

Journal homepage: <http://journals.kunoozu.edu.iq/1/archive> &
<https://www.iasj.net/iasj/journal/340>



Approximately 5% of colon cancers are caused by two genetic disorders: Familial adenomatous polyposis and Lynch syndrome ⁽¹⁾.

The transformation of normal colon cells into precancerous and ultimately cancerous cells requires a combination of somatic and/or genetic changes over approximately 10 to 15 years ⁽²⁾.

Cumin

Cumin (*Cuminum cyminum* L.) is a plant belonging to the Apiaceae family, native to the Mediterranean region, Turkestan and Egypt, and spread to many regions of the world. It is one of the oldest and most important commercial plants, and its cultivation usually requires 3-4 months of summer. It is drought resistant and grows in the Mediterranean climate ⁽³⁾.

Different parts of the fennel plant (leaves, buds, roots, and flowers) have similarities and differences ⁽⁴⁾.

Cumin aldehyde, α -Sabinin, α -Flandren, α -Kadinin, α -Pinene, p-Cymene, Linalool, α -Phellandrene, Safranal, Limonene, Linalyl acetate, Luteolin, 1,8-Cineole, Methyl eugenol, Geraniol, α -Terpineol acetate

Basil

Basil (*Ocimum basilicum* L.) is a plant belonging to the Lamiaceae family, an important oilseed, culinary and medicinal plant. The essential oil of the plant, which grows in tropical and subtropical regions, is used in oral health and dental products as well as in the spice and food industry. O. basilicum, Ocimum, Lamiaceae, Lamiaceae, Class Magnolia, Phylum Magnolia ve Kingdom Plantae cinsinin bir slizidir ⁽⁵⁾

Materials and methods

Materials

Trypsin/EDTA\Capricorn, DMF\Santa Cruz, RPMI 1640\Gibco, MTT Stain\Sigma, Fetal Bovine Serum\Gibco, Ethanol\BDH.

Methods of Extraction

Two hydroalcoholic extracts of basil and cumin were prepared by the reflux and Soxhlet extraction method.



<http://doi.org/10.36582/j.Alkuno.2024.09.07>

Al-Kunooze University College

Journal homepage: <http://journals.kunoozu.edu.iq/1/archive> &
<https://www.iasj.net/iasj/journal/340>



Reflux Extraction

Reflux extraction Crude extract samples were prepared by mixing 25 mg of leaf powder with 150 ml of 70% ethanol and boiling in a reflux at 70°C for 45 min.

Filter the ethanol extract, place it in a vial, and store at -4°C until used for analysis⁽⁶⁾

Soxhlet Extraction

Ethanol extract from Soxhlet extraction equipment, ethanol extract is prepared from dry cumin seeds using continuous Soxhlet extraction equipment.

The crude extract sample was prepared by boiling 15 g of dry cumin seeds (150 ml) with ethanol (70%) in a Soxhlet extractor. Extractions were performed for 10 h and the ethanolic extract of each extract was filtered; both crude extracts were stored at -4°C⁽⁷⁾

High-performance liquid chromatography (HPLC) analysis: HPLC conditions ⁽⁸⁾

- Mobile phase: a mixture of 80% water and 20% trifluoroacetic acid in an acetonitrile solution
- C-18 chromatography column
- 1.0 milliliters per minute (ml/min)
- Column temperature: 22 degrees Celsius
- 20 µL of injection volume
- A UV detector with a wavelength of 290 nm is used for detection.

Cytotoxicity study

Methods

Cell culture

A mixture of cancer cells and normal cell lines were procured from the IRAQ Biotechnology Cell Bank Unit in Basra and cultured in RPMI-1640 with the addition of 10% fetal bovine, 100 units/mL penicillin, and 100 µg/mL streptomycin. After each passage in Trypsin-EDTA, the cells were reseeded at 70% confluence and incubated at 37 °C with 5% CO₂ ⁽⁹⁾.

Cytotoxicity Assays

To test the cytotoxic effect, a 96-well plate was used for the MTT cell viability assay.



<http://doi.org/10.36582/j.Alkuno.2024.09.07>

Al-Kunooze University College

Journal homepage: <http://journals.kunoozu.edu.iq/1/archive> &
<https://www.iasj.net/iasj/journal/340>



Each well was seeded with 1×10^4 cells for each cell line. The cells were exposed to c&u test compounds at a concentration of $\mu\text{g/ml}$ for every compound after 24 hours or after they had grown into a confluent monolayer.

The 72-hour mark was used to determine the cell viability. Take the medium out and mix it with 28 μL of MTT solution (2 mg/mL). Then, put the cells in an incubator at 37 °C for 2 hours.

Add 100 μL of DMSO (dimethyl sulfoxide) after removing the MTT solution and incubate at 37 °C with shaking for 15 minutes⁽¹⁰⁾. The assay was carried out in triplicate and the absorbance at 620 nm (the test wavelength) was measured using a microplate reader.


Following this formula, one may determine the rate of cell growth inhibition (% of cytotoxicity): In this equation, PR is the growth rate, A is the average eye velocity of the untreated well, B is the treated well's eye velocity, and IR is the difference between 100 and PR.

The Results

High Performance Liquid Chromatography HPLC

HPLC analysis of two basil and cumin extracts detected luteolin o -7-glucoside as the major ingredient along with several minor ingredients.

Luteolin o-7-glucoside was determined by comparing the retention time (Rt) with standard luteolin o-7-glucoside (7.25 min) as shown in the figure (3.1).





<http://doi.org/10.36582/j.Alkuno.2024.09.07>

Al-Kunooze University College

Journal homepage: <http://journals.kunoozu.edu.iq/1/archive> & <https://www.iasj.net/iasj/journal/340>



Figure (1) HPLC Chromatogram of standard luteolin o -7- glucoside




Figure (2) HPLC chromatogram of 70% ethanol reflux extraction of basil




Figure (3) HPLC chromatogram of cumin extracted with Soxhlet with 70% ethanol

Cytotoxicity study

Cell line cytotoxicity study shows the following results for colon cancer cell line



<http://doi.org/10.36582/j.Alkuno.2024.09.07>

Al-Kunooze University College

Journal homepage: <http://journals.kunoozu.edu.iq/1/archive> &
<https://www.iasj.net/iasj/journal/340>




Figure (4) Anti proliferative activity of *Cuminum cyminum* crude extract against colon carcinoma cells HCT cell line

Cell viability was evaluated according to MTT assay, and the results showed that the percentage of cell growth inhibition in the treatment was (250, 500 and 1000 µg/ml).




Figure (5) Anti proliferative activity of *Ocimum basilicum* crude extract against colon carcinoma cells HCT cell line

Cell viability was evaluated according to MTT assay, and the results showed that the percentage of cell growth inhibition in the treatment was (250, 500 and 1000 µg/ml).



<http://doi.org/10.36582/j.Alkuno.2024.09.07>

Al-Kunooze University College

Journal homepage: <http://journals.kunoozu.edu.iq/1/archive> &

<https://www.iasj.net/iasj/journal/340>




Figure (6) Anti proliferative activity of luteolin o -7 glucoside standard against colon carcinoma cells HCT cell line

Cell viability was evaluated according to MTT assay, and the results showed that the percentage of cell growth inhibition in the treatment was (250, 500 and 1000 µg/ml).

Discussion

In this study, results showed inhibition in viability of colon cancer cell by *Cuminum cyminum* (17.48% at concentration 250µg , 18.39 at concentration 500µg and 34 at 1000µg) as in figures (4)

Ocimum basilicum showed inhibition in the viability of cancer cell (17.66 at 250 µg , 26.8 500µg and 40.6 at 1000µg) as in figures (5)

While luteolin 0 – glucoside stander showed decrease in the viability of cancer cell by (29.14 at 250 µg 11.11 500µg 39 and at 1000µg) as in figures (6)

Luteolin has demonstrated the ability to hinder the proliferation of HCT116 breast cancer cells by regulating apoptosis and cell cycle arrest in a p53-dependent manner, without inducing autophagy⁽¹¹⁾ .

Chen et al. conducted a study to examine the impact of luteolin on the interruption of the cell cycle in human cancer cells. In 2018, researchers revealed that luteolin induces cell cycle arrest in LoVo human cancer cells at the G2/M phase, leading to death⁽¹²⁾ .



<http://doi.org/10.36582/j.Alkuno.2024.09.07>

Al-Kunooze University College

Journal homepage: <http://journals.kunoozu.edu.iq/1/archive> &
<https://www.iasj.net/iasj/journal/340>



Subsequent molecular investigations revealed that luteolin exerted an impact on the proliferation of LoVo cells by impeding the halt of the cell cycle at the transition from G2 to M phase and deactivating cyclin B1/cell division cycle 2 (CDC2), thereby prompting cell death. Apoptotic protease-activating factor 1 is activated through the involvement of cytochrome c and deoxyadenosine triphosphate ⁽¹³⁾

According to a study conducted in 2022 by Song et al., luteolin had the following effects on colorectal cancer cells: it slowed their proliferation by modulating the MAPK pathway, which in turn caused cell cycle arrest, DNA damage, and apoptosis. Additionally, luteolin had the opposite effect on tumor growth and CRC xenografts. real-time model ⁽¹⁴⁾ .

Furthermore, Song et al. 2022 demonstrated that cisplatin is an effective chemotherapeutic on colorectal cancer cells (CRCs), and that when combined with luteolin, cisplatin enhances its effects by lowering cell survival and increasing the Apoptosis rate of HCT-116 and HT-29 cell counts, in comparison to cisplatin treatment alone.

References

1. Xiao JB, Leng AM, Zhang YQ, Wen Z, He J, Ye GN. CUEDC2: multifunctional roles in carcinogenesis. *Front Biosci (Landmark Ed)*. 2019 Mar 01;24(5):935-946.
2. Bien SA, Su YR, Conti DV, Harrison TA, Qu C, Guo X, Lu Y, Albanes D, Auer PL, Banbury BL, Berndt SI, Bézieau S, Brenner H, Buchanan DD, Caan BJ, Campbell PT, Carlson CS, Chan AT, Chang-Claude J, Chen S, Connolly CM, Easton DF, Feskens EJM, Gallinger S, Giles GG, Gunter MJ, Hampe J, Huyghe JR, Hoffmeister M, Hudson TJ, Jacobs EJ, Jenkins MA, Kampman E, Kang HM, Kühn T, Küry S, Lejbkiewicz F, Le Marchand L, Milne RL, Li L, Li CI, Lindblom A, Lindor NM, Martín V, McNeil CE, Melas M, Moreno V, Newcomb PA, Offit K, Pharaoh PDP, Potter JD, Qu C, Riboli E, Rennert G, Sala N, Schafmayer C, Scacheri PC, Schmit SL, Severi G, Slattery ML, Smith JD, Trichopoulou A, Tumino R, Ulrich CM, van Duijnhoven FJB, Van Guelpen B, Weinstein SJ, White E, Wolk A, Woods MO, Wu AH, Abecasis GR, Casey G, Nickerson DA, Gruber SB, Hsu L, Zheng W, Peters U. Genetic variant predictors of gene expression provide new insight into risk of colorectal cancer. *Hum Genet*. 2019 Apr;138(4):307-326.



<http://doi.org/10.36582/j.Alkuno.2024.09.07>

Al-Kunooze University College

Journal homepage: <http://journals.kunoozu.edu.iq/1/archive> &
<https://www.iasj.net/iasj/journal/340>



3. H. Hajlaoui, H. Mighri, E. Noumi, M. Snoussi, N. Trabelsi, R. Ksouri, A. Bakhrouf, *Food Chem. Toxicol.* 2010, 48, 2186.
4. Akrami F., Rodríguez-Lafuente A., Bentayeb K., Antioxidant and antimicrobial active paper based on Zataria (Zataria multiflora) and two cumin cultivars (Cuminum cyminum). *LWT - Food Science and Technology.* 2015; 60 (2) : 929-933 .
5. Yoo H.S., Won S.B., Kwon Y.H. Luteolin Induces Apoptosis and Autophagy in HCT116 Colon Cancer Cells via P53-Dependent Pathway. *Nutr. Cancer.* 2022;74:677–686. doi: 10.1080/01635581.2021.1903947
6. Tawfeeq AA, Mahdi MF, Abaas IS, Alwan AH. Phytochemical and anti bacterial studies of leaves of Rosmarinus officinalis cultivated in Karbala, Iraq. *Al-Mustansiriyah J Pharm Sci.* 2018;17(2):9.
7. Al-Shammaa DA, Saour KY, Abdul-Khalik ZM. Phytochemical investigation for the main active constituents in Arctium lappa L. cultivated in Iraq. *Iraqi J Pharm Sci (P-ISSN 1683- 3597, E-ISSN 2521-3512).* 2013;22(1):18–24.
8. Gudzenko A. Development and validation of a method for the determination of rosmarinic acid in Mentha piperita L. using solid-phase extraction and RP-HPLC with photodiode array detection. *J Chem Pharm Res.* 2013;5(9):40–5.
9. Al-Shammari AM, Alshami MA, Umran MA, Almukhtar AA, Yaseen NY, Raad K, et al. Establishment and characterization of a receptor-negative, hormone-nonresponsive breast cancer cell line from an Iraqi patient. *Breast Cancer Targets Ther.* 2015;7:223.
10. Al-Shammari AM, Al-Esmaeel WN, Al Ali AAA, Hassan AA, Ahmed AA. Enhancement of Oncolytic Activity of Newcastle Disease virus Through Combination with Retinoic Acid Against Digestive System Malignancies. In: *MOLECULAR THERAPY. CELL PRESS* 50.
11. ahmani, A. H., Alzohairy, M. A., Khan, M. A., & Almatroodi, S. A. (2014). Therapeutic implications of black seed and its constituents in the management of lung cancer. *International Journal of Pharmacology*, 10(4), 206-213.
12. Yoo H.S., Won S.B., Kwon Y.H. Luteolin Induces Apoptosis and Autophagy in HCT116 Colon Cancer Cells via P53-Dependent Pathway. *Nutr. Cancer.* 2022;74:677–686. doi: 10.1080/01635581.2021.1903947
13. Chen Z., Zhang B., Gao F., Shi R. Modulation of G2/M Cell Cycle Arrest and Apoptosis by Luteolin in Human Colon Cancer Cells and Xenografts. *Oncol. Lett.* 2018;15:1559–1565. doi: 10.3892/ol.2017.7475



<http://doi.org/10.36582/j.Alkuno.2024.09.07>

Al-Kunooze University College

Journal homepage: <http://journals.kunoozu.edu.iq/1/archive> &
<https://www.iasj.net/iasj/journal/340>



14. Song Y., Yu J., Li L., Wang L., Dong L., Xi G., Lu Y.J., Li Z. Luteolin Impacts Deoxyribonucleic Acid Repair by Modulating the Mitogen-Activated Protein Kinase Pathway in Colorectal Cancer. *Bioengineered*. 2022;13:10998–11011. doi: 10.1080/21655979.2022.2066926.



<http://doi.org/10.36582/j.Alkuno.2024.09.07>

Al-Kunooze University College

Journal homepage: <http://journals.kunoozu.edu.iq/1/archive> &

<https://www.iasj.net/iasj/journal/340>

