

IMMUNIZATION TRIAL OF SHEEP AGAINST HYDATID CYST INFECTION

Esraa Abdul Wadood

Internal and Preventive Medicine ,College of Veterinary Medicine, University of Basrah,Basrah,Iraq

(Received 6 March 2007, Accepted 19 August 2007)

Keywords: Protoscoleces, Metacystod, Sheep

ABSTRACT

This study was conducted to evaluate the role of vaccination trial of sheep with different types of hydatid cyst antigens (hydatid cyst fluid, protoscoleces and laminated layer).in the prevention and control of hydatid cyst infection .The animals in all groups received *Echinococcus granulosus* egg as challenge after the third injection of antigens .

Serum antibody responded of vaccinated animals with different antigens studied during the period of study , the results revealed that ,the titer of antibody depend on indirect hemagglutination test after 21 days and 6 months were 640 and 320 respectively when hydatid cyst fluid antigen was used Also it was 320 and 160 respectively when protoscoliscs and laminated layer antigen was used .

The animals observed over a year after vaccination and the observation show reduction in the hydatid cyst growth in the vaccinated groups, whereas the control group show growth of small cyst in the liver and lung.

INTRODUCTION

Hydatid disease poses a world-wide problem. It is a parasitic zoonotic disease caused by infection by the metacystode stage (infective larvae) of the dog tapeworm

Echinococcus granulosus , which has a sustaining sheep-dog epizootic cycle difficult to break off (1).

Cystic Echinococcus in farm animals cause considerable economic problem due to loss of the edible liver, significant loss of meat and milk production and values of the fleece from infected sheep may also occur (2)

The vaccine will be used to prevent infection in livestock species such as sheep, which are commonly involved in transmitting the parasite, the vaccine will have its biggest impact in those parts of the world where hydatid disease is most prevalent, such as south America, China, Africa and the Mediterranean region(3).

Prevention is much better than cure ,scientists have also developed effective vaccine for sheep, giving rise to hope that a hydatids vaccine suitable for humans (3).

large controlled studies with sheep have shown that vaccination with arecombinant oncospherical *Echinococcus granulosus* antigen (E G95) induce high degree of protection, reduction of the cyst number in vaccinated animals by approximately 90-100% (4).

A high degree of immunity about 80% persists for six months (in the absences of re infection) and pregnant ewe vaccinated before lambing transfer high levels of antibody to their lambs.

The objective of the present study was to conduct a preliminary trial for production of hydatid cyst vaccines by using inactivated ovine hydatid cyst fluid ,protoscolices and laminated layer antigens.

MATERIALS AND METHODS

Antigen preparation :

Hydatid cyst fluid antigen prepared according to (5).

Protoscolicse antigen prepared according to (6).

Lamminated layer antigen prepared according to (7) .

Protein estimation at 260 and 280 nanometers according to (8) .

Antibody titer in each trial was estimated using Indirect heamagglutination test according to (9).

Experimental animals :

Sixteen ewe(16-23 months age)were divided into four groups equally,each animal in the first group inoculated subcutaneous with 1ml (0.5mg)ovine hydatid cyst fluid antigen with incomplete Freund's adjuvant and reinjected with the same antigen and dose after one week and followed by injection of the same antigen at one month and six month later.

The second group given 1ml (0.5mg) of protoscolices antigen at the same dose, route and interval .

The third group given 1ml(0.5mg/ml) of laminated layer antigen at the same dose , route and interval .

The fourth group given normal saline with adjuvant at same dose and route as a control group.

After (21) day and one month from the last injection blood samples were collected from all animals groups for antibody detection by indirect haemagglutination test (IHA).

At the same time two dogs grew up in cage and treated with Albendazole to ensure the dogs have no any gastrointestinal parasite ,and arecoline 1-2mg/kg as purgative . the parasite burden was determined by the number of protoscolices ingested and then the faeces examined after using aercolin as purgative although proglottides in the faeces is conclusive

(10) , also a post –mortem examination is the most reliable method of diagnosis was based on microscopic identification of *Echinococcus granulosus* egg in sample of duodenal mucus.

Then each ewe in all groups fed 50,000 egg/ animal as challenge (11) after the third injection of antigen so the control groups .after one year from challenge all the animals were slaughtered and the internal organs are inspected for presence of hydatid cyst.

RESULT

Depending on IHA, the titer of antibody due to injection of hydatid cyst fluid antigen after 21 day and 6 month was 640 and 320 respectively , and it was 320 and 160 respectively when protoscolices and laminated layer antigens were used whereas the titer in the control group 20 as table (1) show. ewe were observed for 12 month after the last injection of antigen and after that ewe were slaughtered and the internal organs inspected for present of hydatid cyst, we see there is reduction of cyst growth in comparison with control group which showed multiple cyst in different growth stages in the liver and lung .

There is a reduction in the average number of hydatid cyst in three group of experiment ,established in the ewe many of the vaccinated animals were completely protected against infection .

the ewe in the vaccine trials had no viable hydatid cyst when examined one year after the experiment , that mean possible vaccines is essential in order to supplement the existing methods of breaking the *Echinococcus* life cycle.

DISCUSSION

The result showed that the immunological reaction were same when the groups of animals inoculated with the different antigens of hydatid cyst ,the study showed a good level of immunity in the three groups of animal .This agree with

(12) and (12) ,which studied the serum antibody response in sheep infected with hydatid cyst following parenteral immunization with hydatid cyst fluid ,they found that 60% of the infected animals responded to antigen by rapid production of antibodies.

The reduction in the average number of hydatid cyst is established in the vaccinated ewe ,many of the vaccinated animals were completely protected against infection with *Echinococcus granulosus* eggs . sheep in the vaccine trials had no viable hydatid cyst when examined one year after the experiment, that mean possible vaccines is essential in order to supplement the existing methods of breaking the *Echinococcus granulosus* life cycle .

Vaccination of the intermediate host is has moved forward considerably in recent years following the development of recombinant vaccine against *Echinococcus granulosus* ,arrange of different antigens including cyst fluid (13 ; 14; 15) .

Cyst membranes (16)and protoscoleces (17) had been used as proto type vaccine against hydatid cyst (18) . (19) showed the immunity generated by two injection of the vaccine persists for at least 12 months ,therefor annual vaccination of domestic livestock is recommended . the vaccine conferred a high degree of protection against challenge with different geographical isolates of *Echinococcus granulosus* (20) .

Table (1) mean of IHA test use different antigen

Type of antigen	Titer after 21	Titer after 6 months
Hydatid cyst fluid	640	320
Protoscoliece	320	160
Lamminated layer	320	160
Control	20	20

محاولة تلقيح الأغنام ضد الإصابة بالأكياس المائية

أسراء عبد الودود

فرع الطب الباطني والوقائي، كلية الطب البيطري، جامعة البصرة، البصرة، العراق

الخلاصة

صممت هذه الدراسة لتقييم محاولة تلقيح الأغنام باستخدام مستضدات مختلفة من الأكياس العدرية (مستضد سائل الكيس العدري، مستضد الرؤيسات الأولية والطبقة الصفائحية) للوقاية والسيطرة من الإصابة بداء الأكياس العدرية حيث تم ملاحظة الحيوانات لمدة سنة بعد التلقيح . حيث تم اعطاء الحيوانات في كل المجاميع وضمنها مجموعة السيطرة بيوض ديدان المشوكة الحبيبية كجرعة تحدي بعد الحقنة الثالثة من المستضد . تم ملاحظة مستوى الاجسام المضادة باستخدام اختبار التلازن الدموي غير المباشر حيث كان معدل معيار الاجسام المضادة عند استخدام سائل الكيس العدري بعد احدى وعشرون يوم وستة اشهر 640 و 320 على التوالي وعند استخدام مستضد الرؤيسات الأولية ومستضد الطبقة الصفائحية 320 و 160 على التوالي . ومن خلال ملاحظة الحيوانات بعد سنة من التلقيح وذبح الحيوانات حيث تم ملاحظة اختزال نمو الاكياس العدرية في كل المجاميع الملقحة بينما لوحظ نمو اكياس صغيرة في الرئة والكبد في مجموعة السيطرة .

REFERENCES

1. Ray ,R.De.Pk. and Karak ,K.(2002). Combined role of casoni test and indirect haemagglutiation test in the diagnosis of hydatid disease medical microbiology and parasitology .20,2:79-82 .
2. Torgerson ,P.R.; Dowling,P.M. and Abo-shehada ,M.N.(2001). Estimating the economic effect of cystic Echinococcosis .3.Jordan,a developing country with lower-middle income.Ann.Trop.Med.Hyg.95:595-603 .
3. Lightowlers ,M. (2003).Hydatid vaccine .Pak.J.Med Sci.19 (4) :268-271 .

4. Heath,D.D.; Jensen ,O. and Lightowlers M.W. (2003). Progress in control of hydatidosis using vaccination –areview offormulation and delivery of the vaccine and recommendations for practical use in control programmes .Acta. Tropica 85:133-143.
5. Bombardieri,S. and Gioradano, F. (1974). An evaluation of an Agar gel diffusion test with crude and purified antigen in the diagnosis of hydatid disease. Bull.WHO,15:525-530.
6. Dottorini, S. and Carmelo, T.(1978). *Echinococcus granulosus*: compares between antigens in scolices and hydatid fluid. Inter. J. For Parasitology. 8: 259-265.
7. Taherkhani, H. and Rogan, M.T. (2000). General characterization of Laminated layer of *Echinococcus granulosus* . Iran . J. Med. Sci. 25 (3&4) : 95 – 104.
8. Hudson, L. and Hay, F.C. (1989).Practical Immunology. 3rd Ed. Blackwell Scientific Publications. Oxford .London. Pp. 348-350.
9. Csizmas, L.(1960). Preparation of formalinized erythrocytes. Proc. Scoc. Exp. BioL.Med. 103:157-160.
- 10.Thienpont,D.;Rochette, F.;Vanparijs,O.F.J.(1979). Diagnosing Helminthiasis through coprological examination .Janssen Research Foundation , BeerseBelgium .
- 11.Torgerson,P.(2002). Transmission dynamics of taeniid parasites in animal hosts,P.22-235.In Craig,P. and Pawlowski ,Z. (ed) ,cestode zoonoses :echinococcosis and cysticercosis,an emergent and global problem .IOSPress,Amsterdam, TheNether Lands.
- 12.Marshall,WL.; Michael,DR. and Rhonda,DH. (1986) . serum antibody response following parenteral immunization with hydatid cyst fluid in sheep infected with *Echinococcus granulosus* .Am.J. Trop.Med.Hyg. 35:81-823 .
- 13.Craig P.S.(1997). *Echinococcus granulosus* immunodiagnosis and vaccination ,a perspective.Parasitologyia . 39:345-347 .
- 14.Dada, B.J. and Belino, E.D.(1981). Immunization of sheep against cystic hydatidosis with homologous and heterologous metacestode antigens. Int.J.Zoonoses.8:20-25 .
- 15.Heath,D.D.; Lawrence ,S.B. and Yong,W.K.(1992). *Echinococcus granulosus* in sheep:Transfer from ewe to lamb of (Arc5) antibodies and oncosphere –killing activity, but not protection .Int.J.Parasitol.22:1017-1021 .
- 16.Economides,P.E.; Larrieu,J. and Orlando,D.(2001). Evolution of programmesfor control of *Echinococcus granulosus* (examples) ,P.204-209.In.Eckert, J.;Gemmell, M.A. ;Meslin,F.X. and Pawlowski ,Z.S.(ed)WHO/OIE .manual on *Echinococcus* in humans and animals:a public health problem of global concern. World Organisation for Animal Health ,Paris,France.

17. Pauluzzi, S.; and DeRosa, F. (1969). Experimental vaccine prophylaxis control of experimental secondary Echinococcus granulosus in BALB/C mice. Ann. Sclavo. 11: 518-530.
18. Hernandez, A. and Nieto, A. (1994). Induction of protective immunity against murine secondary hydatidosis. Parasite Immunol. 16: 537-544.
19. Heath, D.D., and Lawrence, S.B. (1996). Antigenic polypeptides of Echinococcus granulosus oncospheres and definition of protective molecules. Parasito. Immunol. 18: 347-357.
20. Roberts, M.G.; Lawson, J.R. and Gemmell, M.A. (1986). Population dynamics in Echinococcosis and cysticercosis: mathematical model of the life-cycle of Echinococcus granulosus. Parasitology. 92: 621-641.