



<http://doi.org/10.36582/j.Alkuno.2024.09.02>

Al-Kunooze University College

Journal homepage: <http://journals.kunoozu.edu.iq/1/archive> &
<https://www.iasj.net/iasj/journal/340>



Effect of Caffeine and Chlorogenic acid on α -galactosidase Enzyme

Al-zamely Oda Mizil Yasser^a, Qasim N. Thewaini^b, Ameera M.Z. Fadli^c

a:Department of Chemistry, College of Science, University of Babylon, Iraq.

b:Department of Biotechnology, College of Biotechnology, Al-Qasim Green University, Iraq.

c: Department of Community health, Kut Technology Institute, Medill Technical University, Iraq.,

Corresponding Author, Email: oda.alzamely@gmail.com,

Abstract

A glycoside hydrolase enzyme known as α -galactosidase (EC 3.2.1.22) hydrolyzes the terminal alpha-galactosyl moieties found in glycolipids and glycoproteins. Bacteria that produce this enzyme are isolated from *Erysipelothrix rhusiopathia*. Following the release of α -galactosidase from the cells, the purification process involves precipitating the protein with 40% ammonium-sulfate, resuspending it, and dialyzing it using a dialyze tube cut-off number (10–14 KD) overnight against phosphate buffer (pH=7). Finally, the protein is further purified using DEAE-cellulose column chromatography with a (33.4) folds. At 45 °C, the enzyme had maximal activity, while 6.5 °C was its optimum pH. Through the use of SDS-PAGE technique, the molecular weight of the enzyme was estimated to be (55 KD). The enzyme's K_m and V_{max} were found to be (6.6 mM) and (833.3 μ mol/min) of protein, respectively, using p-nitrophenyl- α -D-galactopyranoside as the substrate. Following treatments of the purified enzyme with chlorogenic acid and caffeine, studies involving caffeine indicate that the inhibitor has a same affinity for the enzyme as the substrate pNPG. Measurements of the reaction rates at different concentrations of substrate and inhibitor observed a non-competitive inhibition and was shown the K_m value was ineffective (6.66) mM and decrease in V_{max} values (555.5) μ mol/min. While when we treated chlorogenic acid with α -galactosidase that is purified from *Erysipelothrix rhusiopathiae* this substrate acts as uncompetitive inhibition with substrate pNPG as the catalytic site that reflected the enzyme has single active that results from the change of K_m and V_{max} (3.22 mM) and (400 μ mol/min) respectively.

Keywords: coffee, caffeine, chlorogenic acid, α -galactosidase, *Erysipelothrix rhusiopathi*



<http://doi.org/10.36582/j.Alkuno.2024.09.02>

Al-Kunooze University College

Journal homepage: <http://journals.kunoozu.edu.iq/1/archive> &
<https://www.iasj.net/iasj/journal/340>



Introduction

The enzyme known as α -Galactosidases (EC 3.2.1.22) is an exo-type glycoside hydrolase that is responsible for catalyzing the breakdown of α -1,6-galactosidic linkages in galactose-containing oligosaccharides, including melibiose (galactose α -1,6-glucose), stachyose (galactose α -1,6-raffinose), and raffinose (galactose α -1,6-sucrose), which are frequently found in legumes and seeds ^(1; 2; 3). Humans, animals, plants, and microbes all have α -galactosidases ⁽⁴⁾. Different degrees of enzyme activity have been observed in the α -galactosidases isolated from different bacterial species, such as *Bacillus Stearothermophilus* ⁽⁶⁾, *Streptomyces griseoloalbus* ⁽⁷⁾, *Bifidobacterium bifidum* ⁽⁸⁾, and *Rmusbrockianus* ⁽⁵⁾. This enzyme can be utilized to break down α -galactooligosaccharides, improving the nutritional value of foods made with legumes ⁽⁹⁾. The cost of manufacturing determines whether industrial enzymes are commercially viable. The cost of the enzyme is mostly determined by the medium and fermentation technique that are used to isolate stable forms with high yields. As a result, it is best to separate high yielding and stable forms utilizing a medium devoid of repressors and containing product inducers ⁽¹⁰⁾.

Coffee beans, tea leaves, cocoa beans, cola nuts, and other plants are natural sources of caffeine (1,3,7-trimethyl xanthine), a natural alkaloid that is commonly found in coffee, tea, and soft drinks ^(11, 12, 13, 14). In the adult diet, the main sources of caffeine are tea and coffee, whereas in the diet of children, the main sources are chocolate and caffeinated soft drinks ^(15; 16). Arabica coffee beans ⁽¹⁷⁾ have a caffeine value of 10–12 mg/g, whereas Robusta coffee beans ⁽¹⁸⁾ have a caffeine content of 19–21 mg/g. Caffeine's ability to scavenge hydroxyl radicals ⁽¹⁹⁾ indicates that it has an antioxidant activity in addition to its well-known bioactive effects, which include improving alertness and reducing tiredness. A number of epidemiologic and clinical investigations conducted in the past several years have revealed that coffee drinking is linked to health advantages, including a decreased risk of type 2 diabetes ^(20; 21), Parkinson's disease, and Alzheimer's disease ⁽²²⁾; these findings are independent of caffeine intake.



<http://doi.org/10.36582/j.Alkuno.2024.09.02>

Al-Kunooze University College

Journal homepage: <http://journals.kunoozu.edu.iq/1/archive> &
<https://www.iasj.net/iasj/journal/340>



The primary phenolic components in coffee are chlorogenic acids (CGA) ($C_{16}H_{18}O_9$; mol. wt. = 354.31), an ester of caffeic acid and quinic acid. CGA is an ester of trans-cinnamic acids, including caffeic acid, ferulic and p-coumaric acids with (-) quinic acid ^(23; 24). They are thought to possess antioxidant qualities, which may be crucial in preventing oxidative damage to food, cells, and any organ ^(25; 26).

Experiments

Isolation of α -Galactosidase Producing Bacteria from *Erysipelothrix rhusiopathia*

The strain was employed to produce α -galactosidase. It was cultured to prepare the inoculum in tryptic soy agar for 24 hours at 37°C. A 50 mL conical flask containing 5 mL of thioglaculate broth was filled with a single new colony and shaken at 220 rpm for 24 hours at 37°C. A 250 mL Erlenmeyer flask with 50 mL of tryptic soy broth was filled with 1% of the resultant 24-hour culture, which was then kept at 37°C for 24 hours. The inoculum for solid-state fermentation (SSF) was this pre-culture.

Purification of α -galactosidase

Isolating the enzyme from cultivated cells is the initial stage in the purification process. Solid ammonium sulfate was added to the extract crude until it reached 40% saturation ⁽²⁷⁾. Proteins that had precipitated were recovered using chilled centrifugation (14000g for 20 min), resuspended, and dialyzed against phosphate buffer (pH=7.0) for a whole night. After the dialyzed extract (3mL) was equilibrated with 10 mM phosphate buffer (pH=7.0), it was put onto a DEAE-cellulose column (30 × 15mm). The column was cleaned using 2V phosphate buffer that was equilibrated with 10 mM (pH=7.0). Elution was performed at a flow rate of 0.5 ml/min using a gradient sodium chloride solution (0.1 M, 0.5 M, and 1.0 M), and fractions of 3 ml were collected ⁽²⁸⁾. Using a spectrophotometer UV (Nano-drop), the absorbance at 280 nm was measured after the proteins were eluted from the column. Each fraction's α -galactosidase activity was measured, and the fractions with the greatest enzyme activity were combined.



<http://doi.org/10.36582/j.Alkuno.2024.09.02>

Al-Kunooze University College

Journal homepage: <http://journals.kunoozu.edu.iq/1/archive> &
<https://www.iasj.net/iasj/journal/340>



Electrophoresis

Sodium dodecyl sulphate (SDS) was added to the separating gel at a concentration of 10% throughout the electrophoresis process. The sample was handled in accordance with Laemmli's procedure ⁽²⁹⁾. Bromophenol blue was acting as a front run marker after being heated to 100°C for five minutes (Page Run–R WSE– 1100). In an electrophoresis apparatus, the gels were run for 80minutes at 21mA and 150V. Following the run, Coomassie brilliant blue R-250 ⁽³⁰⁾ was used to label them for protein.

Enzyme assay

The test for α -Galactosidase activity using the De et al. ⁽³¹⁾ technique with p-nitrophenyl- α -galactopyranoside (pNPG) at a concentration of 0.003M as the substrate. The addition of sodium carbonate (0.1M Na₂CO₃) terminated the reaction, which was conducted at 45°C. At 410 nm, the quantity of released p-nitrophenol (pNP) was measured. The quantity of α -galactosidase that releases 1 μ mol of pNP (p-nitro phenyl) per minute under the specified test conditions ⁽³²⁾ is known as an enzyme activity unit (U).

Enzyme characterization Effect of temperature

By employing the enzyme assay at various incubation temperatures (30°C, 35°C, 40°C, 45°C, 50°C, 55°C, 60°C, and 65°C), the optimal temperature of α -galactosidase was determined.

Effect of pH

An enzyme assay method was used to investigate the effects of pH on α -galactosidase activity at various pH levels of potassium phosphate buffer solutions (4.0 - 8.0).



<http://doi.org/10.36582/j.Alkuno.2024.09.02>

Al-Kunooze University College

Journal homepage: <http://journals.kunoozu.edu.iq/1/archive> &
<https://www.iasj.net/iasj/journal/340>



Kinetic parameter

Kinetics experiments were conducted at pH 6.5 and 45 °C. Using Lineweaver-Burk and Michaelis-Menten plots, the maximal response rates (V_{max}) and Michaelis-Menten constant (K_m) were determined ⁽³³⁾. By applying Michaelis-Menten kinetics, one may ascertain the kinetic parameters of enzymes in accordance with the (PHYWE SYSTEME GMBH) software. The number of particles transformed in a unit of time is known as the rate of a chemical reaction. It is not necessary to assess the reaction rate for the breakdown of pNP- α -D-galactopyranoside (substrate), although it is dependent on the initial substrate concentration. All that is measured is the conductivity's change over time. Thus, the increase in conductivity is directly related to the rate of reaction.



Preparing the solutions

To calculate the values of the Michaelis-Menten equation a series of dilutions of the substrate (pNP- α -D-Galactopyranoside) was used. First preparer a 0.04 molar (pNP- α -D-Galactopyranoside) stock solution. To make a series of dilution solutions of (pNP- α -D-Galactopyranoside) (0.02; 0.01; 0.005; 0.0025 and 0.00125 mol/L). It is best to have the solution ready as soon as possible before beginning the experiment. In the reaction, the (pNP- α -D-Galactopyranoside) is converted into the electrically conductive products (p-Nitrophenol), which enhance the flow of current. Plotting the specific conductivity values versus time is shown in the "Specific conductivity" graphic. There is a short initial phase followed by an almost linear rise in specific conductivity. As per the program (PHYWE SYSTEME GMBH), the higher the initial concentration of substrate, the faster it increases conductivity (an increase in concentration leads to an increase in conductivity) and converts into V_{max} against [S].

To investigate the impact of caffeine and chlorogenic acid, first make a stock solution (0.04M) from (substrate pNPG plus caffeine) and (substrate pNPG plus chlorogenic acid) then prepare a dilution series of solution (0.02; 0.01; 0.005; 0.0025 and 0.00125 mol/L), using the (PHYWE SYSTEME GMBH) application, the conductivity was measured from the two stock solutions and converted into V_{max} versus [S]



<http://doi.org/10.36582/j.Alkuno.2024.09.02>

Al-Kunooze University College


Journal homepage: <http://journals.kunoozu.edu.iq/1/archive> &
<https://www.iasj.net/iasj/journal/340>



Table 1. A description of the intracellular α -galactosidase purification process from *Erysipelothrix rhusiopathiae*

Purification steps	Protein (mg/ml)	Activity (U/ml)	Specific activity (U/mg)	Total Volume (ml)	Yield Units in total	Yield (%)	Purification (fold)
Crude extract	39.56	730.3	18.46	5	3651.5	100	1
Ammonium Sulfate precipitation.	3.74	593.6	158.7	4	2374.4	65	8.6
DEAE-cellulose chromatography	0.537	331.15	616.6	3	993.45	27.2	33.4

Steps in purification: Activity (U/ml), Purification (fold), Yield (%), Protein (mg), Specific activity (U/mg).





<http://doi.org/10.36582/j.Alkuno.2024.09.02>

Al-Kunooze University College

Journal homepage: <http://journals.kunoozu.edu.iq/1/archive> &
<https://www.iasj.net/iasj/journal/340>




(Fig 1). *Erysipelothrix rhusiopathiae*'s α -galactosidase elution profile after DEAE-cellulose chromatography purification




<http://doi.org/10.36582/j.Alkuno.2024.09.02>

Al-Kunooze University College

Journal homepage: <http://journals.kunoozu.edu.iq/1/archive> &
<https://www.iasj.net/iasj/journal/340>



(Fig.2): Partially purified α -galactosidase produced from *Erysipelothrix rhusiopathiae* on 10% SDS-PAGE.



(Fig. 3). optimum temperature of α -galactosidase purified from *Erysipelothrix rhusiopathiae*




<http://doi.org/10.36582/j.Alkuno.2024.09.02>

Al-Kunooze University College

Journal homepage: <http://journals.kunoozu.edu.iq/1/archive> &

<https://www.iasj.net/iasj/journal/340>



(Fig. 4). optimum pH of α -galactosidase purified from *Erysipelothrix rhusiopathiae*




Fig. (5): Michaelis-Minton for *Erysipelothrix rhusiopathiae*'s purified α -galactosidase (substrate=pNPG)



<http://doi.org/10.36582/j.Alkuno.2024.09.02>

Al-Kunooze University College

Journal homepage: <http://journals.kunoozu.edu.iq/1/archive> &
<https://www.iasj.net/iasj/journal/340>




Fig. (6): Lineweaver-Burk plot for the α -galactosidase from *Erysipelothrix rhusiopathiae* that has been purified



<http://doi.org/10.36582/j.Alkuno.2024.09.02>

Al-Kunooze University College

Journal homepage: <http://journals.kunoozu.edu.iq/1/archive> &
<https://www.iasj.net/iasj/journal/340>




Fig. (7): Lineweaver-Burk plot for *Erysipelothrix rhusiopathiae*'s pure α -galactosidase (substitute with caffeine)




Fig.(8): Lineweaver-Burk plot for *Erysipelothrix rhusiopathiae*'s pure α -galactosidase (substrate containing chlorogenic acid)



<http://doi.org/10.36582/j.Alkuno.2024.09.02>

Al-Kunooze University College

Journal homepage: <http://journals.kunoozu.edu.iq/1/archive> &
<https://www.iasj.net/iasj/journal/340>



Discussion

A number of investigations into the anti-infective qualities of coffee on various microorganisms have been conducted over the past ten years (34, 35), and the results have shown possibilities for improving the management of a variety of medical conditions. Caffeic acid, caffeine, and chlorogenic acids are the primary components in roasted coffee extracts that exhibit this kind of action (36, 37).

As shown in (Table 1) The enzyme was partially purified by many steps. The first ammonium sulfate precipitation and the total enzyme activity was found to be in 40% saturation, which was then dialyzed to remove undesirable protein. Before dialysis, this crude α -galactosidase was assayed to determine the range of its activity. At pH (6.5), as seen in (Fig. 4), maximum enzymatic activity was observed and optimum temperature at (45°C) shown in (Fig 3). The findings agree with those of other α -galactosidases found in *Bifidobacteria* ⁽³⁸⁾ and *Lactobacilli* ⁽³⁹⁾ however, α -galactosidase found in fungi ^(40; 41) demonstrated maximum activity at pH levels that are acidic. After concentrating and further purifying the enzyme preparation using DEAE-cellulose chromatography (Fig. 1), α -galactosidase was purified by a 33.4-fold using the aforementioned purification processes, resulting in an increase in specific activity (616.6 U/mg) and a yield of 27.2%.

The enzyme's purity was assessed using SDS-PAGE of the purified fractions. The figure (2) illustrates the band separation. Standard markers and purified α -galactosidase were electrophoresed in a 10% (w/v) polyacrylamide gel under denaturing conditions. This method was also beneficial for counting the number of tubes that were collected and contained the separated band. The isolated enzyme displayed a single band measuring 55 KDa. This method was employed to gather comparable tubes containing just pure α -galactosidase. The purification α -galactosidase from *L. acidophilus* had a molecular mass of (45 KDa), as determined by SDS-PAGE ⁽⁴³⁾. Our result is different from that obtained by ⁽⁴⁴⁾, which was based on the zymographic analysis. The molecular weight of the purified enzyme from *PSEUDOMONAS* SP. MCCMB3 was approximately (80 KDa). The molecular mass of α -galactosidase was slightly different from that of earlier purified α -galactosidase from other bacteria, such as *Saccharopolyspora erythraea* (45 kDa) ⁽⁴²⁾.



<http://doi.org/10.36582/j.Alkuno.2024.09.02>

Al-Kunooze University College

Journal homepage: <http://journals.kunoozu.edu.iq/1/archive> &
<https://www.iasj.net/iasj/journal/340>



α -galactosidase which is purified from *Erysipelothrix rhusiopathiae* bacteria causing halitosis and detected the optimum temperature at (45 C⁰) and optimum pH (6.5) and the activity with different concentration of substrate pNPG with constant concentration of enzyme, the enzyme α -galactosidase for these physical properties was obey Michaelis-Menten equation by showing the Michaelis-Menten curve as in fig.(5,6). Using p-nitrophenyl- α -D-galactopyranoside (pNPG) as the substrate, the K_m and V_{max} values for the isolated enzyme were ascertained to be 6.66 mM and 833.3 μ mol/min per mg of protein, respectively.

The enzyme's K_m value is greater than the *Lactobacillus reuteri* K_m value of 0.48 mM published by ⁽⁴⁵⁾. On the other hand, it was less than that of the α -galactosidase that was isolated from different microbes using the same substrate (pNPG). Based on the K_m values from *Bifidobacterium adolescentis* (0.9 mM) ⁽⁴⁷⁾ and *Citrobacter freundii* (5 mM) ⁽⁴⁶⁾ that other researchers have published, which could be that the isolated enzyme in this study contain isoenzyme compared with another source which has another different value of K_m and V_{max}.

To examine how caffeine affected the activity of α -galactosidase, which was isolated from *Erysipelothrix rhusiopathiae*, it was found that caffeine inhibited the enzyme (pNPG is a substrate S1) when it was applied to other individuals. A reduction in the V_{max} values of α -galactosidase was seen in (Figure 7) when α -galactosidase reactions with equal quantities of both substrates, pNPG (S1) and caffeine substrate (S2), were assessed kinetically in the presence of increasing doses of caffeine. Caffeine-related experiments indicate that the inhibitor and the enzyme's substrate, pNPG, have similar affinities. Similar affinities are shown by the fact that the molar concentration of caffeine at which it starts to impact enzyme activity is rather close to the concentration of the pNPG substrate. Measurements of the reaction rates at various substrate and inhibitor concentrations revealed a non-competitive inhibition, a decrease in V_{max} values (555.5) μ mol/min that is less than that of the enzyme without inhibitors (833.3) μ mol/min, and an ineffective K_m value (6.66 mM) as shown in Figure (7). By binding to the enzyme at the active site, which is distinct from the substrate binding site, caffeine functions as an inhibitor. As a result, it may bind to both the free enzyme and the enzyme-substrate complex, lowering (V_{max}) while having no impact on (K_m).



<http://doi.org/10.36582/j.Alkuno.2024.09.02>

Al-Kunooze University College

Journal homepage: http://journals.kunoozu.edu.iq/1/archive_&https://www.iasj.net/iasj/journal/340



When we treated chlorogenic acid with α -galactosidase that is purified from *Erysipelothrix rhusiopathiae* this substrate acts as uncompetitive inhibition with substrate pNPG as the catalytic site that is reflected the enzyme have single active ⁽⁴⁸⁾, that is become from the change of K_m and V_{max}

mM) and (400 μ mol/min) respectively as shown in figure (8). We found a decrease in both (K_m) and (V_{max}) because chlorogenic acid binds reversibly to the enzyme-substrate complex, resulting in the formation of an inactive (ESI) complex.

Coffee is an inexpensive, naturally occurring product that may be purchased in bulk. The fact that coffee is ingested in a concentrated form (6–10%) as opposed to numerous medical infusions that have demonstrated such an effect in vitro and are typically eaten at 1-2% ⁽⁴⁹⁾ is another benefit of coffee as an antibacterial beverage. Within the parameters of the study, it can be concluded that using mouthwash on a regular basis to supplement mechanical halitosis management methods may help prevent or block the production of enzymes by bacteria that cause halitosis and lessen the likelihood that unpleasant odors would arise.

References

- 1- Viana PA, DE Rezende ST, Marques VM, Trevizano LM, Passos FML, Oliveira MGA, Bemquerer MP, Oliveira JS, Guimaraes VM (2006), Extracellular α – galactosidase from *Debaryomyces Hansenii* UFV-1 and its use in the hydrolysis of raffinose oligosaccharides. J. Agric. Food Chem. 54:2385-2391.
- 2- Comfort, D. A., Bobrov, K. S., Ivanen, D. R., Shabalin, K. A., et al., (2007). Biochemical analysis of *Thermotoga maritima* GH36 α - galactosidase (TmGalA) confirms the mechanistic commonality of clan GH-D glycoside hydrolases. Biochem. 46: 3319-3330.
- 3- Weignerova, L., Simerska, P., and Kren, V. (2009), α -Galactosidases and their applications in biotransformations. Biocatal. Biotransform. 27, 79-89.
- 4- Brouns, S. J. J., Smits, N., Wu, H., Snijders, A. P. L., Wright, P. C., de Vos, W. M., and van der Oost, J. (2006) Identification of a novel α -galactosidase from the hyper thermo philear chaeon *Sulfolobus solfataricus*. J. Bacteriol. 188: 2392-2399.
- 5- Fridjonsson O, Watzlawick H, Gehweiler A, Rohrhirsch T, Mattes R (1999), Cloning of the gene encoding a novel thermostable α -galactosidase from *Thermusbrockianus* ITI360. Appl. Environ. Microbiol. 65: 3955-3963.
- 6- Talbot G, Sygusch J (1990), Purification and characterization of thermostable α – mannase and α –galactosidase from *Bacillus stearothermophilus*. Appl. Environ. Microbiol. 56:3505-3510.
- 7- Anisha GS, John RP, Prema P (2009). Biochemical and hydrolytic properties of multiple thermostable α –galactosidases from *Streptomyces griseolalbus*: Obvious existence of a novel galactose- tolerant enzyme. Process. Biochem. 44:327-333.



<http://doi.org/10.36582/j.Alkuno.2024.09.02>

Al-Kunooze University College

Journal homepage: <http://journals.kunoozu.edu.iq/1/archive> &
<https://www.iasj.net/iasj/journal/340>



- 8- Goulas T, Goulas A, Tzortzis G, Gibson GR (2009), A novel α -galactosidase from *Bifidobacterium bifidum* with transgalactosylating properties :gene molecular cloning and heterologous expression. *Appl. Microbiol. Biotechnol.* 82:471-477.
- 9- Thananunkul D, Tanaka M, Chichester CO, Li T (1976), Degradation of raffinose and stachyose in soybean milk by α -galactosidase from *Mortierella vinacea*. *gel. J. Food Sci.* 41: 173-175.
- 10-Prakasham, R. S., Subba, Rao, C., Sreenivas Rao, R., Rajesham, S., and Sarma, P. N. (2005), Optimization of alkaline protease production by *Bacillus* sp. Using Taguchi methodology. *Applied Biochemistry and Biotechnology.* 120: 133-144.
- 11-Murphy, S. J., and Benjamin, C. P., (1981), The effects of coffee on mouse development. *Microbios Letters*, 17: 91–99.
- 12- IARC, (1991b), Caffeine. IARC Monographs on the Evaluation of Carcinogenic Risks to Humans, Vol. 51 (Lyon: International Agency for Research on Cancer) pp. 291–390.
- 13-Dlugosz, L., and Bracken, M. B., (1992), Reproductive effects of caffeine: a review and theoretical analysis. *Epidemiologic Reviews*, 14: 83–100.
- 14- Carrillo, J. A., and Benitez, J., (1996). CYP1A2 activity, gender and smoking, as variables influencing the toxicity of caffeine. *British Journal of Clinical Pharmacology*, 41: 605–608.
- 15- Shils, M. E., Olson, J. A., Shike, M., and Ross, A. C., (1999), *Modern Nutrition in Health and Disease*, 9th ed. (Baltimore: Williams & Wilkins).
- 16- Tanda, G., and Goldberg, S. R., (2000), Alteration of the behavioral effects of nicotine by chronic caffeine exposure. *Pharmacology, Biochemistry and Behaviour*, 66: 47–64.
- 17-Fox, G.P.; Wu, A.; Yiran, L.; Force, L (2013), Variation in caffeine concentration in single coffee beans. *J. Agric. Food Chem.* 61: 10772–10778.
- 18- Casal, S.; Beatriz Oliveira, M.; Ferreira, M.A , (2000). HPLC/diode-array applied to the thermal degradation of trigonelline, nicotinic acid and caffeine in coffee. *Food Chem.* 68: 481–485.
- 19- León-Carmona, J.R.; Galano, A (2011), Is Caffeine a Good Scavenger of Oxygenated Free Radicals? *J. Phys. Chem. B* 115: 4538–4546.
- 20- Agardh, E. E., Carlsson, S., Ahlbom, A., Efendic, S., Grill, V., Hammar, N., Hilding, A., Ostenson, C. G. (2004): Coffee consumption, type 2 diabetes and impaired glucose tolerance in Swedish men and women. *J. Intern. Med.*, 255: 645–652.
- 21- Bravi, F., Bosetti, C., Tavani, A., Bagmardi, V., Gallees, S., Negri, E., Fauschi, S., La Vecchia, C. (2007): Coffee drinking and hepatocellular carcinoma risk: a meta- analysis. *Hepatology*, 46: 430–435.
- 22- Lindsay, J., Laurin, D., Verreault, R., Hebert, R., Helliwell, B., Hill, G. B., McDowell, I. (2002), Risk factors for Alzheimer's disease: a prospective analysis from the Canadian study of health and aging. *Am. J. Epidemiol.* 156: 445–453.



<http://doi.org/10.36582/j.Alkuno.2024.09.02>

Al-Kunooze University College

Journal homepage: <http://journals.kunoozu.edu.iq/1/archive> &
<https://www.iasj.net/iasj/journal/340>



- 23-Clifford MN (2000), Chlorogenic acids and other cinnamates, occurrence, dietary burden, absorption and metabolism. *J. Sci. Food Agric.*80: 1033-1043.
- 24- Clarke RJ, Macarae R (1985). *Coffee: volume 1-Chemistry* Elsevier, England pp.1-7.
- 25- Cole RA (1984), Phenolic acids associated with the resistance of lettuce cultivars to the lettuce root aphid. *Ann. Appl. Biol.* 105: 129-145.
- 26- Moreira DP, Monteiro MC, Ribeiro-Alves M, Donangelo CM, Trugo LC(2005), Contribution of chlorogenic acid to the iron reducing activity of coffee beverages. *J. Agric. Food Chem.* 53: 1399-1402.
- 27- Shibuya H., Kobayashi H., Park G. G., Komatsu Y., Sato T., Kaneko R., Nagasaki H., Yoshida S., Kasamo K., Kusakabe I., (1995), Purification and some properties of α -galactosidase from *Penicillium purpuro genum*. *Biosci. Biotechnol. Biochem.*, 59: 2333-5.
- 28- Puchart V., Vrsanska M., Bhat M. K., Biely P., (2001), Purification and characterization of α -galactosidase from a thermo philic fungus *Thermo myceslanuginosus*. *Biochim. Biophys. Acta*, 1524: 27-37.
- 29- Laemmli UK (1970), Cleavage of structural proteins during assembly of the head of bacteriophage T4. *Nature*, 227: 680-685.
- 30- Kapitany RA, Zabrowsky EJ (1973), A high resolution PAS stain for polyacrylamide gel electrophoresis. *Anal. B Chem.* 56: 361-367.
- 31- Dey, PM and Pridham JB, (1972), *Biochemistry of α -galactosidases*, *Advances in Enzymology and Related Areas of Molecular Biology*, Vol. 36: 91-130.
- 32- Schnell, S., Chappell, M. J., Evans, N. D., Roussel, M. R., (2006) , The mechanism distinguishability problem in biochemical kinetics: The single-enzyme, single-substrate reaction as a case study". *Comptes Rendus Biologies.* 329 (1): 51–61.
- 33- Zhao, Y., Ma, S., Sun, Y., Huang, Y., Deng, Y., (2012) , Isolation, identification and enzyme characterization of a thermophilic cellulolytic anaerobic bacterium. *Wei Sheng Wu Xue Bao.* 52(9):1160-6.
- 34- Antonio, A. G., Moraes, R. S., Perrone, D., Maia, L. C., Santos, K. R. N., Iorio, N.L. P., et al. (2010); Species, roasting degree and decaffeination influence the antibacterial activity of coffee against *Streptococcus mutans*. *Food Chemistry*, 118:782–788.
- 35- Zhang L, Xue J, Ji Yao L, Zou L, Hao Y, Zhou X, et al. (2009), Effects of *Galla chinensis* on inhibition of demineralization of regular bovine enamel or enamel disposed of organic matrix. *Arch Oral Biol* 54:817–22
- 36- Daglia M, Papetti A, Grisoli P, Aceti C, Spini V, Dacarro C, et al. (2007), Isolation, identification, and quantification of roasted coffee antibacterial compounds. *J Agric Food Chem* ., 55:10208–13.
- 37- Almeida AAP, Farah A, Silva DAM, Nunan EA, Glo´ria MB. (2006) ; Antibacterial activity of coffee extracts and selected coffee chemical compounds against *Enterobacteria*. *J Agric Food Chem* 54: 8738–43.
- 38- Sakai K, Tachiki T, Kumagai H, Tochikura T (1987), Hydrolysis of α -D- galactosyl



<http://doi.org/10.36582/j.Alkuno.2024.09.02>

Al-Kunooze University College

Journal homepage: <http://journals.kunoozu.edu.iq/1/archive> &
<https://www.iasj.net/iasj/journal/340>



- oligosaccharides in soymilk by alpha-D-galactosidase of Bifido bacterium breve 203. Agric. Biol. Chem. 51: 315-332 .
- 39- Garro M, de Valdez GF, Oliver G, de Giori G (1997), Purification of alpha-galactosidase from *Lactobacillus fermentum*. J. Biotechnol. 45: 103-110.
 - 40- Kotwal SM, Gote M, Sainkar SR, Khan MI, Khire GK (1998), Production, purification and characterization of a constitutive intracellular agalactosidase from the thermophilic fungus *Humicolasp*. Process Biochem. 33: 337-343.
 - 41- Hin-chung W, Chien-A H, His-Lien Y, Wanchuang S (1986), Production, purification, and characterization of alpha-galactosidase from *Monascus pilosus*. Appl. Environ. Microbiol. 23: 1147-1152.
 - 42- Gote MM, Umalkar H, Khan MI and Khire JM, ,(2004), Thermostable alpha galactosidase from *Bacillus stearothermophilus* (NCIM5146) and its application in the removal of flatulence causing factors from soymilk. Process Biochemistry, 39:1723-1729.
 - 43- HEMA. T. A * AND HELEN PAPPA. T(2012), production , optimization and enzymatic removal of oligosaccharides from soy milk by alpha-galactosidase enzyme from rock soil *PSEUDOMONAS* SP. MCCMB3 Int J Pharm Bio Sci (2012) Oct; 3(4): (B) 770 – 781.
 - 44- Tzortzis G, Jay M, Baillon G, Gibson R (2003), Rastall Synthesis of a galacto oligosaccharides with alpha-galactosidase from *Lactobacillus reuteri* of canine origin. Appl. Microbiol. Bio technol. 63: 286-292.
 - 45- Lokage MA, Deepal CM (2001), Kinetic study of extracellular alpha galactosidase from *Citrobacter freundii*. J. Natl. Sci. Foundation Sri Lanka. 134: 97-106.
 - 46- Susanne L, Winfried H, Stefan P (1999), alpha-Galactosidase of Bifido bacterium adolescentis DSM 20083. Curr. Microbiol. 38: 101-106.
 - 47- Hankins C. N., Kindinger J. I., Shannon L. M., Legume, (1980), α -galactosidase form devoid of hemagglutinin activity. Plant Physiol., 66: 375.
 - 48- Brouns, S. J. J., Smits, N., Wu, H., Snijders, A. P. L., Wright, P. C., de Vos, W. M., and van der Oost, J. (2006) Identification of a novel α -galactosidase from the hyper thermo philear chaeon *Sulfolobus solfataricus*. J. Bacteriol. 188: 2392-2399.
 - 49- Bravo, L., Goya, L., & Lecumberri, E. (2007); LC/MS characterization of phenolic constituents of mate (*ilex paraguariensis*, St. Hil.) and its antioxidant activity compared to commonly consumed beverages. Food Research International, 40: 393– 405