

STAINING TECHNIQUE FOR HELMINTH PARASITES BY USE RED BEET (*BETA VULGARIS* L.) EXTRACT

Mohanad . F. A.Al-Amura ; Zahra .A. Hassen and Batool .H. AL-Mhanawi

Department of Microbiology, College of Veterinary Medicine, University of Basrah , Iraq

Department of pathology, College of Veterinary Medicine, University of Basrah , Iraq.

(Received 3 April 2012,Accepted 26 April 2012)

Key words: Helminthes, Trematoda, red beet.

ABSTRACT

Experimentally red beet (*Beta vulgaris* L.) extract had been used to stain different types of parasites *Lecithochirium acutum* (Trematoda), *Taenia* spp.*Dipylidium caninum* and cysticercus cyst for *Taenia* spp. (Cestoda), *Toxascaris leonina* and *Toxocara canis* (Nematoda) and *Neoechinorhynchus iraqensis* (Acanthocephala). Stained helminthes were acquired a good pigmentation with distinction their internal structure, fish's Trematoda *Lecithochirium acutum* appear well –defined of its internal structure, the oral and ventral sucker, cirrus sac, uterus, testes, ovary and vitallaria were appeared pink in color. The whole of mature and gravid segments of *Taenia* spp. were acquired the yellow to brown colour in both cirrus sac and vas deference besides differentiation of eggs in gravid segment, Scolex of *Taenia* spp. get brown in color.in addition segment of *Dipylidium caninum* acquired yellow colour. The cysticercus cyst was taken pink in color with distinguish of its hooks, while both *Toxascaris leonina*, *Toxocara canis* and *Neoechinorhynchus iraqensis* were appear pink to red in colour .

INTRODUCTION

The red beet (*Beta vulgaris*) plant belongs to the Chenopodiaceae family, it has huge roots which have contained different substantial elements like amino acid, sucrose, minerals, vitamins and dyes in fruit and roots (4).

(10) mentioned that *Beta vulgaris* has advantages in medicine such as it had a strong lipotropic and antioxidant activity inside the human body, (16) used the extraction of red beet as an active substance which was given to mice to protect them from exposure to gamma irradiation. Furthermore (8) has revealed that the red beet has an inhibitory effect on lung and skin tumors produced by the Epstein-Barr virus that was experimented in vivo.

In addition, the red beet has been used as a natural colorant in food products which it could be healthier or at least not harmful. Both (12) and (9) have referred that many consumers in the USA have replaced artificial dyes by natural dyes of red beet. In such a case, the juice of it contains all types of natural colorants which include yellow, orange, red and purple colors (1).

The aim of the study includes preparing a dye extract from red beet to stain some types of helminths as a substitution of cheap and easy-to-getting material.

MATERIALS AND METHODS

The following steps are :

- 1- In this experimental study the parasites had been chosen which included, Trematoda is *Lecithochirium acutum*, Cestoda (Scolex , mature and gravid segment of *Taenia* spp , gravid segment of *Dipylidium caninum*, cysticercus cyst. Nematoda is *Toxocara canis* and *Toxascaris leonina* and Acanthocephala *Neoechinorhynchus iraqensis*.
- 2- prepare extract solution from red beet : the plant of red beet was cut to small portions and boiled (1 kg with 1 liter normal saline) and set to filtration.
- 3- Stain of the specimens similar steps for aceto carmin which clarified in Palm (2004).

- A- Transferred the specimens from formalin 10% to the ethanol 70 % for at least 15 minutes .
 - B- The specimens were transferred to red beet solution and was kept in it for 1-2 days .
 - C- The specimens were transferred to 70% ethanol for 15 minutes to wash out excess stain .
 - D- placed the specimens in acid ethanol (2 ml of concentrated HCL in 100 ml of 70% alcohol) in order to remove excessive stain without loss of pigmentation .
 - E-Specimens again transferred to 70 % ethanol for 15 minutes .
 - F- They were transferred in to 90 % ethanol for 15 minutes .
 - G- The specimens were transferred to 100 % ethanol for 30 minutes and to ascending dish of 100 % ethanol.
 - H-The specimens were cleared by clearing agent (methyl salicylate).
- 4- mounted the specimen on slides by canada balsam .

RESULT

Through the picture under microscope helminthes samples were evidently stained by red beet extraction and have been well-defined in external and internal structure , whence the following point well be giving more details :

1- Cestoda : Scolex of *Taenia* spp appear brown in color - fig (1). The whole of mature and gravid segment appear yellow in color, in addition the cirrus, cirrus pouch, vas deference been very definitely and brown in color -fig (2) .The eggs of gravid segment also showed aggregation like chains in dark yellow appearance fig (3) . The eggs in gravid segment of *Dipylidium caninum* appear as multiple cell masses through their acquired yellowish in color fig (4). The cysticercus cyst were appear pink in staining with prominent of its hooks, fig (5,6) .

2- The anterior part of both *Toxocara canis* , *Toxascaris leonina* were appear red in colour by this stain, fig (7,8) whether the posterior end of *Toxascaris leonina* showed pink or slightly red in color with well –defined of anus pore fig (9).

3- Acanthocephala (*Neoechinorhynchus iraqensis*) acquired red in color, fig (10).

4- Trematoda (*Lecithochirium acutum*) appear well –defined of its internal structure that the oral and ventral sucker , cirrus sac yellowish in color , uterus show coils and brown color, testes, ovary pink in color) and vitellaria pink in color and like rose in apparent .fig (11,12) .



Figure (1) : Scolex of *Taenia* spp x 300



figure (2) : mature segments of *Taenia* spp,
(↖) cirrus pouch , (↗) cirrus x 400.




figure (3) show gravid segmenst of *Taenia* spp , eggs as masses () , genital pore () X 300



figure (4) : gravid segment of *Dipylidium caninum* engorgement with eggs. X 400



figure (5) : cysticercus cyst , arrow in hooks
X 300




figure (6) : hooks dendrites of Cysticercus
cyst X 400



figure (7) : Anterior end of *Toxacara canis* X
300



figure (8) : Anterior end of *Toxascaris leonina*
X 400




Figure (9): posterior end of *Toxascaris leonina* () anus region . x 400



figure (10) : *Neoechinorhynchus iraqensis* x 400



figure (11) : anterior end of *Lecithochirium acutum* () arrow refer to testes x 400



figure (12) : posterior end of *Lecithochirium acutum* () arrow refer to vitallaria x 400

DISCUSSION

The applied of this study by used red bet plant extract for staining different type of helminthes (Trematoda, Cestoda, Nematoda and Acanthocephala). (15) referred that necessary to explicit the internal and external parts of parasites in order to laid in classification and identical key. Recently it ran many research to evaluation the effect of natural dyes to stain different parasites, therefore (3) were used the aqueous extraction from red bet and Kujurat flower for staining Scolex of hydatid cyst in order to detection the viability.

The specimen were appeared well pigmentation with distinction of their internal structure. So that the successful staining has been ability the stain to distribution in the tissues of helminthes in different degree for permanent mount. The pigmented specimens were kept without decay for more than one year, also the Cestoda was acquired yellow in color while the Nematoda and Acanthocephala acquired red in color that due to the juice of red bet have red and yellow dyes with a strong stability. whence the (14), (7) and (5) were referred the red beet have many pigment component like Betacyanin (red color) and Betanin, isobetanin and betanidin (yellow color).

Nevertheless the variances helminthes structure might be given another reason to acquisition the dye at different color degree. (13) were mentioned the body or tegument of cestoda composed from muscle bundles, layer of glycocalyx and calcareous corpuscles and other substance, but the internal body of Nematoda have been pseudocoelom, therefore it would be high probably to acquire the red color.

The present result agreeable with (2) when found the kujurat red flower could stain of *Diphyllbothrium latum*, with purple color but the *Ancylostoma* spp. and *Enterobius* sp. were taken a red in color.

The structure of Trematoda like Cestoda but different from its digestive system and bodies more elastic than Cestoda, This Trait might be reason to acquired pink colour, therefore present result agree with (6), when found red colour of whole structure of *Fasciola hepatica* by alizarin and henna.

Conclusion From all above mention the regard red beet extract or its juice has prosperous stain for pigmentation different type of helminthes and as natural dye material

that it would be cheap dye than expensive artificial one and capable to developed it by using some modification to other groups of helminthes such as Leeches.

Aknownlogment : I would like to thank Dr Atheer ali in agriculture college \ basrah unviersity who help me for got some specimen and other effort .

تقنية تصبغ الديدان الطفيلية باستخدام مستخلص الشوندر

Beta vulgaris L.

مهند فارس عبد الحميد الامارة و زهرة عباس حسن و بتول حسين المهناوي
فرع الاحياء المجهرية، كلية الطب البيطري ، جامعة البصرة، البصرة، العراق.
فرع الامراض ، كلية الطب البيطري ، جامعة البصرة، البصرة، العراق.

الخلاصة

تضمنت الدراسة استخدام خلاصة الشوندر (*Beta vulgaris* L.) لتصبغ بعض انواع من الطفيليات وهي المتقوية *Lecithochirium acutum* والشريطيات *Taenia spp.* و *Dipylidium caninum* و الطور المتكيس لـ *Taenia spp.* و الديدان الاسطوانية *Toxascaris leonina* و *Toxocara canis* و شوكية الرأس *Neoechinorhynchus iraqensis*. الديدان المصبغة اكتسبت الصبغة بشكل جيد مع وضوح تراكيبها الداخلية، متقوية الاسماك *Lecithochirium acutum* حيث وضحت تراكيبها الداخلية والمتضمنة المحجم الفمي والبطني و كيس الذؤابة والامعاء و الرحم والخصى والمبيض مع بروز الغدد الحية باللون الوردي. اكتسبت القطع الناضجة والحبل لـ *Taenia spp.* اللون الاصفر- مع ظهور كل من كيس الذؤابة والناقل المنوي باللون الرمادي اضافة وضوح البيوض في القطع الحبلى، اما رؤيس الدودة الشريطية *Taenia spp.* اكتسب اللون الرمادي.بالضافة ان قطعة طفيلي *Dipylidium caninum* اكتسبت اللون الاصفر الطور المتكيس اكتسب اللون الوردي مع وضوح الاشواك ، بينما ظهر كل من الطفيلي *Toxascaris leonina* و *Toxocara canis* و *Neoechinorhynchus iraqensis* اللون الوردي الى الاحمر .

REFERENCE

1-Adriana, P. V.; Cachita ,C. D; Pop, L. and Purcarea, C. (2010). Histo-anatomical specific feature of red beet and sugar beet vitro plantlets . Fascicula : protectia mediului .vol .XV,(1) 740-748

- 2- Al-Adreesy, S. R. (2005).** A comparative study between commercial acid carmin and watery extract of kujarat flowers (*Hibiscus sabdariffa*) for staining some helminthes parasites. Al-Qadisiya J. Vet. Sci., 2(4):69-76.(in arabic)
- 3- Al-Azizz, Z. A. A. (2010) .** The use of some natural dyes for detection of viability of *Echinococcus granulosus* protoscolices in vivo . Minufiya Vet. J., 7(20: 433-439.
- 4- Al-Kubaasy, E. (2004).** Dictionary of grass and medicine plants. Dar Al-Kotob Al-Ilmiyah, Beirut. 566 pp. (in arabic).
- 5- Boris, N.; Zbigniew, P.; Aneta, P.; Pawel, S.; Wayne, T.; Tadeusz, M. and Stawomir, W. (2011).** Betalainic and Nutritional profiles of pigment –enriched red beet root. Food chemistry, 127: 42-53.
- 6- Daryani, A. Sharif, M. and Meigouni, M. (2010).** Staining of *Fasciola hepatica* by natural herbal dyes. Comparative clinical pathology, 20 (4): 305-308 .
- 7- Gasztonyi, M. N.; Daood, H .; Hajos, M.T. and Biase, P. (2001).** Comparison of red beet (*Beta vulgaris* var. *conditiva*) varieties on basic of their pigment component . J. Scie .Food & Agri, 81(9):932-933.
- 8- Govind, J.; Kapadia, A.; Harukuni, T.; Takao, K. and Hoyoku, N. (1996).** Chemoprevention of lung and tumor skin cancer by *Beta vulgaris* (beet) root extraction. Cancer letter, 100: 211-214.
- 9- Henry, B. S. (1996).** Natural food colours. In G. A. F. Hendry & J. D. Houghton (Eds.), Natural food colorants. London: Blackie Chapman & Hall .Pp 40-79.
- 10- Jones, R. G. and Storey, R. (1981).** The physiology and biochemistry of drought resistance in plants. Academic Press, Sydney. 508 pp.
- 11-Palm, H. W. (2004).** The Trypanorhyncha Diesing , 1868. PKSPL-IPB press , Bogor, 710 pp.
- 12-Philip,V. M. (1978).** How will we colors soft drinks of the future ? food prod .dev., 12: 24-25.
- 13-Roberts, L. S. & Janovy, J. J. (2005).** Foundation of Parasitology. 5th ed. Wm. C. Brown Publisher. The McGraw Hill Companies. Inc. 659 pp.
- 14- Usha, M.; Edgard, B.; Carvalho, M. Aand Wayne, R. (1998).** Growth and pigment production by hairy root culture of *Beta vulgaris* L. in bubble columnar reactor . Biotechnology letter, 20 (5): 469-474.

15- Yamaguti, S .(1959). Systema Helminthum: Vol. 2: The Cestodes of vertebrates.1st ed. Interscience Pub. Inc. N.Y. 860 pp.

16- Xiaoling, L. U; Yuping, W. and Zesheng, Z. (2009). Radioprotection activity of betalains from red beet in mice exposed to gamma irradiation . Eur. J. Phar. 615: 223-227.