اختيار تربة رملية مزيجة بكر لدراسة تأثير المبيد Cypermethrin على الإحياء المثبتة
للتروجين عند رطوبة السحمة الحقلية

(30-0)	(Typic quartzipsamments)	(Typic Torrifluent)
()	()	(-)
/ (535.4 -52.3)	/ (248.3- 48.6) ds/m(9.4-2.1)	
		(.0.01)
%2		Cypermethrin
(21 14 7)	/ (1.5 1 0.5)	600
(0.01)		

Abstract

The physical and chemical properties of different cultivated and uncultivated soils have been study. duplicate samples were collected for (0-30)cm depth, of Elluvial soils(Typic Torrifluent) Abukhasseb, Majdiah and (Typic quartzipsamments)from Shuaibah .The statstical analysis showed that there were high significant corlleation (R). Moreover all physical properties perdicated values were highly correlated at 0.01 level . The soils texture(clay,claylome-lomeysand) classify Fin to Cores (Abukhasseb, Majdiah)-(Shuaibah) continuously. Percent of clay in soils were ranged(52.33-535.4)g/kg with increised salinity (2.1-9.4)ds/m,and(48.6-248.3) g/kg calcium carbonate for Abukhasseb and Shuaibah soils respectively. The elluvial soils rejected due to differences properties for sepheir fine texture.There were no significsnt correlated between cultivated and uncultivated soils for same station. Effect of insecticide Cypermethrin was studied on loamysand of uncultivated Shuaibah soil , low cations,organic matter salinity, and high percentage of sand.The insecticide was applied in (0.5,1,1.5) mg/L.The concentration of insecticide and the incubation periode (7, 14 ,21) days affected were highly correlated at 0.01. level bacteria ,fungi and Azotobacter density there is no effect of the concentration and the incubation periode on nitrogenase activity.No correlated between the incubation periode on nitrogenase activity and soil moister.

	653	370	%23-6
/ 100-6	/ 250 -200		(.1997)
.(2002)			
	% 42 -5	²	185

48° 31 - 46°) (31° 18 -29° 07)
(Buring,1960) (35
(1988) (Typictorri Psamnents Catcorous and Maxed Hyperme)
(Al-Taie, 1970) %80-70.

(Mazurak et al., 1954)

.(2002)

. Dougrameji and Clor.1977

.(1997)

Misra et al .(1974) ()
(CO2) (CO2 NH3)
()
(2001) C/N
.Water Use Efficiency

Schlesinger and Pilmonis (1998)

.Hillel (1980)

.(1989)

(30-0)
2 .() ()
,(Page et al.,1982) (Black, 1965)

.(1) 8 30

%2

.(Core method)

600

(T3 T2 T1)

.(1990)

³ 120

/ (1.5,1,0.5)

Control

28

.(pw^{FC})

Nutrient

(21, 14, 7)

. Potato dextrose Agar

Agar

(Alexander, Most Probable Number (MPN)

(Brown et al.

glucose

1982)

Nitrogenase

.1964)

³ 20

5

) C₂H₄

28

2

C₂H₂

³ 3

³ 0.1

Gas Chromotography-GC

(2/ /

. 3

.SPSS

(1)

() (1a)

.Talati et al(1975)

(1b) 0.01
 $ds/m(9.4-2.1)$ / (535.4 -52.33)
 / (248.3- 48.6)
 / (43.85 -21.8)
 / $10^{-2} *(0.079-0.012)$

(2002)

/ (248.3-214.4)

(LoamySand)

/ 810.7 (1a)

ds/m 2.1 (1988)

%20.0 = (pw^{FC}) / 48.6

/ $10^{-2} *(1.68)$

(1980)

()

Cl dimethyl cycle

(2)

(3-phynoxy phenyl)

(2001)

()

(1990)

20-13 (Bohmont, 1999)

.Finch ,(1996)

(2)

1987

0.01

(3)

.Singh and Singh(2005)

Alexander(1982)

(4)

(12.31 CFU * 10⁴)

(14)

Vijay et al.(2005)

. 0.05

(black clay soil)

/ (25 10 5 0)

Cypermethrin

300 – 10

Methidathion

Gonzalez et al.(1992)

/

.Loam

chloropyrife methylpyrimifos

(5)

Radhi et al. (1991)

(pw^{FC})

(6)

%(23. 30 -22. 76-22.75 -21.59) T3 T2,T1 ,C

()

(kijn,1967)

.(Romanov, 1962)

(6a)

Cypermethrin

- .1988 .
- .1990 .
- .1997 .
- .28 (17)
- .1980 .
- .1990 .
- .1987 .
- .2002
- (28) . zea mays L.
- .195-185
- .2001 .
- 110 ACSAD
- .1989.
- .1987 .
- .(1997)
- .1988.
- 108 (10)5
- .136
- Alexander, M. 1982. Introduction to soil microbiology. John Wiley and Sons. New York.
- Al-Taie, E. 1970 . Salt effected and water logged soils of Iraq . Report to
Siminar on methods of an maliration of saline and water logged. soils. Baghdad .
- Black, C.A. 1965 . Method of soil analysis . Part (1). Physical properties Am. Soc. Agron.
Inc. puplisher Madison. Wisconsin , USA.
- Bohmont, Bert L. 1999. The Standard pesticides user, sGuide. 4th pp.531. Brown ,
A.D,1964.Aspects of batteries respons of the ionicenvironment. BactRev.28:296-
329.
- Buring.P.1960.Soil and soil condition of Iraq. Ministry of Agriculture Rep, of Iraq.
- Dougrameji, J.S. and M.AClor.1977.Case study on desertification greater Mussayeb
Project. Iraq United Nation conference on Desertification A Conf. 74/10 .
- Finch,C.1996.Industrial water soluble polymers royal society of chemistry Gonzalez.
Lopez, J., M. V. Martinez .Toledo and V.
- Salmeron. 1992. Effect of insecticide methidathion on agricultural soil microflora .J. of
Biology and Fertility of soils 13:173-175.
- Hillel .D.L. 1980 . Application of siol physics. Academic press Inc. New York ,N.Y.
11.Marcel Dekker Inc .,New York.

(1b)

Source	Silt	Clay
Sand	-0.980**	-0.988**
Silt		0.954**
Source	df	F
Sand	17	12.330
Silt	17	124.65
Clay	17	6.229

** Significant at the 0.01 level

جدول (2) المبيد سايبيرمثرين المستخدم في الدراسة

g/mol			
pyrethroid	Syngenta	cyano(3-phenoxyphenyl)methyl 3-(2,2-dichloroethenyl)-2,2-dimethylcyclopropanecarboxylate	cypermethrin
416.3

*The Pesticide Manual (2002)

F (3)

F				F	S. O. V.
0.261 ^{n.s.}	547.852**	13.735**	24.647**	3	
1.994 ^{n.s.}	367.434**	11.519**	128.061**	2	
0.447 ^{n.s.}	418.306**	4.081*	28.774**	6	
				12	
				23	Total

** Significant at the 0.01 level

(4)

	()			
	21	14	7	
8.17	4	14.25	6.25*	C
16.75	12.25	12.5	25.5	T1
9.33	7.00	11.25	9.75	T2
6.25	3.75	11.25	3.75	T3
	6.75	12.31	11.31	

(colong foring unite) CFU * 104

** Significant at the 0.01 level

: (2)

(ppm / h. / gm soil)

(5)

	()			
	21	14	7	
14.33	3.78	3.63	35.59	C
8.04	3.67	10.4	10.06	T1
7.04	6.76	3.03	11.34	T2
12.16	5.30	8.80	22.38	T3
	4.87	6.46	19.84	

(pw^{FC}) (6a)

% (pw)	()			
	21	14	7	
21.59	21.03	21.86	21.90	C
22.75	22.70	23.10	22.44	T1
22.76	22.91	22.40	22.98	T2
23.30	23.22	23.82	22.87	T3
	22.46	22.79	22.55	

F (6b)

F	Df	S. O. V.
n.s1.045	3	
n.s1.103	2	
n.s0.846	6	
	24	
	36	Total

a. Rsquared =.303 (Adj. Rsquared=-.017