



Research article

***Haemonchus contortus*: Review of recent molecular advances in anthelmintic resistance and vaccination**

Nuha Qasim Mohammed¹

Hiba Shehab Ahmed²

Noor Idan Jarad¹

Amal Hassan Al-Shabbani³

1- College of Veterinary Medicine, University of AL-Qadisiyah, Iraq.

2- College of Biotechnology, University of AL-Qadisiyah, Iraq.

3- College of Pharmacy, University of AL-Qadisiyah, Iraq.

Corresponding Author Email: nuha.allban@qu.edu.iq

Abstract

Haemonchus contortus is one of the world's major financial worms that attack ruminants. It is a blood-sucking parasite founded in the abomasum, particularly in cattle, sheep, and goats. Nematode infections may cause anemia, weight loss, or even death in animals that are severely affected. Current management practices against *H. contortus* largely depend on regular anthelmintic therapies in all countries with varying incidence across various regions.

H. contortus thus aims to form new action techniques in order to overcome resistance to anthelmintic agents. One option is a logical approach focused on a thorough knowledge of the molecular pathways in growth and reproduction cycles in the manufacture in modern anti-parasite drugs and vaccines. Key molecules may be defined as potential drug targets by a simple description of molecular, biochemical functions.

Besides, it is immediately essential to formulate immunological control of farm animal nematode infections. Important prevention has been accomplished after vaccination with native protein extracts, which shows that vaccination is possible. This paper explores the success of *H. contortus* science in the world. Particular fields of concern include epidemiological research, genetic analysis, and anthelmintic resistance identification using traditional and molecular techniques; morphological and chemical research of crucial molecules in mechanism expansion, parasitic organism-host interaction, and vaccine research. In the suggested form of these opinions, areas for potential exploration and alternatives for new or revised prevention strategies are described.

Keywords: Control, Vaccination, genetic, *Haemonchus contortus*, Ruminant.

Introduction

Nematodes have developed to take advantage of a wide variety of natural habitats. Although most stabilize a free lifestyle to reach maturity (Fig. 1), they depend on one or more hosts (1). A complex sequence of morphological features is taking place in several parasitic nematodes associated with migration from their hosts to create a mature infestation (2). Life cycles may include transitional hosts, vectors and, environmental-based stay when confronted with extreme and

unpredictable circumstances, including freezing or dehydration (3). The risks, namely predation, atmosphere, and immune reactions, of a wide variety, both plant and animal hosts have been adapted (1–3). The evolution of parasites affects public health (which amounts to 10 million life-years adjusting to disabilities), or indirectly by major financial failures in plant and animal production and by investing in parasite control (4). Anthelmintic medications are used as a



monitor, provided recurrently in livestock and through large-scale medical preventive initiatives (5). Although the effectiveness of these approaches was initially inevitable, the appearance or documenting

inefficiency in human-infectious organisms of medicinal-resistant veterinary worms challenges continuous prevention efforts (4–7)




Figure 1: The life cycle of *Haemonchus contortus*. <https://u.osu.edu/sheep/2019/07/30/ag-note-parasites-focus-haemonchus-contortus>

Vaccines provide an enticing alternative control method for these parasites: however, while two approved vaccines are presently accessible for veterinary purposes, the vast parasite heterogeneity and immune procedural mechanisms are significantly hindered in the production of vaccine; transcriptomic plasticity after the vaccine attempt may be used to prevent infections (8). It is thus obvious that modern, efficient control methods are essential. Helminths can respond to control measures and therefore evade them because of genetic variation (9). A deeper understanding of the nature and dynamics of this diversity across its spectrum can provide better knowledge of the pathways by which they are modified and can

recognize new goals that can be manipulated for influence (8,9).

The *H. contortus* (trichostrongylide) is a gastrointestinal worm of ruminants in tropical and temperate areas around the world, which has substantial effects on the economic and animal health of the sheep farm in a unique way (10). It also emerges as a prototype of a parasite nematode framework for functional and comparative genome sequencing, primarily due to its quick capability to collect resistance to medicines, its nearly equivalent tractability in controlled circumstances, the growth of enormous genomic inputs, and its special bond with other veterinary and medical nematodes of clad V (12,13).



Resistance to anthelmintics

The fundamental strategy for the containment of parasitic nematodes like *H. contortus* is focused upon the utilization of anthelmintics, but an anthelmintic resistance has over-grown. Resistance to anthelmintic in several areas of the planet was identified and published (6). A concise-short study of the nature of anthelmintic resistance in several nations refers to the occurrence of the resistance in countries like the United States, Mexico, South Africa, Australia, New Zealand, European countries, and China. Albendazole, fenbendazole, and ivermectin were used to treat in dairy goat farm (400 Goats) with confirmed infection (all above 80%) (14). Albendazole and ivermectin (IM) showed low activity both, but fenbendazole displayed high effectiveness (6). This study showed a fecal egg count reduction test (FECRT) of 23,72% for the routine dose rate (5 mg /kg) of albendazole. In this field, ivermectin has not been successful, with a prescribed FECR of 52.29 percent (0.2 mg/kg) and a 2-fold FECR of 89 percent (0.4 mg/kg) (6,14). The drug resistance was tested using albendazole, levamisole, ivermectin, and a mixture of albendazole and ivermectin to check performance against the nematode. The findings were successful in the cases of levamisole, ivermectin, and the combination. The achievement of an FECR was above 95% and an FECR of albendazole of below 95% (6,14).

Molecular identification of anthelmintics resistance

Resistance to Benzimidazole

Three specific β -tubulin SNPs were seen to be linked to *H. contortus* with resistance to benzimidazole (BZ),

identified as F167Y (TTC to TAC), E198A (GAA to GCA) and F200Y (TTC to TAC) (15). A nucleotide sequence modification has resulted in alteration in the target protein and decreased BZ binding to the β -tubulin (16). In five *H. contortus* species, including two sheep and three goats species, a multiplex PCR for the β -tubulin consistent with BZ resistance has been established, which suggests that the resistant genetic makeup is not found in any observed community of the nematode (15–20). In the meantime, an analysis utilizing PCR-SSCP in the area of the F200Y community *H. contortus* of sheep showed a plurality of homozygous prone genotype (21). Also, one analysis employed PCR-RFLP system to detect F200Y in *H. contortus* sheep populations, which revealed that the homozygous prone genotype is the most common (21). In various regions, a resistance was earlier revealed in the detection of all three known SNPs in a β -tubulin gene with PCR-coupled sequencing, which has been commonly used for the management of worm load (21).

Molecular approaches may be utilized, but they could not test resistance at the quantitative rates, as a diagnostic aid in detecting BZ resistance in populations (22). In fact, in combination with this field, molecular testing and biological testing are to be used (23). To date, the molecular research of the identification of BZ resistance in field samples, including several species of nematodes, has not been tested explicitly for this reason (22). Moreover, in fecal samples that included nematode eggs (Fig. 2) directly, molecular trials were not used, which could save a great deal of time. For now and on, whether current approaches or modern molecular production are used (22,23).



Figure (2): The egg shape of *Haemonchus contortus*.

https://www.researchgate.net/publication/262932569_Efficacy_and_Safety_of_Albendazole_against_Haemonchus_Contortus_Infection_in_Goats/figures?lo=1&utm_source=google&utm_medium=organic

Resistance to Ivermectin

IM also relates to the leading anthelmintic groups and has been used disproportionately, contributing to widespread resistance (24). In addition to numerous genes that code for IM targets and efflux pumps, resistance to IM is considered to be very multi-gene based nature (25). SNPs around the entire genome were scanned using a 2b-RAD sequencing tool for further IM resistance-associated SNPs in goats in both sensitive and resistant nematodes (24–26). 2962 SNPs and 2667 SNPs in resistant isolates have been identified (27). Among resistant or sensitive species, identical and relatively smaller genetic variations were observed (28). However, 208 SNPs with a significant variation, 24 of which were SNPs (29). This process is likely to be the primary selection of IM, with seven of the nine genetic markers expected to code for those proteins, which may play a crucial role in the IM target or efflux pump and even in receptor complex factor proteins, including membrane or neurons for transcriptional regulating proteins (30). It was suspected that these genes are correlated with IM tolerance. The findings

of this research revealed genes affecting the detection of IM and correlated with IM resistance in the genome of the nematode (31). Extensive SNP analysis utilizing the 2b-RAD sequencing method may be used to classify the worm (31).


Haemonchus contortus and diapause-presented genetic markers

Haemonchus contortus may evolve diapause while on an infective-free-living L3s (iL3s), or beginning of the L4 (Fig. 3) of the nematode stage in order to safeguard it against a harsh environment (32). Knowing this transformation could reveal major molecules as new drug attractions and provide new insights into protection and management (33). The L3 (iL3s)-based nematodes have been proven to have higher common phenotypic, behavioral, and biological characteristics with the long-standing phase of free-living *Caenorhabditis elegans* (34). The "dauer hypothesis" considers that the iL3 is comparable to that of the dauer stage of *C. elegans* in terms of developmental and



functional applications and controlled by a relatively similar method (35). The insulin-like component regulates the long-term development of *C. elegans* preserved in many parasitic species (35). The insulin-like signaling pathway genes are fork head transcription factor (Hc-daf-16), insulin

receptor (Hc-daf-2), phosphoinositide 3-kinases (PI3Ks) component catalytic subunit (Hc-age-1), regulatory subunit (Hc-aap-1), and phosphoinositide dependent protein kinase-1 gene (Hc-pdk-1) (32–35).



Figure(3): The Larva shape of *Haemonchus contortus*

https://www.researchgate.net/publication/266152011_Evaluation_of_the_Role_of_Galectins_in_Parasite_Immunity/figures?lo=1

They are intestinal genes which regulate the growth of the nematodes (36). It was found that *C. elegans* daf-2 mutant can be rescued by Hc-daf-2, which means that similar functions of Hc-daf-2 to Ce-daf-2. PI3K kinase, Hc-age-1, and Hc-aap-1 are the downstream of Hc-DAF-2. The *C. elegans* age-1-mutant can not be rescued by the Hc-age-1 like Ce-age-1 (36). RNAseq analysis shows that these four genes have the maximum amount of transcription expression in iL3, which displayed that these genes could play a significant role in controlling iL3 against an unpleasant surrounding environment (36). In summary, the results from these experiments indicate that insulin-like

signals are functionally conserved between the two worms. In contrast, a bioinformatic analysis further supports the mapped insulin-like signal route of transcriptomic and genetic information for this parasite (36).

Besides, researchers are actively working on other signals such as a TGF beta signal pathway that is important for the normal growth of *C. elegans*, and their function in *H. contortus* diapause in iL3 must be understood (37). Besides being in a position to protect themselves from drying, *H. contortus* iL3s have also established genes that participate in this biological cycle (38). Two versions of the major gene transcript were called Hc-ubq



and Hc-gst based on their homologous ubiquitin in *C. elegans* and glutathione-S-transferase in *H. contortus*, respectively (37,38). The silencing in L3 of *H. contortus* of Hc-ubq or Hc-gst by RNAi decreased the rate of survival, leading to its resistance (37). The Hc38 silencing, first observed with the blot of Northern and strongly expressed in the intestinal microvilli by in situ position, was decreased by 50 and 48.6 percent, respectively by taking iL3 from *H. contortus* into dsRNA, which were utilized to invade sheep (37,38).

The L4 has short body, low metabolic rate, and crystal rod-like in the intestinal canal to safeguard them in the cold niches (39). Gene expression includes Hc-daf-22 (3-ketoacyl-CoA thiolase), Hc-maoc-1 (enoyl-CoA hydratase), and Hc-hsp-70 (heat-shock protein 70). Hc-daf-22 and Hc-maoc-1 have similarities in *C. elegans*, Ce-daf-22 and Ce-maoc-1, respectively, and may act in the growth of the nematode and its peroxisomal β -oxidation (40). The Hc-hsp-70 overexpression manages the reduction of hsp-1 in *C. elegans*, indicating similar activity to that of the Ce-hsp-1 that helps of parasitic survival and invasion. Hc-fau, a human fau and *C. elegans* Crps30 analogs (ribosomal protein S30 coder), with a conserved domain of the S30 (expressed in the nucleus) and a diverged ubiquitin-like (UBiL) protein domain (cytoplasm-located expression) (41). They act in the egg-laying process and the life-span range of the *C. elegans*, indicating potential activities in L4 diapause based regulation in *H. contortus* (42).

Vaccination

Increasing consumption of anthelmintics for the care and mitigation of human and animal parasitic nematode diseases has created significant problems with anthelmintic and pharmaceutical

resistance around the world (43). The vaccine is an effective method to manage parasitic nematodes, like *H. contortus* (44). Remarkable improvement in the identification of several possible antigens from *H. contortus* over the past two decades has been done in order to enhance influential immune thresholds in the immune system (43,44). Here, recombinant subunit vaccine and DNA vaccination have been synthesized the molecular properties and defensive potency of the key-antigens (44).

Recombinant gene subunit-based vaccines

In terms of the production of consumer vaccines, special attention has been given to gene recombination, an essential biological innovation (14). Partial safeguards with *H. contortus* recombinant antigens has been granted for immunization (45). A 110kDa central membrane glycoprotein complex was the strongest described nominee for the *H. contortus* antigen named H11 (14,45). The immunogen (> 90% decrease in fecal egg counts (FECs), > 75% lowering of the weight on a worm) (45). Consequently, some scientists anticipate the recombinant vaccine for potential large-scale development to be produced to promote the realistic usage of the hosts for immune defense (14). *Escherichia coli* had H11-1 and H11-2 variant expression, which had slightly higher aminopeptidase activity than H11-1 with enzyme activity (14,45). It has been inoculated twice with H11-1, H11-2 and H11-1 and H11-2 mixed with phosphate buffered saline. Partial protection has been afforded through immunization with H11-1 and H11-2 blends (29% reduction in FECs and 18% lowering of worm load) (14,45). The three (H11-1, -2, and -3) fragments of H11 gene have also been inserted in the vector for yeast expression and the lithium chloride



process used to convert recombinant *Pichia Pastoris* X-33. RT-PCR identified transcriptions and SDS-PAGE, and Western blot verified glycosylation of the proteins. Even so, the recombinant molecule containing three isoforms of H11 did not show protection (14,45).

As a medium, *Caenorhabditis elegans* were also attempted to convey H11 to boost safety apart from *E. coli* and *P. pastoris*. (46) In this experiment, a flanking area of 1517bp 5' and part of the first H11 exon. *Caenorhabditis elegans* are subcloned into the upstream region of the pPD95,77 vector, the *contortus* and homologous genes of the *C. elegans* were respectively sub-clones (47). Microinjected into the *C. elegans* gonads, respectively. The findings revealed numerous trends of transcriptional expression powered by free-living and blood-sucking nematodes from their promoters, demonstrating the heterogeneous supply of *C. elegans* (46). The trans-HPS recombinant has been developed (a 1710 bp isoform gene fragment of the H11 gene) (46). Vaccination from transgenic worms with crude Trans-HPS led to a decrease in the FEC's by 38% and a decrease in worm weight by 25% (46,47). However, *E. coli* presented that in the immunization studies sheep were not secured by a gene fragment from 670 bp to 1710 bp isoform H11. In addition to possible antigen H11, the recombinant galectin Hco-gal-m / f (obtained respectively from male and female nematodes) is expressed from *H. contortus* in *E. coli* Vaccinations with 200 µg protein reduced 48-46 percent fecal egg output, with Freund's adjuvant. Vaccination with a mixture of Hco-gal-m / f recombinant proteins has a function to play in the defense of goats (46,47).

DNA-based vaccines

DNA vaccination is a bioengineered DNA defense strategy that allows cells to actively produce antigens and vaccinated animals to develop safe immune responses

to diseases (48). In contrast with traditional vaccines, DNA vaccines have several benefits, including a broader variety of specific immune responses (48). This is a new approach to managing infectious parasites. For the illustration, direct injection of antigen-specific immune-based plasmid codes for exogenous antigenic release (48). Several observations of partial safety in goats after DNA vaccination were reported and identified. Immunization of goats between 8 and 10 months of age, with HC29, DNA encoder (48). Different antibodies and partially immune defense (36% declines in FECs and worm burdens), comparison to goats who obtained just PBS, have been caused by the GPX (*contortus* glutathione peroxidase) (48). Vaccination with DNA Vaccines, comprising three fragments, encoding sections for H11-1 and caprine interleukin 2 (IL- 2), led to a significant reduction of the fecal output, the abomasal worm and lymphocytes by 57 and 47 percent of specific serum immunoglobulins G (IgG), IgA, CD4 + T lymphocytes and CD8 + T lymphocytes (48).

In *H. contortus*, Glyceraldehyde-3-phosphate dehydrogenase, as a protection against experimental infections of *H. contortus* in goats was evaluated with DNA vaccine (49). The analysis indicates that the vaccinated adds essential qualification peripheral and local mucosal immune responses and supported the development of lymphocytes of CD4+T and B but supplied only limited protection in comparison with control groups (35% decrease in FECs, and 38% decrease in the worm weight) (49). DNA vaccine may be much more secure (46% decrease in FECs and 51% decrease in worm burden) against the resulting disease in the goats through vaccination of disorganized muscle family member (Dim-1) (49).

The processes recognition of the immune control in order to identify useful vaccine antigens. Several experiments



were performed during a parasite invasion to examine the function of immune suppression by studying the interplay of parasite and host cell molecules (50). To achieve so, venipuncture has isolated the target peripheral blood mononuclear cells (PBMCs) of goat blood. PBMC is the mixture of activated cell subpopulations, primarily lymphocytes, monocytes and dendritic cells (T cells, B cells and NK cells) (50). The process for communicating host cells with several essential parasite molecules is crucial for any subpopulation. For instance. The Hco-gal-m / f (rHco-gal-m / f) recombinant *Contortus* Galectin peptides are grown using PBMC of caprines and investigated in the PBMC the impact of rHco-gal-m / f on the induction of apoptosis (50). The pathways underlying immunomodulation caused by rHco-gal-m / f were also studied together in a proteomic and transcriptomic way (50). PBMC. The results indicate that rHco-gal-m/f can bind to the surface of the PBMC and serve as an inflammatory reaction to ease *H. contortus* immune escape (50).

Based on the analysis, two H-co and F binding partners, Transmembrane Protein 147 (TMEM147), and Transmembrane Protein 63A (TMEM63A) were also discovered through further yeast two-hybrid testing (51). The communication between galectin and TMEM147 mainly mediates cell division, death of cells, and the transcription of cytokine in the PBMC (51). Along with TMEM63A, this membranous protein participates in the control of galectin and the development of the PBMC in the phagocytosis and nitric oxides. Even so, in the migratory galectin control and IFN- β transcript of goat PBMC, TMEM63A performs a significant role than TMEM147 (51). These research results provide a unique view for the clarification by nematodes and parasite-

host interaction of the processes involved in immune avoidance (51).

As a vaccine nominee, *H. contortus* excretory and secretory items include different proteins that may activate or suppress the host's immune system and are implicated in the pathophysiology of worms (52). Work suggested that the PBMC in vitro is suppressed by *H. contortus* excretory and secretory items. The items were used to inhibit IL-4, IFN- α supply, boost cytokine suppressive IL-10, strengthen inflammatory modulator IL-17, and repress the manufacturing of the nitric oxide (52). Protein engagement with the PBMC in vivo using liquid chromatography/tandem mass spectrometry from these materials in various developmental stages showed a total of 407 particles were detected interacting with the PBMC, 14-3-3 protein in all parasites as PBMC-interacting proteins (52). IL-4 output declined, and the PBMC in vitro proliferation repressed by the 14-3-3 isoform 2 (rHcftt-2) (52).

In addition, an important immunogenic component was also observed in 24 kDa *H. contortus* excretory / secretory proteins (53). The immune interactions between HcES-24 and Goat PBMC recombinant protein have shown an improvement in IL 4, IL-10, IL-17, and cell migration. Nonetheless, PBMC multiplication and NO development were greatly inhibited by the relationship (53). The results showed that the rHcES-24 had significant regulatory impacts on the PBMC (53).

Conclusion

The use of molecular-based approaches is critical in identifying the resistance type and its mechanism for the best utilization of therapeutic drugs against *Haemonchus contortus*.

References

1. Jones JT, Haegeman A, Danchin EGJ, Gaur HS, Helder J, Jones MGK, et al. Top 10 plant-parasitic nematodes in molecular plant pathology. *Mol Plant Pathol*. 2013;14(9):946-

61. <https://doi.org/10.1111/mpp.12057>
2. O'Connor LJ, Walkden-Brown SW, Kahn LP. Ecology of the free-living stages of major trichostrongylid parasites of sheep. *Vet Parasitol.* 2006;142(1-2):1-15. <https://doi.org/10.1016/j.vetpar.2006.08.035>
3. Blaxter M, Koutsovoulos G. The evolution of parasitism in Nematoda. *Parasitology.* 2015;142(Suppl 1):S26-39. <https://doi.org/10.1017/S0031182014000791>
4. Bundy DAP, Appleby LJ, Bradley M, Croke K, Hollingsworth TD, Pullan R, et al. 100 Years of Mass Deworming Programmes: A Policy Perspective From the World Bank's Disease Control Priorities Analyses. *Adv Parasitol.* 2018;127-54. <https://doi.org/10.1016/bs.apar.2018.03.005>
5. McKellar QA, Jackson F. Veterinary anthelmintics: Old and new. *Trends Parasitol.* 2004;20(10):456-61. <https://doi.org/10.1016/j.pt.2004.08.002>
6. Kaplan RM, Vidyashankar AN. An inconvenient truth: Global worming and anthelmintic resistance. *Vet Parasitol.* 2012;186(1-2):70-8. <https://doi.org/10.1016/j.vetpar.2011.11.048>
7. Hay SI, Abajobir AA, Abate KH, Abbafati C, Abbas KM, Abd-Allah F, et al. Global, regional, and national disability-adjusted life-years (DALYs) for 333 diseases and injuries and healthy life expectancy (HALE) for 195 countries and territories, 1990-2016: A systematic analysis for the Global Burden of Disease Study 2016. *Lancet.* 2017;390(10100):1260-344. [https://doi.org/10.1016/S0140-6736\(17\)32130-X](https://doi.org/10.1016/S0140-6736(17)32130-X)
8. Hewitson JP, Maizels RM. Vaccination against helminth parasite infections. *Expert Rev Vaccines.* 2014;13(4):473-87. <https://doi.org/10.1586/14760584.2014.893195>
9. Sallé G, Laing R, Cotton JA, Maitland K, Martinelli A, Holroyd N, et al. Transcriptomic profiling of nematode parasites surviving vaccine exposure. *Int J Parasitol.* 2018;48(5):395-402. <https://doi.org/10.1016/j.ijpara.2018.01.004>
10. Laing R, Kikuchi T, Martinelli A, Tsai IJ, Beech RN, Redman E, et al. The genome and transcriptome of *Haemonchus contortus*, a key model parasite for drug and vaccine discovery. *Genome Biol.* 2013;14(8). <https://doi.org/10.1186/gb-2013-14-8-r88>
11. Doyle SR, Laing R, Bartley DJ, Britton C, Chaudhry U, Gilleard JS, et al. A Genome Resequencing-Based Genetic Map Reveals the Recombination Landscape of an Outbred Parasitic Nematode in the Presence of Polyploidy and Polyandry. *Genome Biol Evol.* 2018;10(2):396-409. <https://doi.org/10.1093/gbe/evx269>
12. Gilleard JS. *Haemonchus contortus* as a paradigm and model to study anthelmintic drug resistance. *Parasitology.* 2013;140(12):1506-22. <https://doi.org/10.1017/S0031182013001145>
13. Coghlan A, Tyagi R, Cotton JA, Holroyd N, Rosa BA, Tsai IJ, et al. Comparative genomics of the major parasitic worms. *Nat Genet.* 2019;51(1):163-74. <https://doi.org/10.1038/s41588-018-0262-1>
14. Wang C, Li F, Zhang Z, Yang X, Ahmad AA, Li X, et al. Recent research progress in China on *Haemonchus contortus*. *Front Microbiol.* 2017;8:1509. <https://doi.org/10.3389/fmicb.2017.01509>
15. Kwa MSG, Veenstra JG, Roos MH. Benzimidazole resistance in *Haemonchus contortus* is correlated with a conserved mutation at amino acid 200 in β -tubulin isotype 1. *Mol Biochem Parasitol.* 1994;63(2):299-303. [https://doi.org/10.1016/0166-6851\(94\)90066-3](https://doi.org/10.1016/0166-6851(94)90066-3)
16. Prichard RK. Genetic variability following selection of *Haemonchus contortus* with anthelmintics. *Trends Parasitol.* 2001;17(9):445-53. [https://doi.org/10.1016/S1471-4922\(01\)01983-3](https://doi.org/10.1016/S1471-4922(01)01983-3)
17. Rufener L, Kaminsky R, Mäser P. In vitro selection of *Haemonchus contortus* for benzimidazole resistance reveals a mutation at amino acid 198 of β -tubulin. *Mol Biochem Parasitol.* 2009;168(1):120-2. <https://doi.org/10.1016/j.molbiopara.2009.07.002>
18. Kwa MSG, Veenstra JG, Van Dijk M, Roos MH. β -Tubulin genes from the parasitic nematode *Haemonchus contortus* modulate drug resistance in *Caenorhabditis*

- elegans. *J Mol Biol.* 1995;246(4):500-10.<https://doi.org/10.1006/jmbi.1994.0102>
19. Silvestre A, Cabaret J. Mutation in position 167 of isotype 1 β -tubulin gene of *Trichostrongylid* nematodes: Role in benzimidazole resistance? *Mol Biochem Parasitol.* 2002;120(2):297-300.[https://doi.org/10.1016/S0166-6851\(01\)00455-8](https://doi.org/10.1016/S0166-6851(01)00455-8)
20. Ghisi M, Kaminsky R, Mäser P. Phenotyping and genotyping of *Haemonchus contortus* isolates reveals a new putative candidate mutation for benzimidazole resistance in nematodes. *Vet Parasitol.* 2007;144(3-4):313-20.<https://doi.org/10.1016/j.vetpar.2006.10.003>
21. Zhang Z, Gasser RB, Yang X, Yin F, Zhao G, Bao M, et al. Two benzimidazole resistance-associated SNPs in the isotype-1 β -tubulin gene predominate in *Haemonchus contortus* populations from eight regions in China. *Int J Parasitol Drugs Drug Resist.* 2016;6(3):199-206.<https://doi.org/10.1016/j.ijpddr.2016.10.001>
22. Roeber F, Jex AR, Gasser RB. Advances in the diagnosis of key gastrointestinal nematode infections of livestock, with an emphasis on small ruminants. *Biotechnol Adv.* 2013;31(8):1135-52.<https://doi.org/10.1016/j.biotechadv.2013.01.008>
23. Kotze AC, Prichard RK. Anthelmintic Resistance in *Haemonchus contortus*. History, Mechanisms and Diagnosis. *Adv Parasitol.* 2016;93:397-428.<https://doi.org/10.1016/bs.apar.2016.02.012>
24. Redman E, Sargison N, Whitelaw F, Jackson F, Morrison A, Bartley DJ, et al. Introgression of ivermectin resistance genes into a susceptible *Haemonchus contortus* strain by multiple backcrossing. *PLoS Pathog.* 2012;8(2).<https://doi.org/10.1371/journal.ppat.1002534>
25. Luo X, Shi X, Yuan C, Ai M, Ge C, Hu M, et al. Genome-wide SNP analysis using 2b-RAD sequencing identifies the candidate genes putatively associated with resistance to ivermectin in *Haemonchus contortus*. *Parasites and Vectors.* 2017;10(1).<https://doi.org/10.1186/s13071-016-1959-6>
26. Lespine A, Ménez C, Bourguinat C, Prichard RK. P-glycoproteins and other multidrug resistance transporters in the pharmacology of anthelmintics: Prospects for reversing transport-dependent anthelmintic resistance. *Int J Parasitol Drugs Drug Resist.* 2012;2:58-75.<https://doi.org/10.1016/j.ijpddr.2011.10.001>
27. Khan S, Nisar A, Yuan J, Luo X, Dou X, Liu F, et al. A whole genome re-sequencing based GWA analysis reveals candidate genes associated with ivermectin resistance in *Haemonchus contortus*. *Genes (Basel).* 2020;11(4).<https://doi.org/10.3390/genes11040367>
28. Laing R, Maitland K, Lecová L, Skuce PJ, Tait A, Devaney E. Analysis of putative resistance gene loci in UK field populations of *Haemonchus contortus* after 6 years of macrocyclic lactone use. *Int J Parasitol.* 2016;46(10):621-30.<https://doi.org/10.1016/j.ijpara.2016.03.010>
29. Alam MBB, Omar AI, Faruque MO, Notter DR, Periasamy K, Mondal MMH, et al. Single nucleotide polymorphisms in candidate genes are significantly associated with resistance to *Haemonchus contortus* infection in goats. *J Anim Sci Biotechnol.* 2019;10(1):30.<https://doi.org/10.1186/s40104-019-0327-8>
30. Becker GM, Davenport KM, Burke JM, Lewis RM, Miller JE, Morgan JLM, et al. Genome-wide association study to identify genetic loci associated with gastrointestinal nematode resistance in Katahdin sheep. *Anim Genet.* 2020;51(2):330-5.<https://doi.org/10.1111/age.12895>
31. Luo X, Shi X, Yuan C, Ai M, Ge C, Hu M, et al. Genome-wide SNP analysis using 2b-RAD sequencing identifies the candidate genes putatively associated with resistance to ivermectin in *Haemonchus contortus*. *Parasit Vectors.* 2017;10(1):31.<https://doi.org/10.1186/s13071-016-1959-6>
32. Li FC, Gasser RB, Lok JB, Korhonen PK, He L, Di W Da, et al. Molecular characterization of the *Haemonchus contortus* phosphoinositide-dependent protein kinase-1 gene (Hc-pdk-1). *Parasites and Vectors.* 2016;9(1).<https://doi.org/10.1186/s13071-016-1351-6>

33. Li FC, Gasser RB, Lok JB, Korhonen PK, Wang YF, Yin FY, et al. Exploring the role of two interacting phosphoinositide 3-kinases of *Haemonchus contortus*. *Parasites and Vectors*. 2014;7(1).
34. Li F, Lok JB, Gasser RB, Korhonen PK, Sandeman MR, Shi D, et al. Hc-daf-2 encodes an insulin-like receptor kinase in the barber's pole worm, *Haemonchus contortus*, and restores partial dauer regulation. *Int J Parasitol*. 2014;44(7):485-96.<https://doi.org/10.1016/j.ijpara.2014.03.005>
35. Hu M, Lok JB, Ranjit N, Massey HC, Sternberg PW, Gasser RB. Structural and functional characterisation of the fork head transcription factor-encoding gene, Hc-daf-16, from the parasitic nematode *Haemonchus contortus* (Strongylida). *Int J Parasitol*. 2010;40(4):405-15.<https://doi.org/10.1016/j.ijpara.2009.09.005>
36. Mohandas N, Hu M, Stroehlein AJ, Young ND, Sternberg PW, Lok JB, et al. Reconstruction of the insulin-like signalling pathway of *Haemonchus contortus*. *Parasites and Vectors*. 2016;9(1):64.<https://doi.org/10.1186/s13071-016-1341-8>
37. Hartman D, Donald DR, Nikolaou S, Savin KW, Hasse D, Presidente PJA, et al. Analysis of developmentally regulated genes of the parasite *Haemonchus contortus*. *Int J Parasitol*. 2001;31(11):1236-45.[https://doi.org/10.1016/S0020-7519\(01\)00248-X](https://doi.org/10.1016/S0020-7519(01)00248-X)
38. Yang Y, Ma Y, Chen X, Guo X, Yan B, Du A. Screening and analysis of Hc-ubq and Hc-gst related to desiccation survival of infective *Haemonchus contortus* larvae. *Vet Parasitol*. 2015;210(3-4):179-85.<https://doi.org/10.1016/j.vetpar.2015.03.020>
39. Guo X, Zhang H, Zheng X, Zhou Q, Yang Y, Chen X, et al. Structural and functional characterization of a novel gene, Hc-daf-22, from the strongylid nematode *Haemonchus contortus*. *Parasites and Vectors*. 2016;9(1).<https://doi.org/10.1186/s13071-016-1704-1>
40. Ding H, Shi H, Shi Y, Guo X, Zheng X, Chen X, et al. Characterization and function analysis of a novel gene, Hc-maoc-1, in the parasitic nematode *Haemonchus contortus*. *Parasites and Vectors*. 2017;10(1):1-14.<https://doi.org/10.1186/s13071-017-1991-1>
41. Zhang H, Zhou Q, Yang Y, Chen X, Yan B, Du A. Characterization of heat shock protein 70 gene from *Haemonchus contortus* and its expression and promoter analysis in *Caenorhabditis elegans*. *Parasitology*. 2013;140(6):683-94.<https://doi.org/10.1017/S0031182012002168>
42. Yan B, Guo X, Zhou Q, Yang Y, Chen X, Sun W, et al. Hc-fau, a novel gene regulating diapause in the nematode parasite *haemonchus contortus*. *Int J Parasitol*. 2014;44(11):775-86.<https://doi.org/10.1016/j.ijpara.2014.05.011>
43. Tak IR, Dar JS, Dar SA, Ganai BA, Chishti MZ, Ahmad F. A comparative analysis of various antigenic proteins found in *Haemonchus contortus*--a review. *Mol Biol (Mosk)*. 2015;49(6):883-90.<https://doi.org/10.1134/S0026893315060217>
44. Bassetto CC, Amarante AFT. Vaccination of sheep and cattle against haemonchosis. *J Helminthol*. 2015;89(5):517-25.<https://doi.org/10.1017/S0022149X15000279>
45. Newton SE, Munn EA. The development of vaccines against gastrointestinal nematode parasites, particularly *Haemonchus contortus*. *Parasitol Today*. 1999;15(3):116-22.[https://doi.org/10.1016/S0169-4758\(99\)01399-X](https://doi.org/10.1016/S0169-4758(99)01399-X)
46. Yanming S, Ruofeng Y, Muleke CI, Guangwei Z, Lixin X, Xiangrui L. Vaccination of goats with recombinant galectin antigen induces partial protection against *Haemonchus contortus* infection. *Parasite Immunol*. 2007;29(6):319-26.<https://doi.org/10.1111/j.1365-3024.2007.00949.x>
47. Zhou QJ, Yang Y, Guo XL, Duan LJ, Chen XQ, Yan BL, et al. Expression of *Caenorhabditis elegans*-expressed Trans-HPS, partial aminopeptidase H11 from *Haemonchus contortus*. *Exp Parasitol*. 2014;145(1):87-98.<https://doi.org/10.1016/j.exppara.2014.08.005>
48. Zhao GW, Yan RF, Muleke CI, Sun YM, Xu LX, Li XR. Vaccination of goats with

DNA vaccines encoding H11 and IL-2 induces partial protection against *Haemonchus contortus* infection. *Vet J.* 2012;191(1):94-100.<https://doi.org/10.1016/j.tvjl.2010.12.023>

49. Yan R, Wang J, Xu L, Song X, Li X. DNA vaccine encoding *Haemonchus contortus* Actin induces partial protection in Goats. *Acta Parasitol.* 2014;59(4):698-709.<https://doi.org/10.2478/s11686-014-0298-z>
50. Wang W, Yuan C, Wang S, Song XK, Xu LX, Yan RF, et al.

Transcriptional and proteomic analysis reveal recombinant galectins of *Haemonchus contortus* down-regulated functions of goat PBMC and modulation of several signaling cascades in vitro. *J Proteomics.* 2014;98:123-37.<https://doi.org/10.1016/j.jprot.2013.12.017>

51. Li Y, Yuan C, Wang L, Lu M, Wang Y, Wen Y, et al. Transmembrane protein 147 (TMEM147): Another partner protein of *Haemonchus contortus* galectin on the goat peripheral blood mononuclear cells (PBMC). *Parasites and Vectors.* 2016;9(1).<https://doi.org/10.1186/s13071-016-1640-0>
52. Gadahi JA, Ehsan M, Wang S, Zhang ZC, Wang Y, Yan RF, et al. Recombinant protein of *Haemonchus contortus* 14-3-3 isoform 2 (rHcftt-2) decreased the production of IL-4 and suppressed the proliferation of goat PBMCs in vitro. *Exp Parasitol.* 2016;171:57-66.<https://doi.org/10.1016/j.exppara.2016.10.014>
53. Gadahi JA, Li B, Ehsan M, Wang S, Zhang Z, Wang Y, et al. Recombinant *Haemonchus contortus* 24 kDa excretory/secretory protein (rHcES-24) modulate the immune functions of goat PBMCs in vitro. *Oncotarget.* 2016;7(51):83926-37.<https://doi.org/10.18632/oncotarget.13487>