# Possible role of C-reactive protein for detection of gram positive urinary tract infection in children.

#### Haider SH.Aubais (M.SC.) College of veterinary Babylon university

#### Abstract

Uropathogenic strains of gram positive bacteria from 90 child aging from 1 to 9 year were studied from first February /2008 to end June/2008 for their susceptibility profile .the various isolates were *Staphylococcus aureus* (62), *Streptococcus pneumoniae*(16) and *Bacillus* spp.(12) *S. aureus* was the commonest causative agent for UTI(68.8%).Males were more infected (70%)than females and UTI were more in the children in 4 to 6 years (40%) than other ages groups. Antibiotics susceptibility pattern of these isolates against 28 different antibiotics revealed that Norfloxacin ,Nitrofurantion and Amikacin were effective for treatment of UTI (75-100%). Amoxicillin, Azithromycin, Bacitracin, Nitrofurantion , Carpeniciln, Cefoxitin, Ceftizoxime, Cephalexin and methicillin were not found to be effective for treatment of UTI (0-20%). The concentration of CRP was found to be more than 24 mg/dL in most patients (74.4%) and this level refers to true bacterial infection while the value of WBC was > 15000 / mm in (86.6%) of patients and ESR was found to be >35mm/h in (85.5%) of patients. In this study we found that CRP concentration were neither completely sensitive nor specific for detecting UTI infection but addition of CRP testing to that of WBC and ESR ensures the detecting of UTI.

#### Introduction

Urinary tract infection (UTI) with its diverse clinical syndromes and affected host groups remains one of the most common but widely misunderstood and challenging diseases encountered in children in clinical practice [1]. Approximately 2%-8% of children have urinary tract infection (UTI), depending of the age and gender [2]. Despite the wide spread availability of antibiotics, UTI remains the most common bacterial infection in the human populations [3]. Since patterns of antibiotics resistance in a wide variety of pathogenic organisms may verity even over short periods and depend on site of isolation and on different environments, periodic evaluation of antibacterial activity is needed to update this information [4]. Measurement of C-reactive protein (CRP), an acute phase protein synthesized by hepatocytes, is valuable in distinguishing systemic bacterial infection [5]. Many studies have addressed the value of CRP in detection bacterial urinary tract infection in addition to white blood cell and absolute neutrophil and erythrocyte sedimentation rate (ESR) [4,6, 7]. The present study recorded cases of UTI in children and studied spectrum of gram positive bacterial isolates and their antibiotics susceptibility and investigated the value of CRP in urinary tract infection in compared with WBC and ESR to determined the diagnostic properties of quantitative CRP associated with clinically undetectable serious bacterial infection in children 1-9 vears of age.

# **Materials and Methods**

In this prospective study we studied 90 child (rectal temperature > 38  $^{\circ}$ C) their ages, ranging from 1-9 years who presented to the pediatric clinic of emergency department of Babylon maternity and children hospital from first May/2008 to end July/ 2008, all children were hospitalized.

Every child underwent a history and physical examination and a full evaluation for sepsis was performed, including total WBC count, ESR, CRP, blood culture, gram stain, urine culture and urine analysis (UA).

Blood and urine specimens were cultured using stander media and techniques.

All urine specimens were obtained by clean container with parent's assistance.

All urine analysis was performed in a certified clinical laboratory.

Specimens were analyzed with both standard UA and haemocytometer WBC counts, the centrifuged urine specimens were examined microscopically on Neubauer chamber by the same technician.

C-reactive protein test done by using latex kit (Humatex CRP, Germany) for qualitative and semi-quantitative determination.

A loop calibrated to deliver 0.01 ml from Brain heart infusion broth vials inoculated with the urine specimen and after 24 to 48 hour at 37°C and appeared turbid was used to inoculate plate's containing blood agar and nutrient agar.

All plates were incubated at 37°C and examined at 24-48 hours for colony count and bacterial identification.

For standard UA, pyuria was defined as at least 5 WBCs/ hfp. For hemocytometer WBC count, pyuria was defined as al least 10 WBCs/ mL In this study we used 28 different antibiotics to estimate the sensitivity of UTI causative agents, using disk diffusion agar using Muller Hinton agar.

Quantitative urine cultures were performed in the microbiology laboratory. [9,10].

#### **Result and discussion**

The study group consisted of 90 children ranging in age from 1-9 years. Amongst the patients more than 68% showed *staphlococcos aureus* as the common isolate while *Streptococcus pneumoniae* and *Bacillus* spp was found in 17.7% and 13.3% of patients respectively (Figure 1). In another study of urine tract infection *S. aureus* was found to be the commonest gram positive isolated organism followed with *Streptococcus* spp. [11]. *Staphylococcus saprophyticus* was found to be 70.1% from all isolates of UTI in children from kabala.[12].



Figure 1: Percentage of Etiological agents causes UTI in children

the distribution of UTI causes. UTI were more in children 4- 6 years than in other age groups and was represented 40% of all patients(Table1). Males were more infected (70%) than females and this result agreed with another study [11].While differ from another study which found that males were less infected(13.5%) than females(15.3%) [12]. This study showed significant differences of *S. auerus* with age and gender but showed non significant relation of *St. pneumoniae* and Bacillus spp with age and gender factors ( $P \le 0.01$ ).

unu ugo,						
Age group (Year)	Gram Positive UTI Causes					
	Staphylococcus aureus		Streptococcus pneumoniae		Bacillus spp.	
	Male	Female	Male	Female	Male	Female
<1	12	2	5	1	2	2
1-3	4	1	3		1	-
4-6	22	7	-	4	1	2
7-9	7	7	2	1	4	-
Total	45	17	10	6	8	4
	62(68.8%)		16(17.7%)		12(13.3%)	

 Table 1: Distribution and frequency of UTI causes isolates according to gender and age.

Antibiotics susceptibility pattern of study isolates showed that most isolates of *S. auerus* were sensitive against Norfloxacin, Oxacillin and Amikacin, 90%, 90% and 80% respectively. But all isolates of this bacterium were resistance against Amoxillin, Cefoxatin and Ofloxacin. All *St. pnuemoinae* isolates were sensitive agiants Norfloxacin(100%) but resistance(100%) against

Amoxicillin+Clavulanic acid (AMC), Bacitracin, Carpenicillin, Cefoxitin and Streptomycin . *Bacillus* spp isolates were sensitive against Amikacin (100%) Chloromphenicol (95%) and Nitrofuration(85%),Kanamycin (85%) but showed different degree of sensitivity toward other antibiotics (Figure 2). Cefoxatin in the present study was found to be no effective for UTI as all the uropathogenes showed high degree of resistance to it, and this result agreed with the other study which showed that. Norfloxacin was effective for treatment of UTI and third generation Cephalosporin was not found to be effective for treatment of UTI [4]. In another study Gentamycin was the drug of choice for most of strains causes UTI while high degree of resistance showed against Tetracycline and Norfloxacin [13].



AK \Amikacin, AX \Amoxicillin, AMC \Amoxicillin + Clavulanic acid, AZM \Azithromycin, , B \Bacitracin, PY \Carbenicillin, CDZ \Cefodizime, FOX \Cefoxitin, ZOX \Ceftizoxime, CL \Cephalexin, C \Chloromphenicol E \Erythromycin, CN\Gentamycin, K\Kanamycin, L\Lincomycin, ME\Methicillin, F\Nitrofurantoin, NOR\Norfloxacin, OFX \Ofloxacin, OX\Oxacillin, T\Oxytetracyclin, P\Penicillin G. RF \Rifamycin, SP\Spiramycin, S\Streptomycin, TE\Tetracycline, TOB\Tobramycin, TMP\Trimethoprim.

Figure 2: Antimicrobial potency and spectrum for 28 selected antimicrobial agents tested against most frequently occurring UTI pathogens

Figure 3 showed the value of diagnostic test in which the CRP level was more than 24 mg/dL in 67 patients (74.4%) of all patients' studies, and ESR was more than 35 mm/h in 77 patients (85.5%) while WBC was found to be more than 15000/ mL in 78 of patients(86.6) with positive blood culture.



Figure 3: The percentage of diagnostic tests in presence of a positive blood culture

This result in line with another study which found the value of CRP above 85 mg/L for UTI compared with the value of WBC which found more than 15000mm<sup>3</sup>/L [7]. In another study used WBC, ESR and CRP to distinguish a SBI in infant showed that non of these parameters alone commonly used by physicians is a reliable diagnostic tool to rule out SBI in infants with fever without source of infection and according to this study the value of WBC > 15000, ESR> 20 mm/h and CRP= 2 mg/dL [14]. The value of CRP in this study appears to be higher in most children and the other who has low level of CRP may be treated with antibiotic before the study. Because of level of CRP remain elevated with on going inflammation and tissue destruction, but with resolution

they decline rapidly because of relatively short half-life of 4-7 hours [6]. The value of CRP in this study appears to be higher in most children and the other who has low level of CRP may be treated with antibiotic before they were studied and this lead to decreased level of CRP in the sera of patients[15]. In this study the elevation of CRP concentrations is non completely sensitive for detecting infection in patients with UTI, but could be ensured the infection and this result agreed with another study which explained that CRP concentration were neither completely sensitive nor specific for detecting infection in patients

with bacteremia. [16].

# References

1-Drekonja, D.M.and Johnson, J.R. (2008). Urinary tract infection. ., Pri.Car., 35(2):345-367

2Shaw,K.N.;Gorelick,M.G.;McGowan,K.I.;McDaniel,Y.M.andSchwartz;J.S .(1998).Prevalence of urinary tract infection in febrile young children in the emergency department. PEDIAT.,102(2):123-128
3-Sharma,S. (1997). Current understanding of pathogenic mechanisms in UTIsAnn.Nat.Acad.Med.Sci., 33(1):31-38.

- 4-Gupta,V.;Yadav;A.and Joshi;R.M.(2002)..Indian antibiotics resistance pattern in uropathogens.J.Med.Microbiol., 20(2):96-98.
- 5-Stanley, T.; Barrett, D.and Salmon; C.E. (1991). Viral and bacterial infection in childhood: the value of C-reactive protein . NZ.Med.J., 104:138-139.
- 6-Daniel,J.;Isaacman,M.D.;Bonnie,L.andBurke;M.S.(2002).Utility of C-reactive protein for detection of occult bacterial infection in children. Arch.Pediat.Adolesc.Med., 156: 905-909.
- 7-Shaol,R.;Lahad,A.;Tamir,A.;Lanir,A.and Srugo;I.(2008).C-reactive protein as apridictor of true bacteremia in children. Med.Sci.Monit., 14(5):255-261).
- 8-Ashkenazi,S.;Even,T.S.;Samra,Z.and Dinari;G .(1991).Uropathogens of various populations and their antibiotics susceptibility Pediat.Infect.Dis., 10:742-746.
- 9-Vandepitte, J.; Engbaek, K.; Plot, P. and Henck; C.(1991). Basic

laboratory procedures in clinical bacteriology. WHO Geneva

Switzerland.

- 10-MacFaddin, J.F. (2000). Biochemical tests for identification of medical bacteria.. 3<sup>rd</sup>.ed. Lippincott Williams and Wilkins. Philadelphia.
- 11-Al-Nassiri,Q.N.A.T .(2002).Biology of cell wall deffective microbes from persistent pyuria and persistent hematuria patients.Athesis of PhD. College of since, Babylon university.
- 12-Tuama,M.A.(2006).Urinary tracact infectin under five years in children in pediatrics hospital of karballa Athesis of M.Sc.. College of medicine, Babylon university.
- 13Yengkokpam,C.;Ingudam,D.;yengkokpam,I.S.and Jha;B.K.(2007).Antibiotic susceptibility pattern of urinary tract isolates in Imphal(Manipur) ,Indian.Nepal.Med.Col. J .,9(3):170-172.
- 14—Cuelo,G.C.A.; Tamez,G.L.and Valde;C.J.(2008).Total white blood cell ,erythrosedimentation rate and C-reactive protein for detection of serious bacterial infections in 0-to 90 days-old infants with fever without asource. .Ann.Ped.(Barc).,68(2):103-109.

- 15-Ehl,S.;Gering,B.;Bartmenn,P.;HOGL,j.and Pihland;F.(1997).C-reactive protein is auseful marker for guiding duration of antibiotics therapy in suspected neonatal bacterial infection.Pediat.,(2):216-222.
- 16-McCabe, R.E. and Remington, J.S. (1984). C-reactive protein in patients with bacteremia J.Clin. Microbiol., 20:317-319.

الدور المحتمل لبروتين اعادة التنشيط-C في التحري عن التهاب المجاري البولية المتسببة بالبكتيريا الموجبة لصبغة غرام عند الاطفال.

# حيدر شخير عبيس جامعة بابل/كلية الطب البيطري

تم دراسة العزلات الموجبة لصبغة غرام والمسببة لالتهاب المجاري البولية عند 90 طفل تراوحت اعمار هم بين 1-9 سنة للفترة من اول شهر شباط / 2008 لى نهاية حزيران/ 2008 وتم دراسة حساسيتها تجاة المضادات الحيوية. تم الحصول على 62 عزلة من بكتيريا Bacillus sureus ووجد ان Staphylococcus aureus و 12 عزلة من Staphylococcus aureus و 12 عزلة من بكتيريا Bacillus spp ووجد ان Staphylococcus aureus الكثر شيوعا في احداث التهاب لمجاري البولية وبنسبة 8.80% . ووجد ان الذكور اكثر عرضة للاصابة من الاناث بمعدل %7.20 اظهرت دراسة حساسية 28 مضاد حيوي تجاة العزلات البكتيرية بان المضادات الاتاث بمعدل %70.20 اظهرت دراسة حساسية 28 مضاد حيوي تجاة العزلات البكتيرية بان المضادات الولية وينسبة 75-100 و كانت المضادات الحيوية معاند في علاج التهاب المجاري البولية وينسبة 100.200 وكانت المضادات الحيوية معاد في علاج التهاب المجاري دوfoxitin وarpenicillin ومناد التهاب المجاري البولية المحادي البولية وينسبة 100.200 وكانت المضادات الحيوية معالج في علاج التهاب المجاري البولية المحادي دومناد المعاد معاد معاد معاد معادي المعاد المحادي وي تحاد التعربية عند معاد في علاج التهاب المجاري البولية وينسبة 25-000 % . كما وجد ان مستوى بروتين اعادة التنشيط 2 في المحاري الولية المرضي كان اكبر من 2001 هو معاد التهاب المجاري البولية التهاب المجاري البولية التراوح البولية وينسبة 25-2001 % وكانت المضادات الحيوية معالة في علاج التهاب المجاري البولية التراوح المرضى كان اكبر من 2002 % . كما وجد ان مستوى بروتين اعادة التنشيط 2 في الماد الال الم نسبة الحساسية بين 15000 % . كما وجد ان مستوى بروتين اعادة التشيط 2 في مصال 14.7% من المرضى كان اكبر من 2500 % من المرضى ومعدل ترسيب كريات الدم الحمر اكبر من 35 المرضى كان اكبر من 15000 / 1500 هو معاد المرامي ومعدل ترسيب كريات الدم الحمر اكبر من 35 المرضى كان اكبر من 2001 / 10 في 20.80 % من المرضى ومعدل ترسيب كريات الدم الحمر اكبر من 35 الموضى كان اكبر من 2001 / 10 في 20.80 % من المرامي وحد ان اجراء فحص بروتين اعادة التنشيط - 2 لم يكن حساس او متخصص في تشخيص التهاب المجاري البولية في حين ان اضافة فحص - 2000 هو التها.