

Effect of Essential Oils Extracted from the Peels of Two Species of *Citrus* on Some Fungi

Estabraq A. Mahmoud

Department of Biology, College of Education/Ibn-Al-Haitham, University of Baghdad

Abstract

This study investigated the effect of essential oils extracted from peel of *Citrus limon* and *Citrus reticulata* on two species of fungi: *Penicillium expansum* and *Fusarium proliferatum* and also effect of two fungicides: Hymexazol and Benomyl against this fungi. Results showed that the essential oils of *C. limon* inhibited the radial growth of *P. expansum* and *F. proliferatum* at concentration 4.5 and 5%, respectively. However, the essential oil of *C. reticulata* inhibited this growth at concentration 5.5 and 6%, respectively. Moreover, the two fungicides inhibited radial growth of this fungi. In conclusion, there is a positive relationship between the increasing of concentration and the percentage of inhibiting of radial growth of fungi.

Keywords: Essential oils, citrus peels, fungi.

تأثير الزيوت الطيارة المستخلصة من قشور نوعين من الحمضيات في بعض انواع الفطريات

استبرق عز الدين محمود

قسم علوم الحياة، كلية التربية، ابن الهيثم، جامعة بغداد

خلاصة

هدفت الدراسة الحالية فحص تأثير الزيوت الطيارة المستخلصة من قشور نبات الليمون *Citrus limon* واللائكي *Citrus reticulata* في نوعين من الفطريات (*Penicillium expansum* و *Fusarium proliferatum*) وتأثير اثنين من المبيدات الفطرية (Hymexazol و Benomyl) تجاه نفس الفطريات. أظهرت النتائج وجود تثبيط كامل للنمو الشعاعي للفطرين *P. expansum* و *F. Proliferatum* عند التركيز 4.5 و 5% على التوالي للزيت الطيار لنبات الليمون 5.5 و 6% على التوالي للزيت الطيار لللائكي. كذلك تثبيط النمو الشعاعي للفطرين من قبل اثنين من المبيدات الفطرية، في حين أظهرت التركيزات الأدنى زيادة في معدل التثبيط مع زيادة التركيز لكل المواد المستعملة بالدراسة سواء كانت زيت طيار او مبيدات.

كلمات مفتاحية: زيوت أساسية، قشور الحمضيات، الفطريات.

Introduction

Mandarin and lemon belong to family Rutaceae and order Geraniales which return to sub-class Archichlamyae (1). The origin of lemon (*Citrus limon*) is still uncertain. It must have originated somewhere in southeastern Asia while China is one of the native homes of the mandarin (*Citrus reticulata*) (2). Essential oils are complex natural mixtures of volatile secondary metabolites, isolated from plants by hydro- or steam distillation and by expression (3). The main constituents of essential oils are monoterpenes and sesquiterpenes including carbohydrates, alcohols, ethers, aldehydes and ketones which are responsible for the fragrant and biological properties of aromatic and medicinal plants (4). Citrus essential oils are present in fruit flavedo in high quantities. Peels consist of the epidermis covering the exocarp consisting of irregular parenchymatous cells, which are completely enclosing numerous glands or oil sacs. Citrus essential oils are a mixture of volatile compounds and which mainly consisted of monoterpene hydrocarbons. The terpene fraction can constitute from 50 to more than 95% of the oil; however, it makes little contribution to the flavor and fragrance of the oil (5). Limonenes are found in the essential oil of various citrus leaves and fruit peels and have inhibited properties of both insects and fungi (6). Many studies investigated the essential oils of *Citrus limon* and *Citrus reticulata* against the growth of fungi (7,8,9,10,11,12,13,14,15,16,17).

The high quantity of citrus peel found as industrial waste and citrus industry presents a potential pollution problem, which would be reduced if the waste could be utilized as food for animals (18). Thus the aim of this study uses this waste as fungicide against some fungi and evaluates the effect of fungicides on the growth of fungi and this study is novel in Iraq.

Material and Methods

Plant collected: The samples of *Citrus lemon* and *Citrus reticulata* collected in March 2012 from local market after the fruits had been washed, they were cut into six equal portions and the flesh was removed and used directly without drying (15).

Fungi: *Penicillium expansum* isolate was obtained from University of Baghdad, College of Science, Biology Department, while fungi isolate *Fusarium proliferatum* was obtained from fungi Laboratory postgraduate in University of Baghdad, College of Education Pure Science, Ibn al-Haitham, Biology Department and re-diagnosed and confirmed for them.

Fungicides: Pesticides Hymexazol and Benomyl (benzimidazole) have been obtained from the local markets and prepared according to the described method.

Extraction of essential oils by steam distillation:

The extraction of essential oils have been extracted according to (12) protocol briefly, a 100 gm from fresh peels of plant and placed in the round bottom flask and filled with 1000 ml of distilled water then distillation apparatus was connected to the flask and take at 60 °C. Distillation was continued until there was no more difference in successive readings of the oil volume. The yield of essential oil (%) was calculated as follows (16):

$$\text{Yield of essential oil}\% = \frac{\text{Volume of essential oil (mL)}}{\text{Fruit peel sample (g)}} \times 100$$

Oils collected in distilled dark bottles until used.

Test of sensitive of fungi against essential oils

Each of Petri dish contained potato dextrose agar (PDA) and prepared the concentration below:

C. limon against *P. expansum* (1, 2, 3, 4, 4.5) (v/v)%

C. limon against *F. proliferatum* (1, 2, 3, 4, 4.5, 5) (v/v)%

C. reticulata against *P. expansum* (1, 2, 3, 4, 5, 5.5) (v/v)%

C. reticulata against *F. proliferatum* (1, 2, 3, 4, 5, 5.5, 6) (v/v)%

Dishes were inoculated with the fungus by cutting a 4 mm-diameter disc from pure cultures of *P. expansum* and *F. proliferatum* growing on PDA using a cork borer. This was done for each of concentration as well as for control (without essential oil). The cultures were incubated at 27 °C in incubators for 6 days. Radial growth rate and inhibition ratio as estimated by measuring the maximum diameter of colonies was measured after 3-6 day and the ratio diameter/time was calculated by used the formula as shown below:

$$\% \text{ Growth inhibition} = \frac{DC - DT}{DC} \times 100 \quad (2)$$

DC = The diameter of mold colony from control plate.

DT = The diameter of the mold colony growth in experiment plate which contains the essential oil (9).

Test of sensitive of fungi against fungicide

By using the method of (18), the radial growth rate and inhibition ratio was measured according to formula (2). The concentrations of fungicides were prepared: Hymexazol against *P. expansum* and *F. proliferatum* (50, 100, 150, 200, 225) (w/v)% from the stock solution 36%.

Benomyl against *P. expansum* and *F. proliferatum* (50, 100, 150, 200, 225) (w/v)% from the stock solution 50%.

Statistical Analysis

All determinations were made in triplicates and the data is reported as mean \pm SE for (n = 3). Analysis of Variance (ANOVA) method was used for statistical analysis and at probability levels (0.05, 0.01, 0.001) for the purpose of evaluating the differences in the results of transactions in terms of being significant (influence of material) or not significant differences (as a result of laboratory errors) (19).

Results and Discussion

-Essential oils: The results had been show that for every 100 grams of *C. limon* and *C. reticulata* peels contain (0.12, 0.08) ml [(0.12%, 0.08%) according to formula (1)] of essential oils respectively, this result is consistent with (20) they found that every 100 grams of *C. reticulata* peels contain 0.089 essential oils extracted by steam distillation.

-Sensitive of fungi against essential oils: The results showed that the treatment of fungi, *P. expansum* and *F. proliferatum* with essential oil extracted from the peel of fruits, *C. lemon* and *C. reticulata* have antifungal activity and gradually influenced the growth of these two isolates by increasing low concentration killer below the level of probability (0.001, 0.05, 0.01). The sensitivity of fungi were differed from two essential oil (Figure 1, 2, 3 and 4), where the complete inhibitory effect of essential oil of *C. lemon* on *P. expansum* and *F. proliferatum* were (4 and 4.5)% respectively, and for *C. reticulata* were (5.5 and 6)% . This results are agreed with (7) and (11) which found the essential oils of *C. lemon* and *C. reticulata* have effective toward *Penicillium* sp. and agreement with (8) that found there was an inhibitory of the essential oil *C. lemon* on *Fusarium* spp. and *Penicillium* spp. grown while the recent study was disagreed with studies (9) and (13) those found concentration 10% from *C. lemon* inhibited growth of *Fusarium* spp., moreover, the activity of fungicidal against the myceliae growth of *Fusarium* spp. was observed at a concentration of 4% of lemon essential oils in study (10), which is consistent nearly with what was found in this study (figure 2). Study (14) found that the essential oil of lemon extract has inhibition effect toward four food-borne fungal strains including *Fusarium* spp. The results showed also that the essential oils of *C. lemon* has a significant effect on *Penicillium* sp. growth than *Fusarium* spp. at below the level of probability (0.001, 0.05, 0.01) (Figure 1,2). These findings are agreement with (11) that concluded hydrodistilled-essential oils from

all citrus cultivars were strong antifungal agents toward *Penicillium* spp. While the essential oil of *C. lemon* showed inhibitory action more than essential oil of *C. reticulata* at below the level of probability (0.001, 0.05, 0.01) (figure 1,2,3 and 4), this agreement with (7) which is found that the essential oil of *C. lemon* have more inhibitory action against *Penicillium* spp. than *C. reticulata*. The essential oil of *C. reticulata* significantly inhibited the growth of *Penicillium* spp. more than *Fusarium* spp. at below the level of probability (0.001, 0.05, 0.01) (figure 3,4), this found agreed with (14) which reached that *Penicillium* spp. more affected than other fungi including *Fusarium* spp. and the essential oil extracted from *C. reticulata* has inhibition activity and agreement with (17) that found the essential oil of *C. reticulata* has effective inhibition toward *F. proliferatum*, but less than the impact of essential oil of *Citrus aurantifolia*.

This inhibitory action can be explained by the demonstration of the action of essential oils on the wall of fungi whose structure and function are altered and the transport of nutrients is modified (9). Essential oils extracted from aromatic plants such as *C. lemon* and *C. reticulata* that possess antimicrobial activity, while diameters of treated colonies were smaller than control group, depending on the concentration of oil (14). Extracts of citrus plants contain antifungal compounds that can be used as alternative to synthetic fungicides including fumigants and contact pesticides. The prospect of using citrus for development of natural fungicides is appealing and acceptable because citrus peels are readily available, environmentally safe, and less risky for developing resistance in pests, less hazardous to non target organisms and pest resurgence, less adverse effect on plant growth, less harmful to seed viability and quality and above all less expensive. Based on these findings, citrus plant extracts are viable and can be possible alternative to synthetic pesticides for control of fungal diseases [9].

In a number of citrus species, the bitterness causative factors are limonoids, limonine being one of the potential antipest compounds known. A few other citrus limonoids including nomiline, nomilinic acid, ichangin and obacunonic acid are also bitter. Among these, limonine and nomiline are known to deter feeding in lepidopterans and coleopterans with variable efficacies. It appears that furan and epoxide groups play an important role in the activity of these compounds (21). Tangerine oil was obtained from the peel of *Citrus reticulata* is pleasant in taste and rich in aroma and is mainly used in food and beverages as flavoring agent.

As the essential oil is rich in a wide variety of secondary metabolites, such as tannins, terpenoids, alkaloids and flavonoids, that are found to have effective as antimicrobial properties. The monoterpenes affect the structural and functional properties of lipid fraction of the plasma membranes of bacteria and yeasts, causing leakage of intercellular material and exit of critical molecules and ions leading to death of microbes. Terpenoids affect respiratory enzymes inhibiting microbial oxygen uptake and oxidative phosphorylation (12).

-sensitive of fungi against fungicide

The results had been shown in figure 5, 6, 7 and 8 suggested that there is a significant effectiveness of the fungicides used against two fungal isolates (*P. expansum* and *F. proliferatum*) at below the level of probability (0.001, 0.05, 0.01) and fungicide hymexazol more effective on two fungus than the fungicide Benomyl (Figure 5, 6, 7, 8) while the fungus *P. expansum* was more influenced by both fungicides (Figure 5,7) at below the level of probability (0.001, 0.05, 0.01). Benomyl used against *Fusarium* sp., when infected guava wilt disease where added at concentration 0.2% (22) and inhibited the growth of *Fusarium* sp. by 83.64% (9), as also hymexazol used against *Fusarium* sp. (23). Study (24) was concluded that hymexazol at

the dose recommended by the manufacturer significantly reduced disease incidence for a period of 2 months after transplanting. However Benomyl is unstable and easily decomposed to methyl 2-benzimidazole carbamate in aqueous solutions (25).




Figure (1): Inhibition% for different concentrations of essential oil of peel fruits of *C. limon* in surface growth of the fungus *P. expansum*.

Less significant difference (LSD) at 0.05 level for days 1.06 for concentrations 1.05 and interference 1.74.




Figure (2): Inhibition% for different concentrations of essential oil of peel fruits of *C. limon* in surface growth of the fungus *F. proliferatum*.

Less significant difference (LSD) at 0.05 level for days 1.33 for concentrations 1.00 and interference 1.72.




Figure (3): Inhibition% for different concentrations of essential oil of peel fruits of *C. reticulata* in surface growth of the fungus *F. expansum*.
 Less significant difference (LSD) at 0.05 level for days 1.01 for concentrations 0.76 and interference 1.31.




Figure (3): Inhibition% for different concentrations of essential oil of peel fruits of *C. reticulata* in surface growth of the fungus *F. proliferatum*. Less significant difference (LSD) at 0.05 level for days 2.68 for concentrations 2.02 and interference 3.47.




Figure (5): Inhibition% for different concentrations of Hymexazol in surface growth of the fungus *P. expansum*.

Less significant difference (LSD) at 0.05 level for days 1.33 for concentrations 1.32 and interference 2.19.




Figure (6): Inhibition% for different concentrations of Hymexazol in surface growth of the fungus *F. proliferatum*.

Less significant difference (LSD) at 0.05 level for days 1.62 for concentrations 1.60 and interference 2.66.




Figure (7): Inhibition% for different concentrations of Benomyl in surface growth of the fungus *P. expansum*.

Less significant difference (LSD) at 0.05 level for days 1.99 for concentrations 1.97 and interference 3.26.




Figure (8): Inhibition% for different concentrations of Benomyl in surface growth of the fungus *F. proliferatum*.

Less significant difference (LSD) at 0.05 level for days 2.03 for concentrations 1.53 and interference 2.62.

References

1. Bhattacharyya, B. and Johri, M. M. (1998). Flowering plant taxonomy and phylogeny. Navosa publishing house, New Delhi:753 pp.
2. Nicolosi, E. (2007). Origin and taxonomy. In: Khan, I. A. (Ed.). Citrus Genetics, Breeding and Biotechnology. Biddles Ltd, Kings Lynn, UK: 370 pp.
3. Tripathi, P. and Shukla, A. K. (2010). Exploitation of botanicals in the management of phytopathogenic and storage fungi. In: Arya, A. and Perelló, A. E. (Eds.). Management of fungal plant pathogens. MPG books Group, UK: 388 pp.
4. Kalemba, D. and Kunicka, A. (2003). Antibacterial and antifungal properties of essential oils. Curr. Med. Chem., 10:831-829.
5. Chanthaphon1, S.; Chanthachum, S. and Hongpattarakere1, T. (2008). Antimicrobial activities of essential oils and crude extracts from tropical *Citrus* spp. against food-related microorganisms. Songklanakarin J. Sci. Technol., 30(Suppl.1): 125-131.
6. Koul, P.; Walia, S. and Dhaliwal, G. S. (2008). Essential Oils as Green Pesticides: Potential and Constraints. Biopestic. Internat., 4(1): 63-84.
7. Hall, D. J. and Fernandez, Y. J. (2004). *In vitro* evaluation of selected essential oils as fungicides against *Penicillium digitatum* sacc. Proc. Fla. State Hort. Soc, 117:377-379.
8. Leite de Souza, E.; Lima, E.; Freire K. R. and Paiva de Sousa, C. (2005). Inhibitory action of some essential oils and phytochemicals on the growth of various moulds isolated from foods. Braz. arch. biol. technol., 48(2): 245-250.
9. Okwu, D. E.; Awurum, A. N. and Okoronkwo, J. I. (2007). Phytochemical composition and *in vitro* antifungal activity screening of extract from citrus plants against *Fusarium oxysporum* okra plant *Hibiscus esculentus*). African Crop Conference Proceedings, 8: 1755-1785.
10. El-Mougy, N. S.; El-Gamal, N. G. and Abdel-Kader, M. M. (2007). Control of wilt and root rot incidence in *Phaseolus vulgaris* L. some plant volatile compounds. J. Plant Prot. Res., 47(3):255-265.
11. Viuda-Martos, M., Ruiz-Navajas, Y., Fernandez-Lopez, J. and Perez-Alvarez, J. (2008). Antifungal activity of lemon (*Citrus lemon* L.), mandarin (*Citrus reticulata* L.), grapefruit (*Citrus paradise* L.) and orange (*Citrus senesis* L.) essential oils. Food Control, 19(12):1130-1138.
12. Shahzad1, K.; Nawaz1, S.; Ahmad, R.; Akram, N. and Iqbal, Z. (2009). Evaluation of antimicrobial, antifungal and antioxidant activity of essential oil of *Citrus reticulata* fruit (Tangerine fruit peel). *Pharmacologyonline*, 3: 614-622.
13. Adejumo T. O. and Langenkämper, G. (2012). Evaluation of botanicals as biopesticides on the growth of *Fusarium verticillioides* causing rot diseases and fumonisin production of maize. J. Microbiol. Antimicrob., 4(1): 23-31.
14. Ferdeş, M. and Ungureanu, S. (2012). Antimicrobial activity of essential oils against four food-borne fungal strains. U.P.B. Sci. Bull., Series B, 74 (2): 87-98.
15. Kirbaşlar, F. G.; Tavman, A. T.; Dülger and Türker (2009). Antimicrobial activity of Turkish *Citrus* peel oils. Pak. J. Bot., 41(6): 3207-3212.
16. Oh, H. J.; Ahn, H. M.; Kim, S. S.; Yun, P. Y.; Gyeong Lyong Jeon, G. L.; Ko, Y. H. and Riu, K. Z. (2007). Composition and Antimicrobial Activities of Essential Oils in the Peel of Citrus Fruits. J. Appl. Biol. Chem. 50(3): 148-154.
17. Roy, S. D.; Bania, R.; Chakraborty, G.; Goswami, R.; Laila, R. and Ahmed, S. A. (2012). Pharmacognostic, phytochemical, physicochemical property and antimicrobial activity studies of lemon peel oil. J. Nat. Prod. Plant Resour., 2(3):431-435.
18. Solel, Z. and Pinkas, Y. (1984). A modified selective medium for detecting *Phytophthora cinnamomi* on avocado roots. Americ. Phytopathol. Soc., 74(4): 506-508.
19. Zar, J.H. (1999). Biostatistical 4th. Prentice hall upper saddle river, analysis. New jersey: 663 pp.

20. Ayoola¹, G. A.; Johnson¹, O. O. ; Adelowotan, T. ; Aibinu, I. E. ; Adenipekun, E.; Adepoju-Bello, A.A. ; Coker, H. A. B. and Odugbemi, T.O. (2008). Evaluation of the chemical constituents and the antimicrobial activity of the volatile oil of *Citrus reticulata* fruit (Tangerine fruit peel) from South West Nigeria. *Afr. J. Biotechnol.*, 7(13): 2227-2231.
21. Koul¹, O. and Walia, S. (2009). Comparing impacts of plant extracts and pure allelochemicals and implications for pest control. *Perspect. Agricult., Veterin. Sci., Nutr. Nat. Res.*,4(049): 1-30.
22. Sagar, B. V. and Varma, P. K. (2012). Practical manual of diseases of horticultural crops and their management. Acharya Ng Ranga Agricultural University, College of Agriculture, Rajendranagar-500030: 72 pp.
23. Doulias, K.; Vloutoglou, I.; Elena, K.; Paplomatas, E. J. and Lasvaris, D. (2000). Studies on pathogens causing damping-off of sugarbeet seedlings and on the effectiveness of fungicides as seed coats. *Phytopathol. Mediterr.*, 39: 313-338. (Abstract).
24. Pavlou, G.C. (2006). Preliminary study on hymexazol efficacy to control root and stem rot of cucumber caused by *Fusarium oxysporum* f. sp. *radicis-cucumerinum*. *Phytopathol. Mediterr.*, 45:163–200. (Abstract).
25. Abayasekara, C.; Ratnayake, S. and Adikaram, N.K.B. (1998). Resistance of banana fruit to fungal disease an overview. In: Johnson, G. I. ; Highly, E. and Joyce, D. C. (Eds). *Disease Resistance in fruit*, ACIAR Proceedings No. 80: 39-45 Canberra.