

Prevalence of Silent Bacteriuria in Patients with Diabetes Mellitus

Firas Elias Douri

ABSTRACT:

BACKGROUND:

Diabetes mellitus and its complications continue to carry a major health problem. There is evidence that diabetics are more prone to urinary tract infection.

AIM OF THE STUDY:

The study was done to assess the prevalence of significant bacteriuria in patients with no urinary symptom.

METHODS:

Midstream urine samples were collected from 200 diabetic patients. There were 102 females and 98 males and 80 urine samples from non diabetic persons as control were screened for bacteriuria.

RESULTS:

Out of 200 diabetic urine samples 18 patients [9%] had significant bacteriuria while 2 urine samples [2.5%] of non diabetic samples has significant bacteriuria. Three types of bacteria were isolated from urine samples, *Escherchia coli*, *Staphylococcus*, and *Proteus* and *Escherchia coli* was the high in number among the other genera.

CONCLUSION:

Analysis of the results showed significant bacteriuria in diabetic patients compared with non diabetic patients {p value < 0.05}.

KEY WORDS: bacteriuria , diabetes mellitus

INTRODUCTION:

Diabetes Mellitus is a common metabolic disorder and result in several abnormalities of the host defense system that might increase the risk of certain infections. Infection in diabetic patients is the most serious complication which continues to be a big health problem all over the world. Diabetic foot ulcer were the major cause of infection followed by pulmonary tuberculosis, Urinary tract infection, pneumonia and skin and subcutaneous infections^(1,2). There is a view that urinary tract infection is more common in diabetic patients⁽³⁾ but remain in debate because of varying study designs⁽⁴⁾.

According to the 1998 WHO criteria, diabetes mellitus was defines as fasting glucose concentration of at least 6.1 mmol/l (110 mg/dl) or a two-hours postprandial glucose concentration of at least 10.0 mmol/l (180 gm/dl) or use of glucose lowering medication (tablets or insulin)⁽⁵⁾.

The presence of pyuria and bacteriuria are two most important indicators of urinary tract infections⁽⁶⁾. Many urinary tract infections are asymptomatic. In contrast with men; the prevalence of asymptomatic bacteriuria is three to four times higher among women with diabetes and tends to have persistent or recurrent bacteriuria than those without this condition^(7,8,9).

The bladder urine of normal persons inhibits and kills bacteria. Polymorph nuclear leukocytes in the bladder wall also appear to play a role in killing bacteria.

Asymptomatic bacteriuria was defined as the presence of at least 10^5 colony forming units (cfu) /ml of the same single species plus pyuria in one culture of clean voided specimen of midstream urine sample from an individual without symptoms of UTI^(8,9). we defined contaminated urine as the presence of at least 3 different microorganisms in one urine sample, these specimen were excluded

A symptomatic bacteriuria also defined as $\geq 10^5$ / ml

Department of Medical Microbiology / AlKindy
College of Medicine / University of Baghdad

SILENT BACTERIURIA IN DIABETES

of one single species in the midstream urine of person without urinary symptoms {frequency, dysuria, haematuria, loin pain.....etc.}, when it occurs in infants, pregnant women, immunocompromized persons and in abnormal urinary tract investigations and treatment is required because of high risk of pyelonephritis in those patients. The recognition of the number and type of bacteria in the urine is very important diagnostic procedure. Quantitative estimation of the number of bacteria in urine specimens make it possible to distinguish contamination from true bacteriuria and bacterial colony count of $\geq 10^5$ /ml has been the criterion traditionally used for this purpose. However in symptomatic women with pyuria, colony counts of 10^2 - 10^4 *Escherichia coli*, *Proteus*, *Klebsiella* or *Staphylococcus saprophyticus* per milliliters of midstream urine usually indicate infection, not contamination and should not be disregarded. In certain circumstances (antibiotics, high urea concentration, high osmolarity, low PH) inhibit bacteria multiplication resulting in relatively low bacteria colony counts despite infection, for this reason antiseptic solution should not be used in washing the periurethral area before collection of urine specimen.

Rapid screening methods for detection of bacteriuria have been developed as alternative to standard culture methods, these methods have been developed for measuring bacterial concentration such as a new automated flow cytometry-based urine bacterium analyzer^(10,11) in addition to other rapid screening methods such as Yellow IRIS⁽¹²⁾ that provide results rapidly usually in 1-2 hours. These techniques exhibit sensitivity of 95%-98%. The sensitivity of these tests fall to 60%-80% when 10^2 - 10^4 colony forming units per ml is the standard of comparison.

Microscopy of urine from symptomatic patients can be of great diagnostic value. Microscopical bacteriuria, which is best assessed with Gram's stained and centrifuged urine is found in more than 90% of specimens from patients whose infections are associated with colony counts of at least 10^5 per ml and this finding is very specific. Bacteria can not usually be detected microscopically in infections with lower colony count (10^2 - 10^4 /ml). The detection of bacteria by urinary microscopy constitutes firm evidence of infection but the absence of microscopically detected bacteria does not exclude the diagnosis.

Leukocyte esterase dipstick method is less sensitive than microscopy in identifying pyuria^(13,14) but is useful alternative when microscopy is not feasible.

PATIENTS AND METHODS:

Participants

The study was conducted in diabetic clinic in AlKadhmyia hospital from June 2007 to November 2007. Midstream urine samples were collected from 200 diabetic patients and screened for asymptomatic bacteriuria, age 16-70 years attending the diabetic clinic, there were 102 females and 98 males, 122 having non insulin dependent diabetes mellitus (NIDDM) and 78 patients having insulin dependent diabetes mellitus (IDDM).

Midstream urine samples were collected from 80 non diabetic patients attending the medical outpatient clinic; they were 50 females and 30 males.

The criteria for inclusion in the study are:-

- 1-no urinary tract symptoms.
- 2-no antibiotic administration within previous two weeks
- 3-no pregnancy for females
- 4-no functional or structural abnormalities of the urinary tract.

Full history taking, good physical examination, ultrasonography and blood sugar reading have been done to fulfill the inclusion criteria.

Urine analysis

All urine samples were either transport to the microbiology laboratory for culture within 2 hours of collection or refrigerated. Urine culture was performed according to standard procedure(8): urine was screened for the presence of either more than 5 leukocytes or 10 microorganisms were seen in the slide then the urine was plated onto blood agar and MacConkey plates, all urine samples were plated using quantitative loop and the plates were incubated at 37°C aerobically, the results were read after 24 hours. Microorganisms were identified with API system (Analytic Profile Index system) [bioMerieux]. If the urine was considered contaminated the patient was asked to submit another sample.

RESULTS:

Out of 200 diabetic urine samples examined 18 [9%] had significant bacteriuria, while 2 urine samples out of 80 [2.5%] non diabetic urine samples had significant bacteriuria.

In both groups [the diabetic and the control group], females are more than males in having significant

SILENT BACTERIURIA IN DIABETES

bacteriuria [77.8% in diabetics and 100% in non diabetics].

Out of 18 positive isolates from diabetic urine samples 10 [55.5%] were from NIDDM group and 8 [44.5%] were from IDDM group, this difference was statistically not significant.

The results of microbial culture on blood agar and MacConkey agar and API system showed the prevalence of the isolates of Escherchia coli, Proteus and Staphylococcus aureus in both group, Escherchia coli was the most common isolates 12 [66.7%] in diabetic samples while in control group (non diabetic) was 2 [100%] samples.

Table 1: Samples of significant bacteriuria in diabetic and non diabetic patients P. value < 0.05 (statistically significant)

Urine culture	Diabetic samples	Non diabetic samples
Positive	18 [9%]	2 [2.5%]
Negative	182 [91%]	78 [97.5%]

Table 2: Distribution of bacterial isolates from urine samples of diabetic and non diabetic patients.

Organism	Diabetic samples	Non diabetic samples
E.coli	12 [66.7%]	2 [100%]
Proteus	4 [2.2%]	non
Staph. aureus	2 [11.1%]	non
total	18	2

Table 3: features of diabetic patients who had significant bacteriuria $\geq 10^5$ organism/ml}

Sex	Females = 14[77.8%] Males = 4 [22.2%]
Age	Females = 8>50 years 6< 50 years Males = 4 > 50 years
Type of Diabetic	IDDM =8 [44.5%] NIDDM = 10 [55.5%]
Number of pus cells	All more than 5 cell /H.P.F.
Types of organisms isolated	<u>Escherchia coli</u> = 12 [66.7%] , <u>Proteus</u> = 4 [22.2%] <u>Staphylococcus aureus</u> = 2 [11.1%]
Duration of Diabetes	4 = 2-5 years 6 = 5-10 years 4 = 10-15 years 4 = > 15 years

DISCUSSION:

In the majority of urinary tract infections bacteria gain access to the bladder and then to the renal pelvises via the urethra. Hematological spread of infection may be blamed but in rare cases renal paronychymal infections may be a result of this descend.

This study has revealed a higher prevalence of asymptomatic bacteriuria among diabetic patients than non diabetic patients, this is comparable with the study by Nicolle in 2005⁽¹⁵⁾ also other studies^(9,16,17), although some investigators demonstrated no significant difference in the prevalence of asymptomatic bacteriuria between diabetic and non diabetic women [p value=0.02]^(18,19).

It also revealed that a higher prevalence of asymptomatic bacteriuria in females in both groups [diabetic and non diabetics], which confirms the results of a study by Nicolle et al in 2006⁽²⁰⁾. As most other previous studies^(21,22), *Escherchia coli* was the most prevalence organism in both groups, in few studies ,the microorganism was different for example *Klebsiella* was the most common organism^(23,24).

The factors that predispose to periurethral colonization with Gram negative bacilli remain poorly understood but probably include alteration of normal perineal flora.

The higher prevalence of bacteriuria among females could be due to:-

- Presence of short urethra in females.
- The female urethra appears to be particularly prone to colonization with colonic gram-negative bacilli because of its proximity to the anus.
- Sexual intercourse causes the introduction of bacteria to the bladder.
- Absence of prostatic secretion which contain bactericidal properties.

The explanation of the above results may be due to:

- The presence of significant amount of glucose in diabetic urine, which serves as a favorite media for growth of bacteria^(19,25).
- a change in bacteria adhesion to the uroepithelium ,partly as a result of abnormal intracellular calcium metabolism which lead to decrease in Tam-horsfall protein which usually adhere to the bacteria and prevent attachment to uroepithelium is involved in the pathogenesis in the urinary tract infection in diabetic patient⁽²⁶⁾.

- Abnormal intracellular calcium metabolism lead to granulocytes dysfunction which leads to more attachment of bacteria to uroepithelium and more infection⁽²⁶⁾.
- *Escherchia coli* with type 1 fimbriae adheres better to uroepithelial cells in women with diabetes mellitus than to those without Diabetes⁽²⁷⁾.

There are studies consistently documents that the prevalence of asymptomatic bacteriuria is not influenced by type or duration of diabetic or by the quality of diabetic control and this is in agreement with the results of this study⁽²⁶⁾.

The presence of pus cells in the GUE is sensitive for the presence of infection as in patients in this study who had significant bacteriuria(> 3 pus cell / H.P.F.) but presence of pus cells is less specific and it occurs in other conditions like nephrocalcinosis, interstitial nephritis, polycystic kidney disease and anatomical abnormalities⁽²⁸⁾.

CONCLUSION:

The prevalence of asymptomatic significant bacteriuria is higher among patients with diabetic compared to non diabetic patients and it is more common in females than males.

Escherchia coli is the most common organism that causing asymptomatic bacteriuria in both diabetics and non diabetics.

REFERENCES:

1. Feleke Y, Mengistu Y, Enquesslassie F. Diabetic infections: clinical and bacteriological study at Tikur Anbessa Specialized University, Addis Ababa, Ethiopia. *Ethiop Med J*. 2007;45,171-9
2. Unachukwu CN, Obunge OK, Odi OJ. The bacteriology of diabetic foot ulcers in Port Harcourt, Nigeria. *Niger J Med*. 2005;14,173-176.
3. Patterson VT, Andriole, Bacterial urinary tract infections in diabetes, *Infect. Dis. Clin. North Am*. 1997;11,735-750
4. Gul M; Kurutas E, Ciragil, P, Cetinkaya A, Kilinc M, Aral M, Buyukbese MA. Urinary tract infection aggravates oxidative stress in diabetic patients. *Tohoku J Exp Med*. 2005; 206, 1-6
5. Nicolle LE. Asymptomatic Bacteriuria in diabetic women. *Diabetes Care* 2000;23,722-723.
6. Hikmat K., Abdulla K. Microscopic pyuria in the diagnosis of symptomatic urinary infection. *Fac Med Baghdad* 1998;40,447-452.

SILENT BACTERIURIA IN DIABETES

7. AlHeeti KJ. Bacteriuria in Iraqi pregnant women in Baghdad. 1999; M.Sc. thesis is submitted to the college of Medicine, University of Baghdad.
8. Geerlings SE, Stolk RP, Camps MJ et al. Asymptomatic Bacteriuria may be considered a complication in women with diabetes. *Diabetes Care* 2000;23,744-9
9. Zhanel GG, Nicolle LE, Harding GK. Prevalence of asymptomatic Bacteriuria and associated host factors in women with diabetes mellitus. The Diabetic Urinary Infection Study Group. *Clin Infect Dis.* 1995;21,316-22
10. Okada H, Shirakawa T, Gotoh A, et al. Enumeration of bacterial cell numbers and detection of significant Bacteriuria by use of a new flow cytometry-based device. *Clin Microbiol.* 2006;44,3569-9.
11. Okada H, Sakai Y et al. Detection of significant Bacteriuria by automated urinolysis using flow cytometry. *J of Clin Microbiol.* 2000;38,2870-2872
12. Barlett RCDA, Zern I et al. Screening for urinary tract infection with the Yellow IRIS. *Lab Med.* 1992;23,599-602
13. Al-Daghistani HI, Abdel-Dayem M. Diagnostic value of various urine tests in the Jordanian population with urinary tract infection. *Clin Chem Lab Med.* 2002;40,1048-51.
14. Bachman JW, Heise JM, Naessens JM, Timmerman MG. A study of various to detect asymptomatic UTIs in an obstetric population. *JAMA* 1993;270,1971-4.
15. Nicolle LE. Urinary tract infection in diabetes. *Curr-Opin-Infect-Dis.* 2005;18,49-53.
16. Kayima TK, Otieno LS, Twahir A, Njene E. Asymptomatic Bacteriuria among diabetes attending Kenyatta National Hospital, East Afr. *Med J.* 1996;37,524-6.
17. Epoke J, Odigwe CO, Augonwn GOC. Prevalence of significant Bacteriuria in diabetic patients in Calabar, Nigeria. *Diabetic Internat.* 2000;10,16-17.
18. Brauner A, Fodin N, Hylander B, Ostenon CG. Bacteriuria, bacterial virulence and host factors in diabetic patients. *Diabet. Med* 1993;10,550-4.
19. Bonadio M, Boldrini E, Forotti G et al. A symptomatic Bacteriuria in women with diabetes, influence of metabolic control. *Clin Inf Dis.* 2004;38,41-45.
20. Nicolle LE, Zhanel GG, Harding GK. Microbiological outcomes in women with diabetes and untreated asymptomatic Bacteriuria. *World J Urol.* 2006;24,61-5.
21. Boroumand MA, Sam L et al. A symptomatic Bacteriuria in type 2 Iranian diabetic women: a cross sectional study. *BMC Women's Health* 2006,6:4 (page number not for citation purposes).
22. Razak SA, Khaki I. Some virulence factors of E. coli associated with various forms of urinary tract infection. *J Fac Med Baghdad* 1999;41,261-272.
23. Alebiosu CO, Osinube OA, Olajubu FA. Significant asymptomatic Bacteriuria among Nigerian type 2 diabetes. *J Natl Med Assoc.* 2003;95,344-351.
24. Bonadio M, Meini M, Spitalevi P, Giglic: Current microbiological and clinical aspects of urinary tract infections. *Eur Urol* 2001;40,439-44.
25. Boyko EL, Fihn SD, et al. Diabetes and the use of acute urinary tract infection among postmenopausal women. *Diabetic care* 2002;25,1778-1783.
26. Lindsay E, Nicolle MD. *New Eng J Med* 2000;343,1037-1039.
27. Geerling SE, Meiland R, Hoepelman IM. Urinary tract infection in women with diabetes mellitus. *Ned Tijdschr Geneesk* 2001;145,1832-6.
28. Walter E, Stamm. Urinary tract infections and pyelonephritis, principles of internal medicine, New York 1998, 817-823.