


## ***Haemonchus contortus*: Conventional and molecular identification and characterization via sequencing-based phylogenetic analysis in sheep**

Athraa Najm Abed      Monyer Abdulamier Abd Alfatlawi 

<sup>1</sup>Department of Veterinary Microbiology, College of Veterinary Medicine, University of Al-Qadisiyah, Al-Diwaniyah City, Iraq.

**Submitted:** July 10, 2024

**Revised:** July 31, 2024

**Accepted:** August 02, 2024

**Abstract** The purpose of this study, conducted from 01-October-2023 to 28-February-2024, was to determine the prevalence and genetic diversity of *Haemonchus contortus* Infection from sheep in Al-Diwaniyah slaughterhouse, it was carried out to investigate the level of infection at two levels stages; adult's animal, and human, and also helped us to evaluate the genetic similarities between local isolates from Iraq in comparison with other strains of *H. contortus* from all over the world. In the slaughterhouse in Al-Diwaniyah City, 250 carcasses were sampled. The samples were divided into two tubes, one of them used for microscopic examination by lacto phenol-solution and the other used for molecular tests by using 70% ethanol. Targeting the  $\beta$ -tubulin region by PCR was done, which was followed with the positive PCR-based sequencing. The findings of this study revealed high prevalence of *H. contortus* infection rate in sheep which leads to parasite transmission among the animals. In addition, a high level of genetic similarities was seen between local isolates from Iraq and strains from Sweden, Canada, Switzerland, the USA, and the Netherland. Overall, this study concludes that the level of infections by *H. contortus*. High genetic similarities, with world strains, indicating high diversity and common ancestor.

**Keywords:** Enteral parasites, *Haemonchus contortus*, roundworms

**Introduction** *Haemonchus contortus*, one of the most important gastrointestinal nematode parasites in the abomasum of small ruminants, primarily infects at the wetland map bordering the northern tropical and subtropical zones. The traditional control method using chemical anthelmintics is developing drug resistance (1-3).

*Haemonchus contortus* is a hematophagous parasitic nematode of the abomasum, being one of the most harmful due to the mass production of eggs that facilitates its transmission and the economic losses by non-productivity that it causes to livestock producers (3). Its control is not only anthropocentric; it is also influenced by the presence of other gastrointestinal nematodes, such as *Trichostrongylus*, which also affect the quality of life and productivity of the domestic ruminants. The reproductive strategy of this nematode counts on a very short egg-to-egg generation time, the ability to survive for a long time without feeding, and a reduced number of cell division cycles, making it difficult to detect them and apply the appropriate management alternative. The lack of genetic and physiological knowledge of this parasitic nematode has delayed the development of effective treatments, but the interest of the scientific community has allowed advances to be made in

understanding which areas of their biology this nematode might exploit to survive and replicate (5-7). Sheep are the animals most affected by these parasites, which cause anemia and even death if infection levels reach high values. However, other susceptible hosts include goats, wild ruminants, camelids, and domestic cattle; in the latter, the presence of these parasites directly compromises the economic losses associated with milk production and reproduction (5, 8, 9).

*H. contortus* is an intestinal nematode with a direct life cycle and proportional dependence on the environment for transmission of its infective stages. The life cycle depends on the release of eggs containing larvae (L1) by adult females in the host digestive system and environmental factors like temperature, humidity, and oxygen. The suitable conditions favor the development of cuticular larvae (L2) and directly expose infective third-stage larvae (L3) to the external environment. In moderate climates, they favor the survival and transmission of the infective larvae to new hosts throughout the year. The ability of the L3 to cope with oral ingestion by the ruminant allows them to remain free on the pasture for long periods (10, 11).

The genus *Haemonchus* can cause haemonchosis, severe damage, and increased susceptibility to abomasal gastritis. This condition can be fatal in sheep

and goats. According to previous reports, this nematode caused the highest economic damage to small ruminant production. Since the 70s of the last century, parasitic nematodes of the *Haemonchus* species have developed resistance to anthelmintic drugs practically used in control. According to the FAO, parasitic nematode resistance to anthelmintic agents is a widespread problem worldwide; some countries have reported resistance to all families of anthelmintic drugs. Widespread use of one or more drug groups can cause a high level of resistance and multiple drug resistance in some parasitic nematodes (12-13).

The aim of this study was to determine the prevalence & genetic diversity of *H. contortus* Infection from sheep in Al-Diwaniyah slaughterhouse, it was carried out to investigate the level of infection at three levels stages; adults, animal, and human, and also helped us to evaluate the genetic similarities between local isolates from Iraq in comparison with other strains of *H. contortus* from all over the world.

#### Material and Methods

##### Ethical approval

The current study was approved by the Ethical Committee in the College of Veterinary Medicine, University of Al-Qadisiyah, Al-Diwaniyah City, Iraq.

##### Samples

These samples were randomly collected from the sheep abomasum in three different months from October 2023 to February 2024 during the slaughtering sheep at the abattoir of Al-Diwaniyah slaughterhouse. Each month, 50 carcasses were examined and were randomly selected from male and female sheep, 25 of each sex. The samples obtained were submitted for macroscopic examination where each sample was cut into half, we put one part in one disposable cup which contained lactophenol solution which is used in the microscopic examination and the other part of each sample was put in containers which contains 70% ethanol which used in the molecular tests.

##### Microscopic examination

For microscopic studies, the worms were taken directly onto glass slides and examined under a universal light microscope with 10X and 40X magnifications to confirm their identity. Additionally, the worm was examined for its morphology. Preserving the worm in lactophenol solution helps in its clear visualization of structures and is essential for

identifying and differentiating it from other parasitic species (14).

##### Molecular techniques

The DNA was extracted using DNA extraction kit (AddBio, Korea) and following the kit protocol. The  $\beta$ -tubulin gene region of this extracted DNA was amplified using Polymerase Chain Reaction (PCR). The primers were designed for the  $\beta$ -tubulin region with the following sequences and amplified the 785 bp:  $\beta$ -tubulin-F 5' GGAACAATGGACTCTGTTCG 3'  $\beta$ -tubulin-R 5' GAATCGAAGGCAGGTCGT 3' (Accession number KF269107). Designed by Zongze et al (2018) the PCR compound was as followed: 10 ml of master mix (AddBio) 2 ml of forward primer 0.5 pmol/20  $\mu$ l 2 ml of reverse primer 0.5 pmol/20  $\mu$ l 4 ml of PCR water 2  $\mu$ l of DNA template The thermocycling conditions were as followed: Initial denaturation 95°C 5 min (1 cycle) Denaturation 95°C 35 s (40 cycles) Annealing 55°C 30 s (40 cycles) Extension at 72°C 30 s (40 cycles) Final extension 72°C 5 min (1 cycle).

PCR products were loaded on a 1.5% agarose gel for electrophoresis to determine the appearance and size of the amplicons. Here, 5  $\mu$ l of PCR products was loaded into each gel well and conducted electrophoresis at 100 volts and 80 amperes for 1 hour. The gel was then placed in the gel documentation system and detect the DNA bands.


##### Sequencing

First, 10 positive PCR samples were chosen for DNA sequencing and phylogenetic analysis. The products were purified, and then forwarded samples for the Sanger sequencing system (PCR products were sent to Macrogen Company in Korea). The nucleotide sequences from the Sanger sequencing system were trimmed for noise signals and sent to the NCBI-GenBank to obtain accession numbers. The species then were determined using DNA sequencing and phylogenetic tree analyses by the MEGA X software with multiple sequence alignments based on Clustal W alignment analysis. The comparison of the local isolates with other global sequences from the NCBI-Blast database included species typing and evolutionary relationships were investigated.

##### Results


The number of infected sheep and the relative percentages for each month were 8/50 (16%), 11/50 (22%), 14/50 (28%), 17/50 (34%), and 25/50 (50%) for October, November, December, January, and February, respectively.

In summation, 75 out of the 250 sheep analyzed tested positive for the infection, which gives an overall infection figure of 30%. We then calculated how the infection rates changed month by month and found it to be highly significant ( $p$ -value = 0.003). Figure (1) shows a steady rise in the infection rate. Macroscopic and microscopic examinations showed that the sheep were infested with *Haemonchus contortus* worms




**Figure 1:** Trend percentage in the infection rates by *Haemonchus contortus* in sheep.

The postmortem and microscopic appearance of the presence of the *H. contortus* is shown in Figure 2.



**Figure 2:** Macroscopic and microscopic appearance of *Haemonchus contortus* from sheep. Male and female. The PCR amplification revealed positive PCR products, which are shown in Figure 3.



**Figure 3:** Agarose gel electrophoresis of PCR products for  $\beta$ -tubulin gene region of *Haemonchus contortus* from sheep. Product size: 785 bp. Ladder: (100-3000 bp).


A high level of genetic similarities was seen between local isolates from Iraq and strains from Sweden, Canada, Switzerland, the USA, and The Netherland (Table 1 and Figure 4).

**Table 1:** Homology Sequence identity (%) between local isolates of *Haemonchus contortus* from sheep and world isolates.

| Accession number (Current study) | Accession number (World isolates) | Country     | Identity (%) |
|----------------------------------|-----------------------------------|-------------|--------------|
| PP393534                         | MK382798                          | Sweden      | 100          |
| PP393535                         | MK382800                          | Sweden      | 95.75        |
| PP393536                         | MK382797                          | Sweden      | 96.08        |
| PP393537                         | MK382761                          | Sweden      | 95.92        |
| PP393538                         | KF483614                          | Canada      | 95.92        |
| PP393539                         | DQ469245                          | Switzerland | 95.59        |
| PP393540                         | MK382772                          | Sweden      | 95.42        |
| PP393541                         | MK382788                          | Sweden      | 95.26        |
| PP393542                         | KX246652                          | USA         | 94.60        |
| PP393543                         | X80046                            | Netherlands | 94.60        |

## Discussion

The number of infected sheep slaughtered at the Al-Diwaniyah slaughterhouse increased during the study period, rising from 16.0% (October) to 50.0% (February). The trend suggests that the burden of the parasite *H. contortus* is increasing in the sampled population, with an overall infection rate of 30%. These data are in agreement with *H. contortus* infections reported from multiple locations around the world, though specific prevalence rates can vary by region and environmental conditions and based on management practices.



**Figure 4:** Phylogenetic tree of PCR products for  $\beta$ -tubulin gene region of *Haemonchus contortus* from sheep.

Other studies conducted in other parts of Iraq report different prevalence of *H. contortus* infection, a study in Baghdad report the prevalence among sheep to be 25.8 percent and a study in Basrah report 21.4 percent prevalence (15, 16). However, the higher prevalence in the current study %30 may be due to differences in the sampling period, climatic conditions, animal management and their associated environmental factors. Another important factor is seasonal variation and its effects on *H. contortus* prevalence on livestock, temperature of days and months were higher during the summer season, and humidity also increased, which influences the survival and transmission of the parasite. The trend, during the cooler months, was rising and may be associated with the most favorable of the time for *H. contortus* development and transmission.

Meanwhile, in the broader Middle Eastern region, previous studies have estimated the prevalence of *H. contortus* infections in the sheep at 28.3% (17), 33.7% (18) and 7.6% (19). Again, these results are similar to this current study, which indicates that there is a consistent appearance of *H. contortus* across the Middle East region and similar climates contribute to its persistence. Furthermore, we expect that

movement of livestock between and within the neighboring countries is the main reason behind the consistent appearance of *H. contortus* and similar prevalence rates across the region.

Internationally, wide variation in the prevalence of *H. contortus* infections have been reported. In countries practicing intensive livestock industry with routine worming of animals, researchers have reported low prevalence rates. Regassa et al (20) record a 12.5% prevalence which is important due to the stringent control measures and regular use of anthelmintics. In tropical and subtropical forests favored by climatic conditions for the survival of larvae of *H. contortus*, such high figures are prevalent; Mannan et al (21) reports 45.3% prevalence. In regions where intensive control measures are a rarity and climatic conditions are favorable for larval development, the parasite affects a significant percentage of the sheep population.

The detection and molecular analysis of the parasites present revealed high genetic identity of the isolates with various global strains of *H. contortus*, ranging from 94.60% to 100% sequence identities to strains from Sweden, Canada, Switzerland, USA and the Netherlands. These results revealed the genetic conservation of *H. contortus* across geographical regions, and also suggested close evolutionary relationships between local and international strains, indicative within sources and modes of transmission. Similar patterns have been observed using comparative studies. Laing et al. (22) described high levels of genetic similarity between *H. contortus* isolates from different parts of the world. These studies supported the hypothesis of a single origin of strains and global gene flow, leading to the potential dissemination of resistant alleles in association with high levels of gene flow. In conclusion, the high genetic similarities suggest that integrated and coordinated control of *H. contortus* infections is important because resistance genes can rapidly disseminate if not monitored (23-25).

#### Conclusion

In conclusion, the study revealed a very high prevalence of *H. contortus* infections in sheep in the Al-Diwaniyah region (33.8 to 50%) with an increase in rates of infections over the study period. The findings of the study on the molecular diagnostics and phylogenetic analyses showed a high degree of genetic similarities between local strains and strains circulating internationally, which urges for a close



monitoring of the local and the spread of *H. contortus* strains to minimize their economic impacts on livestock production. Finally, a comparison of the results of the present study with some studies from Iraq, the Middle East and worldwide shows the presence of *H. contortus* in sheep with persistent prevalence and genetic richness, which suggests the importance of integrated and region-specific HA control strategies.

#### Conflict of interest

There is no conflict of interest in this study as stated by the authors.

#### Acknowledgment

The authors would like to present their gratitude to the staff of the Parasitology Laboratory, College of Veterinary Medicine, University of Al-Qadisiyah.

#### References

1. Pervin HH. Prevalence of gastrointestinal parasitic infections and haemoprotozoan diseases of buffalo in coastal areas of Chattogram division, Bangladesh. 2023. doi:10.60015/bjvas/V08I2A10
2. do Nascimento JM, Brito SV, Teixeira AA, Frederico RG, Rodrigues AA, do Nascimento Sousa Filho JG, da Cunha IA. Potential distribution modelling for *Haemonchus contortus* (Nematoda: Trichostrongylidae) in South America. Parasitology Research. 2024 May;123(5):1-2. <https://doi.org/10.1007/s00436-024-08247-5>
3. Sharma P, Kaur A, Batish DR, Kaur S, Chauhan BS. Critical insights into the ecological and invasive attributes of *Leucaena leucocephala*, a tropical agroforestry species. Frontiers in Agronomy. 2022 May 24;4:890992. <https://doi.org/10.3389/fagro.2022.890992>
4. Lagos S, Moutzourelis C, Spiropoulou I, Alexandropoulou A, Karas PA, Saratsis A, Sotiraki S, Karpouzias DG. Biodegradation of anthelmintics in soils: does prior exposure of soils to anthelmintics accelerate their dissipation?. Environmental Science and Pollution Research. 2022 Sep;29(41):62404-22. <https://doi.org/10.1007/s11356-022-19964-8>
5. Kuiseu J, ZINSOU FE, Olounlade PA, ALOWANOU GE, Adenile AD, Dansou CC, Hounzangbe-Adote S, Babayemi OJ, Edorh PA. Prevalence, effects and alternative control methods of *Haemonchus contortus* in small ruminants: a review. Journal of Veterinary Medicine and Animal Health. 2021 Apr 30;13(2):84-97. doi:10.5897/JVMAH2020.0868
6. Jiménez-Penago G, Hernández-Mendo O, González-Garduño R, Torres-Hernández G, Torres-Chablé OM,

Maldonado-Simán E. Mean corpuscular haemoglobin concentration as haematological marker to detect changes in red blood cells in sheep infected with *Haemonchus contortus*. Veterinary Research Communications. 2021 Dec;45(4):189-97. <https://doi.org/10.1007/s11259-021-09800-8>

7. Adduci I, Sajovitz F, Hinney B, Lichtmannsperger K, Joachim A, Wittek T, Yan S. Haemonchosis in sheep and goats, control strategies and development of vaccines against *Haemonchus contortus*. Animals. 2022 Sep 8;12(18):2339. <https://doi.org/10.3390/ani12182339>

8. Flay KJ, Hill FI, Muguiro DH. A Review: *Haemonchus contortus* Infection in Pasture-Based Sheep Production Systems, with a Focus on the Pathogenesis of Anaemia and Changes in .... Animals. 2022. <https://doi.org/10.3390/ani12101238>

9. Werszko J, Wilamowski K, Kraszewska O, Bakier S, Pyziel AM. First molecular identification of *Haemonchus contortus* (Nematoda: Trichostrongylidae), a blood-sucking gastric nematode of artiodactyles, in the ground beetle *Carabus granulatus* (Coleoptera: Carabidae). Medical and Veterinary Entomology. 2024 Mar 27. <https://doi.org/10.1111/mve.12715>

10. Aleuy OA, Kutz S. Adaptations, life-history traits and ecological mechanisms of parasites to survive extremes and environmental unpredictability in the face of climate change. International Journal for Parasitology: Parasites and Wildlife. 2020 Aug 1;12:308-17. <https://doi.org/10.1016/j.ijppaw.2020.07.006>

11. Chylinski C, Cortet J, Cabaret J, Blanchard A. *Haemonchus contortus* Adopt Isolate-Specific Life History Strategies to Optimize Fitness and Overcome Obstacles in Their Environment: Experimental Evidence. Animals. 2023. <https://doi.org/10.3390/ani13111759>

12. DANSOU CC, KUISEU J, OLOUNLADE PA, AZANDO EV, LAGNIKA L, ABOH AB, EDORH PA, HOUNZANGBE-ADOTE MS. Haemonchosis: a review on dreaded strongylosis that affects the zootechnical performance of sheep. Journal of Parasitology and Vector Biology. 2021 May 31;13(1):58-70. doi:10.5897/JVPB2020.0405

13. Tilahun T. Review on Prevalence of *Haemonchus contortus* in Ethiopian. Journal of Pathology Research Reviews and Reports. SRC/JPR-145. 2021;3(4):2-4. [https://doi.org/10.47363/JPR/2021\(3\)136](https://doi.org/10.47363/JPR/2021(3)136)

14. Klaif SF, Jassim A, Alfatlawi MA, Ali MA. Major-surface-protein-4-gene-based detection of *Anaplasma marginale* isolated from sheep in Al-Diwaniyah province, Iraq. *Iraqi J Vet Sci.* 2022;36(3):85-88. doi:10.33899/ijvs.2021.129230.1635
15. Ahmad N, Khan SA, Majid HA, Ali R, Ullah R, Bari A, Akbar NU, Majid A. Epidemiology and phylogeny of *Haemonchus contortus* through internal transcribed spacer 2 gene in small ruminants. *Frontiers in Veterinary Science.* 2024 Apr 9;11:1380203. <https://doi.org/10.3389/fvets.2024.1380203>
16. Jabar Jasim H, Mijbas Mohammed Alomari M, Abbas Ali N, Khalid Alani Z, Abed SM, Nasser Kadim A. Prevalence, haematological and molecular studies of *Haemonchus contortus* isolated from goat at AL-Muthanna Province, Iraq. *Arch Razi Inst.* 2023 Feb 28;78(1):287-295. doi: 10.22092/ARI.2022.359004.2353.
17. Ahmed AI, Aziz KJ, Abdullah SO. Prevalence of gastrointestinal nematodes parasites from sheep and evaluation of some anthelmintic resistance in Erbil Governorate. *Bas J Vet Res.* 2015;14(1):1-11. doi:10.33762/bvetr.2015.99659
18. Mushonga B, Habumugisha D, Kandiwa E, Madzingira O, Samkange A, Segwagwe BE, Jaja IF. Prevalence of *Haemonchus contortus* Infections in Sheep and Goats in Nyagatare District, Rwanda. *J Vet Med.* 2018 Sep 6;2018:3602081. doi: 10.1155/2018/3602081.
19. Al-Qureishy SA. Prevalence of cestode parasites in sheep slaughtered in Riyadh City, Saudi Arabia. *J Egypt Soc Parasitol.* 2008;38(1):273-80. doi:10.1016/j.sjbs.2017.01.056
20. Mushonga B, Habumugisha D, Kandiwa E, Madzingira O, Samkange A, Segwagwe BE, Jaja IF. Prevalence of *Haemonchus contortus* infections in sheep and goats in Nyagatare District, Rwanda. *J Vet Med.* 2018;3602081:9. doi:10.1155/2018/3602081.
21. Regassa D, Birhane A, Getahun Y, Chalchisa A. Prevalence and Associated Risk Factors of *Haemonchus contortus* in Sheep Slaughtered at Bahir Dar Municipal Abattoir, Ethiopia. *J Parasitol Res.* 2024 May 15;2024:1433264. doi: 10.1155/2024/1433264.
22. Mannan, M.A., Chowdhury, S., Hossain, M.A. et al. Genetic variability of *Haemonchus contortus* isolates in small ruminants from slaughterhouses in Bangladesh. *Parasitol Res* 122, 3101–3107 (2023). <https://doi.org/10.1007/s00436-023-08000-4>
23. Laing R, Kikuchi T, Martinelli A, Tsai IJ, Beech RN, Redman E, Holroyd N, Bartley DJ, Beasley H, Britton C, Curran D, Devaney E, Gilabert A, Hunt M, Jackson F, Johnston SL, Kryukov I, Li K, Morrison AA, Reid AJ, Sargison N, Saunders GI, Wasmuth JD, Wolstenholme A, Berriman M, Gilleard JS, Cotton JA. The genome and transcriptome of *Haemonchus contortus*, a key model parasite for drug and vaccine discovery. *Genome Biol.* 2013 Aug 28;14(8):R88. doi: 10.1186/gb-2013-14-8-r88.
24. Alfatlawi MSH, Alfatlawi MA. Molecular and phylogenetic study of *Theileria* spp isolated from ticks in Al-Diwaniyah City, Iraq. *Iraqi J Agricul Sci.* 2019;50(1):475-79. doi: <https://doi.org/10.36103/ijas.v50i1.313>
25. Alfatlawy HH, Alfatlawi MA. Morphological and molecular identification of *Parabronema skrjabini* of camels (*Camelus dromedary*) in Najaf province. *Iraqi J Vet Sci.* 2021;35(3):507-12. doi: 10.33899/ijvs.2020.127101.1459