

4-23-2025

## Verification and Demonstration of the Ability of Plasma Generated from Dielectric Barrier Discharge to Oxidize

N. Yasoob A.

*Department of Physics, College of Science for Women, University of Baghdad, Baghdad, Iraq,*  
noorya\_phys@csu.uobaghdad.edu.iq

Follow this and additional works at: <https://bsj.uobaghdad.edu.iq/home>

---

### How to Cite this Article

Yasoob A., N. (2025) "Verification and Demonstration of the Ability of Plasma Generated from Dielectric Barrier Discharge to Oxidize," *Baghdad Science Journal*: Vol. 22: Iss. 4, Article 22.  
DOI: <https://doi.org/10.21123/bsj.2024.11217>

This Article is brought to you for free and open access by Baghdad Science Journal. It has been accepted for inclusion in Baghdad Science Journal by an authorized editor of Baghdad Science Journal.



## RESEARCH ARTICLE

# Verification and Demonstration of the Ability of Plasma Generated from Dielectric Barrier Discharge to Oxidize

N. Yasoob A. 

Department of Physics, College of Science for Women, University of Baghdad, Baghdad, Iraq

## ABSTRACT

The oxidation ability of cold plasma is a major influence on bacterial inactivation, lipid oxidation, medical equipment sterilization, and applications in air and wastewater purification, for this reason, the ability of cold plasma to oxidize materials is an important factor that must be studied. In this study, the ability of cold plasma generated by a dielectric barrier discharge (DBD) to oxidize materials was revealed and proven by using different exposure times (5, 10, 15, 20 min) and several measurements: the decolorization of indigo carmine from dark blue to yellow, the significant reduction in pH, and the increase in oxidation reduction potential (ORP) and electrical conductivity (EC). Proven spectrophotometrically in the presence of nitrogen and its oxide ions. It was observed that the exposure time of 15 min is the maximum period during which oxidation occurs, after which saturation occurs. Numerous reactive oxygen and nitrogen species are created upon the interaction of gas phase plasma and water vapor (supported by excited species, energies and plasma electrons).

**Keywords:** Dielectric barrier discharge (DBD), Decolorization, Electrical conductivity EC, Indigo carmine, Oxidation

## Introduction

Cold plasma is regarded as among the most modern technologies for sophisticated oxidation, cold plasma finds application in the fields of wastewater treatment and air purification, as well as in the treatment of various pollutants,<sup>1,2</sup> sterilizing medical equipment,<sup>3</sup> lipid oxidation,<sup>4</sup> biodecontamination,<sup>5</sup> and others.<sup>6,7</sup> Consequently, to maximize the chemical and biological effects of atmospheric non-thermal plasma discharges, research into the oxidation capacity of atmospheric plasma discharges is necessary.

The presence of nitrate and nitrate, a decrease in the medium's pH, an increase in conductivity, and chemical degradation are some indicators of plasma discharge's ability to oxidize. Nonthermal plasma can generate both oxidative and reductive radicals. There are several types of systems for generating nonther-

mal plasma such as DBD,<sup>8</sup> FE-DBD,<sup>9–11</sup> plasma jet,<sup>12</sup> and microwave plasma.<sup>13</sup>

A sophisticated oxidation process used to break down organic molecules in water is called non-thermal plasma (NTP). Whereas the exact chemical constitution of air, cold plasma discharge remains unclear, it is known that these discharges produce reactive oxygen species, and highly oxidizable nitrogen compounds. When the plasma interacts with the surrounding air, ionized air produces RONS. Delivery to the target, tissue, or fluid is what RONS are. At first, it was believed that RONS were the root cause of free radical aging.<sup>6</sup> Nonetheless, it is widely acknowledged that the therapy of disease may benefit from the use of exogenous RONS in small amounts. Several investigations have suggested a potential connection between angiogenesis, migration, apoptosis, and plasma-generated RONS in cell culture conditions.<sup>14–17</sup>

Received 19 March 2024; revised 28 May 2024; accepted 30 May 2024.  
Available online 23 April 2025

E-mail address: noorya\_phys@cs.w.uobaghdad.edu.iq (N. Yasoob A.).

<https://doi.org/10.21123/bsj.2024.11217>

2411-7986/© 2025 The Author(s). Published by College of Science for Women, University of Baghdad. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The oxidation effects of impulsive plasma discharge have been studied in this research. Surface discharges spread across an interface between liquid and air, while indirect discharges propagate through the liquid sample's bulk. The oxidizing capability was analyzed by estimating the degree of decolorization of IC dye in water solutions, Indigo carmine was chosen to serve as a chemical sensor. Numerous elements significantly contribute to these processes, including: reducing pH, oxidation reduction potential (ORP), Electrical conductivity (EC), concentration and transmittance of solution which are responsible for the oxidizing effect in principle.

In this paper, DBD is used to obtain cold plasma to investigate its chemical oxidation, a plasma oxidation process involves the use of active neutral and charged oxygen species to oxidize surfaces by discharging the ambient air, dielectric barrier discharges (DBDs), an advanced oxidation technology based on free radical processes, have been found to completely decompose. At the interface between gas and liquid, the NTP discharge caused the transient species (OH, O, and NO radicals) to develop. The indirect discharge treatment by using dielectric barrier discharge (DBD) possesses an efficiency of oxidation. The main effect that can be observed visually is the decolonization rate of indigo carmine, other factors related to the impacts on the chemical characteristics of the water that's a laboratory measured, such as pH, oxidation reduction potential (ORP) and electrical conductivity (EC).

## Materials and methods

### *Dielectric barrier discharge system*

DBD system is used to generate cold plasma to study the mechanism of oxidation capability of cold plasma, it consists of two electrodes made from stainless steel covered by Teflon and the quartz barrier between the electrodes as shown in Fig. 1.

The system was energized using a high voltage AC power supply, the magnitude of applied voltage about (15 kV), and consisted mainly of high-voltage electrodes, the cold plasma was obtained by ionizing the ambient air and applied on the sample for different times of exposure 0, 5, 10, 15, 20 min, and the electrode was separated from the solution surface by (2–3 mm) for discharge and plasma generation.

### *Indigo carmine*

Indigo carmine pigment (IC) served as a chemical sensor for examining the cold plasma's ability for oxidation. Indigo carmine is a food coloring and safe to use, nontoxic, a powder is typically a dark blue. The

formula of IC is (C<sub>16</sub>H<sub>8</sub>N<sub>2</sub>Na<sub>2</sub>O<sub>8</sub>S<sub>2</sub>), the molecular weight is 446.35 g/mol.<sup>18</sup> The solution was prepared by using deionized water to obtain a solution with concentration (0.35 mM) and pH 5.5. An essential component in assessing a solution's efficacy is its pH of non-thermal plasma oxidation in the medium. The pH, EC, ORP were measured by (PH-2603). The optical emission and transmission of the solution were measured by UV-visible spectrophotometer (SEC 2000 spectra system). The solution should be stored in a cool, dark place.

## Results and discussion


### *Oxidation effect on coloring, pH, electric conductivity, ORP of IC*

The study examined the oxidizing impacts of transient discharges that propagate through the majority of the sample. A 10 mL sample of IC solution was put into a Petri dish and exposed to plasma for each separate time, just like with the surface discharges. (0, 5, 10, 15, 20 min), the color was changed gradually from dark blue to green yellowish as shown in Fig. 2(a).


We show that the color of indigo solution changes after being exposed to plasma, initially transferring to dark green after 5 min of cold plasma exposure, then becoming light green after 10 min of plasma, and after 15, 20 min turning to yellowish, the oxidation or acidic effect of the decrease in pH caused to change color from blue to yellow as shown in Fig. 2.

After 24 hours of exposure, the color of the solution that's exposed for 5 min becomes light green, and the sample exposed for 10, 20 min it becomes yellow, as shown in Fig. 2. Finally, after 72 h, the color of IC solution for all time becomes yellow for all time, which proves the lifetime of the plasma effect is long and not temporary. The reason for this behavior is due to the ability of plasma to oxidize, and it is considered a preliminary indicator to prove the ability of plasma to oxidize, it agree with.<sup>19</sup>

The electrical conductivity, pH, and oxidation reduction potential of the pigment solutions were measured before and after plasma treatment. An improvement in the solutions' electrical conductivity from 0.12 to 0.34 ms/cm was observed as shown in Fig. 3(a) and an increase in ORP, the ORP measurement is related to the oxidation potential and formed the main species from non-thermal plasma as shown in Fig. 3(b). A drop in the pH of aqueous dye solutions was seen in all treatment cases, which could be the result of nitric and nitrous acid production, coming from the formation of the nitrogen and its compounds nitrate and nitrate, non-thermal plasmas can activate



**Fig. 1.** High voltage Dielectric barrier discharge system.



**Fig. 2.** Decolorization of indigo carmine for different time of exposure to cold plasma and for different days (a): 0-day, (b) 1 day, (c) 3 days.

the reactants  $N_2$  and  $H_2/H_2O$ , which then break down into reactive species like N, H, and OH radicals. These reactive species will eventually recombine to form intermediate compounds and  $NH_3$  as indicated

by Eqs. (8) to (19)<sup>19–21</sup> Significantly, with increasing oxidation, the pH decreased from  $\sim 5.5$  to a minimum value of  $\sim 3.18$  as shown in Fig. 3(c). All these indicate an oxidation effect and agree with.<sup>22,23</sup>




Fig. 3. Effect of cold plasma on (a), conductivity (b) ORP, (c) pH.


The initial sample solution that was not exposed was tested as a reference. The control sample in this instance was distilled water. Fig. 4 illustrates how the unexposed sample's transmittance differs from that of distilled water. The transmittance ( $T\%$ ) is less than 100% in the visible range which agrees with 23. This indicates that, in comparison to pure water, the IC solution has a reduced transmittance. Otherwise, a greater amount of visible light is absorbed by the IC sample solution.

Next, the exposed sample for plasma is utilized as the target sample, and the original unexposed IC solution is used as the control sample if the transmittance is less than 100%. One measures the differential transmittance spectrum. Two instances of the differential transmittance spectra of the "exposed sample to unexposed sample" were exposed to 5, 10, 15, and 20 min. The transmission increased gradually after 5 min and the peak shifted into blue green region about 452–539 nm as shown in Fig. 4(b), while after 10 min the transmittance rose and the peak shifted to a green region (550 nm) as shown in Fig. 4(c), even though 15 and 20 min a raised to same result as shown in Fig. 4(d), (e) that means the solution reached to saturation after

15 min, the transmittance shifted to yellow region (586 nm). As a result to change the pH the color may be change from blue to yellow. Consequently, since this is the wavelength range where blue to yellow light falls, it is important to look into changes in the wavelength of the differential spectrum between 450 and 590 nm, its agree with.<sup>24</sup>

#### Optical emission analysis

Fig. 5 represents the emission spectroscopy of CAP by DBD for the wavelength (250–500 nm), the reactive species produced by the DBD system are shown in the spectrum. The reference<sup>16</sup> was followed to identify the emission bands. It can be seen that modest emission bands for NO lines were found in the 250–300 nm wavelength range.<sup>16,17</sup> The direct electron impact excitation is responsible for the dispersion of the  $N_2(C_3\Pi_m)$  excited state (having enough energy to pass the threshold) from the ground state  $N_2(X_1\Sigma_g^+)$ . The characteristic photons of the second positive band head (0–0) with a wavelength of 337.1 nm are released by the following radiative decays. The



**Fig. 4.** The transmission of Indigo solution before and after exposed to cold plasma (a) untreated (b) 5 min. (c) 10 min. (d) 15 min. (e) 20 min.

wavelengths of 315.29, 337, 358, and 380.4 nm could be analytic of the low-intensity  $N_2$  second-positive system.  $N_2$

$$(C_3\Pi_u) \nu' = 0 \Rightarrow N_2 B_3\Pi_g \nu'' = 0 \quad (1)$$

The dominant nitrogen spectrum lines are 315.29 nm to 470 nm. The posterior radioactive degradation of distinct photons is dissolved from the second positive-band header (0-0) and has a wavelength of 337 nm.






Fig. 5. Optical emission spectrum of cold plasma by DBD.











Fig. 6. Reactive species production in plasma discharge, gas phase, gas-liquid phase, and liquid phase represented graphically.

Consequently, the species has been identified in Fig. 6 produced at the cold plasma gas-liquid interface; the primary species generated by cold plasma migrate by diffusion to the liquid phase and then stabilized by water forming the secondary species<sup>24</sup> as shown in Fig. 6.





Eqs. (2) to (19)<sup>24,25</sup> represented the products of a process known as indirect discharge, in which plasma generated in the gas phase comes into contact with water vapor and is assisted by both plasma electrons and the energies of excited species, produces a lot of reactive oxygen species in addition to nitrate, nitrite, and NO radicals.

## Conclusion

The oxidation capability of cold plasma is a major influence on many applications such as inactivation of bacteria, lipid oxidation, sterilization of medical equipment, and applications in air and wastewater purification. Indigo carmine dye (IC) served as an effective chemical sensor for examining the cold plasma's ability for oxidation. It was shown that microscope by decolorization of solution and microscope is done by measuring electric conductivity EC, pH, oxidation reduction potential (ORP) and optical emission spectroscopy and observed the main RONS species that produced, we show that the oxidation increased with increased time exposure for plasma by increased in EC, ORP and decreasing pH, and arrived to saturation time after 15 min from being exposed to cold plasma for 10 ml from solution.

## Authors' declaration

- Conflicts of Interest: None.
- I hereby confirm that all the Figures and Tables in the manuscript are mine. Furthermore, any Figures and images, that are not mine, have been included with the necessary permission for republication, which is attached to the manuscript.
- No animal studies are present in the manuscript.
- No human studies are present in the manuscript.
- Ethical Clearance: The project was approved by the local ethical committee at the University of Baghdad.

## References

1. Aggelopoulos CA. Recent advances of cold plasma technology for water and soil remediation: A critical review. *Chem Eng* J. 2022;428(4):131657. <https://doi.org/10.1016/j.cej.2021.131657>.
2. Razuqi NS, Muftin FS, Murbat HH, Abdalameer NK. Influence of dielectric-barrier discharge (DBD) cold plasma on water contaminated bacteria. *Annu Res Rev Biol*. 2017;14(4):1–9. <http://dx.doi.org/10.9734/ARRB/2017/34642>.
3. Vichiansan N, Kaai N, Leksakul K, Lumyong S, Han JG. Sterilization of medical equipment by plasma technology. *Chiang Mai J Sci*. 2018;45(4):1811–1826. <http://epg.science.cmu.ac.th/ejournal/>.
4. Pérez-Andrés JM, Cropotova J, Harrison SM, Brunton NP, Cullen PJ, Rustad T, *et al*. Effect of cold plasma on meat cholesterol and lipid oxidation. *Foods*. 2020;9(12):1786. <http://dx.doi.org/10.3390/foods9121786>.
5. Mravlje J, Regvar M, Vogel-Mikuš K. Development of cold plasma technologies for surface decontamination of seed fungal pathogens: Present status and perspectives. *J Fungi*. 2021;7(8):650. <https://doi.org/10.3390/jof7080650>.
6. Barjasteh A, Dehghani Z, Lamichhane P, Kaushik N, Choi EH, Kaushik NK. Recent progress in applications of non-thermal plasma for water purification, bio-sterilization, and decontamination. *Appl Sci*. 2021;11(8):3372. <https://doi.org/10.3390/app11083372>.
7. Mohammed SA, Al-Khazrajy OS, Abdallah M, Aadim KA, Al-Mamari A, Al-Owaisi H, *et al*. Removal of dyes from aqueous solutions using non-thermal plasma. *Environ Process*. 2023;10(4):1–19. <https://doi.org/10.1007/s40710-023-00677-0>.
8. Kadum E. Effect of the dielectric barrier discharge plasma on the optical properties of CDS thin film. *Baghdad Sci J*. 2019;16(4(Suppl.)):1030–1035. [http://dx.doi.org/10.21123/bsj.2019.16.4\(Suppl.\).1030](http://dx.doi.org/10.21123/bsj.2019.16.4(Suppl.).1030).
9. Yasoob AN, Murbat HH, Khaleel KJ, eds. The influence of cold atmospheric pressure plasma on TSH and thyroid hormones in male rats. In *AIP Conf Proc*. 2020;2213:020015. <http://dx.doi.org/10.1063/5.0000125>.
10. Yasoob AN, Khaleel KJ, Murbat HH. Study of the effect of cold plasma on the reproductive endocrinology of male rats. *Plant Arch*. 2020;20(1):526–31. [https://plantarchives.org/SPECIAL%20ISSUE%2020-1/104\\_526-531\\_.pdf](https://plantarchives.org/SPECIAL%20ISSUE%2020-1/104_526-531_.pdf).
11. Yasoob AN, Khaleel K, Murbat H, eds. Effect of cold atmospheric pressure plasma on fertility hormones for female rats. In *IOP Conf Ser. Mater Sci Eng*. 2020;757:012069. <http://dx.doi.org/10.1088/1757-899X/757/1/012069>.
12. Abbas BH, Abdalameer NK, Mohammed RS. Synthesis of copper oxide nanostructures by ar plasma jet and study of their structural and optical properties. *Iraqi J. of Sci*. 2024;65(4):1999–2006. DOI: 10.24996/ij.s.2024.65.4.18.
13. Noori AS, Mageed NF, Abdalameer NK, Mohammed MK, Mazhir SN, Ali AH, Jaber NA, *et al*. The histological effect of activated aloe vera extract by microwave plasma on wound healing. *Chem Phys Lett*. 2022;807:140112. <https://doi.org/10.1016/j.cplett.2022.140112>.
14. Kim S, Kim C-H. Applications of plasma-activated liquid in the medical field. *Biomedicines*. 2021;9(11):1700. <http://dx.doi.org/10.3390/biomedicines9111700>.
15. Halliwell B, Gutteridge JM. Free radicals in biology and medicine: Oxford university press, USA. 2015;488–489 p. <http://dx.doi.org/10.1093/acprof:oso/9780198717478.001.0001>.
16. Chen Z, Yu S, Lin L, Cheng X, Gjika E, Keidar M. Effects of cold atmospheric plasma generated in deionized water in cell cancer therapy. *plasma process polym*. 2016;13(12):1151–1156. <https://doi.org/10.1002/ppap.201600086>.
17. Mahdi SS, Aadim KA, Khalaf MA. New spectral range generations from laser-plasma interaction. *Baghdad Sci*



- J. 2021;18(4):1328. <https://doi.org/10.21123/bsj.2021.18.4.1328>.
18. de Keijzer M, van Bommel MR, Keijzer RH-d, Knaller R, Oberhumer E. Indigo carmine: understanding a problematic blue dye. *Studies in Conservation*. 2012;57(sup1):S87–S95. <https://doi.org/10.1179/2047058412Y.0000000058>.
19. Krosuri A, Wu S, Bashir MA, Walquist M. Efficient degradation and mineralization of methylene blue via continuous-flow electrohydraulic plasma discharge. *J Water Process Eng* 2021;40:101926. <https://doi.org/10.1016/j.jwpe.2021.101926>.
20. Krosuri A, Wu S, Bashir MA, Walquist M. Efficient degradation and mineralization of methylene blue via continuous-flow electrohydraulic plasma discharge. *J Water Proc Eng*. 2021;40:101926.
21. Zhou D, Zhou R, Zhou R, Liu B, Zhang T, Xian Y, *et al*. Sustainable ammonia production by non-thermal plasmas: Status, mechanisms, and opportunities. *Chem Eng J*. 2021;421:129544. <https://doi.org/10.1016/j.cej.2021.129544>.
22. Li S, Timoshkin IV, Maclean M, MacGregor SJ, Wilson MP, Given MJ, *et al*. Oxidation and biodecontamination effects of impulsive discharges in atmospheric air. *IEEE Trans Plasma Sci*. 2016;44(10):2145–55. <https://doi.org/10.1109/TPS.2016.2581317>.
23. Reddy PMK, Raju BR, Karuppiiah J, Reddy EL, Subrahmanyam C. Degradation and mineralization of methylene blue by dielectric barrier discharge non-thermal plasma reactor. *Chem Eng J*. 2013;217:41–7. <https://doi.org/10.1016/j.cej.2012.11.116>.
24. Li S, Timoshkin IV, Maclean M, MacGregor SJ, Wilson MP, Given MJ, *et al*. Steady-state corona discharges in atmospheric air for cleaning and decontamination. *IEEE Trans Plasma Sci*. 2013;41(10):2871–8. <https://doi.org/10.1109/TPS.2013.2264903>.
25. Zeghioud H, Nguyen-Tri P, Khezami L, Amrane A, Assadi AA. Review on discharge plasma for water treatment: Mechanism, reactor geometries, active species and combined processes. *J Water Process Eng*. 2020;38:101664. <https://doi.org/10.1016/j.jwpe.2020.101664>.

## اثبات وتوضيح قابلية البلازما المتولدة بتفريغ الحاجز العازل على الاكسدة

نور يعسوب عبد الخضر

قسم الفيزياء، كلية العلوم للبنات، جامعة بغداد، بغداد، العراق.

### الخلاصة

إن قابلية البلازما الباردة على الأكسدة تلعب الدور الأساسي في قتل البكتيريا وأكسدة الدهون وتعقيم المعدات الطبية وتطبيقاتها في تنقية الهواء ومياه الصرف الصحي، ولهذا السبب فإن قدرة البلازما الباردة على أكسدة المواد هي عامل مهم يجب دراسته. في هذه الدراسة تم التحقق من قابلية البلازما الباردة الناتجة عن تفريغ حاجز العزل الكهربائي (DBD) على أكسدة المواد من خلال استخدام أوقات تعريض مختلفة (5, 10, 15, 20 دقيقة) وإثباتها من خلال عدة قياسات: تغير لون الصبغة القرمزية النيل من اللون الأزرق الداكن. تدريجياً إلى الأصفر مع تغير ازمان التعريض، الانخفاض الكبير في الاس الهيدروجيني مما يدل على حامضية الوسط بسبب الأكسدة، والزيادة في جهد الأكسدة (ORP) والتوصيلية الكهربائية (EC)، تم إثبات ذلك طيفياً من خلال لوحظ وجود العديد من أنواع الأكسجين والنيوتروجين التفاعلية عند تفاعل بلازما طور الغازي وبخار الماء بتعزيزه (الأنواع التفاعلية المثارة، والطاقات والالكترونات الناتجة من التفاعل). وقد لوحظ أن مدة التعرض البالغة 15 دقيقة هي أقصى فترة تحدث فيها الأكسدة وبعدها يحدث التثبع حيث بقي اللون الأصفر لم يتغير وكذلك بالنسبة للطيف.

**الكلمات المفتاحية:** تفريغ حاجز العزل الكهربائي، إزالة اللون، التوصيلية الكهربائية، صبغة النيل القرمزية، الأكسدة.