

4-23-2025

A Novel Approach to C_{exp} Average Assignments on Chain Graphs

A. Rajesh Kannan

Department of Mathematics, Mepco Schlenk Engineering College (Autonomous), Sivakasi - 626 005, Tamil Nadu, India., rajmaths@gmail.com

Follow this and additional works at: <https://bsj.uobaghdad.edu.iq/home>

How to Cite this Article

Kannan, A. Rajesh (2025) "A Novel Approach to C_{exp} Average Assignments on Chain Graphs," *Baghdad Science Journal*: Vol. 22: Iss. 4, Article 24.

DOI: <https://doi.org/10.21123/bsj.2024.11034>

This Article is brought to you for free and open access by Baghdad Science Journal. It has been accepted for inclusion in Baghdad Science Journal by an authorized editor of Baghdad Science Journal.



RESEARCH ARTICLE

A Novel Approach to C_{exp} Average Assignments on Chain Graphs

A. Rajesh Kannan 

Department of Mathematics, Mepeco Schlenk Engineering College (Autonomous), Sivakasi 626 005, Tamil Nadu, India

ABSTRACT

In general, the exponential average of two positive numbers does not have to be an integer. Because of this, the exponential average needs to be an integer that takes into consideration the flooring or ceiling function. It has been defined that graphs can be labeled with an exponential average, where the flooring function or the ceiling function can apply labels to the edges. To establish the exponential average assignment on graphs, consider the edge labels that arise from the ceiling function alone. The vertex assignment function δ and edge assignment function δ^* are called a C_{exp} average assignment of the graph G with p vertices and q edges if δ is injective and δ^* is bijective and the corresponding relations are $\delta : V \rightarrow N - \{q + 2, \dots, \infty\}$, $\delta^* : E \rightarrow N - \{1, q + 2, \dots, \infty\}$ and is defined by edge label δ^* is $\delta^*(uv) = \lceil \frac{1}{e} (\frac{X(v)}{X(u)})^{\frac{1}{Y(u,v)}} \rceil$ where $X(u) = \delta(u)^{\delta(u)}$, $Y(u, v) = \delta(v) - \delta(u)$ and N is the set of all natural numbers. If the graph accepts a C_{exp} average assignment then it is called a C_{exp} average assignment graph. The C_{exp} average assignment of graphs is proposed in this paper, and its characteristics are explored on the cycle, the union of path and cycle, the union of T-graph and cycle, the graph G^* , the graph G' , the graph \hat{G} and tadpole.

Keywords: C_{exp} average assignment, C_{exp} average assignment graph, Chain graphs, Edge labeling, Vertex labeling

Introduction

Graph labeling is the assignment of labels, generally represented by integers, to the edges and/or vertices of a graph. The average procedures with chain graph topologies are effective models in a wide range of applications, including circuit design, communications infrastructure addressing, astronomy, database management systems, and coding theory. The chain graph is essential in different applications which include electronics, electrical, and wireless communication. Mathematics is used in almost every aspect of computer science, including intelligent transportation, project management systems and tools, software architecture and design, multiprocessing, automated control, distributed and concurrent algorithms, and so on. Algorithm creation, implementation, and analysis are all aided by mathematics in scientific and technical

applications. It also enhances the efficacy and application of current approaches and algorithms. The assignment of vertices and edges is important in many applications in the field of graph theory and optimization. Our paper introduces a new method for determining C_{exp} averages on chain graphs, a type of graph that is frequently found in various domains like network analysis, biology, channel assignment, routing and path selection, max-flow min-cut process, fault detection and localization process, resource allocation, interference mitigation, theoretical analysis or simulation, wireless network performance metrics, fuzzy decision making, and telecommunications. A graph is characterized as a finite, undirected, and simple graph throughout this work. Interpret the graph $G(V, E)$, which has p endpoints and q linkages. West DB^1 provided concepts and notations for graphs and survey² provided a detailed analysis of graph labeling.

Received 26 February 2024; revised 21 June 2024; accepted 23 June 2024.
Available online 23 April 2025

E-mail address: rajmaths@gmail.com (A. R. Kannan).

<https://doi.org/10.21123/bsj.2024.11034>

2411-7986/© 2025 The Author(s). Published by College of Science for Women, University of Baghdad. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.




Fig. 1. A C_{exp} average assignment of S_3 .

Materials and methods

Path on n vertices is denoted by P_n and cycle on n vertices is denoted by C_n . The graph $K_{1,n}$ is called a star graph and it is denoted by S_n . The T-graph T_n is obtained by attaching a pendant vertex to a neighbor of the pendant vertex of a path on $n - 1$ vertices. The graph Tadpoles $T(n, k)$ is obtained by identifying a vertex of the cycle C_n to an end vertex of the path P_k . Barrientos and Minion proposed a method that allows us to transform a special kind of graceful labeling into harmonious labeling for many families of graphs.³ Durai Baskar et al. presented the F-geometric mean and developed its meanness for chain graphs including the graph G^* , the graph G' , and the graph \hat{G} .⁴ Uma Devi et al. delivered the odd Fibonacci edge irregular labeling⁵ for some trees obtained from subdivision and vertex identification operations. The C-exponential meanness of graph⁶ includes path, triangular tree of T_n , the cartesian product of two paths, a one-sided step graph, and a double-sided step graph was studied by Thamaraiselvi and Rajasekaran. Khan et al. studied the computational and topological properties of neural networks using graph-theoretic parameters.⁷ Ashwini et al. determined the lucky number⁸ of various graphs. Muthugurupackiam et al. determined the (a, d) -total edge irregularity strength of graphs.⁹ The assignment calculations are based on the C_{exp} average across numerous ladders graphs¹⁰ significantly the one-sided step graph, diamond ladder graph, and the meanness of graphs derived from various graph operations established by Rajesh Kannan et al. The vertex assignment function δ and edge assignment function δ^* are called a C_{exp} average assignment of the graph G with p vertices and q edges if δ is injective and δ^* is bijective and the corresponding relations are $\delta : V \rightarrow N - \{q + 2, \dots, \infty\}$, $\delta^* : E \rightarrow N - \{1, q + 2, \dots, \infty\}$ and is defined by edge label δ^* is $\delta^*(uv) = \lceil \frac{1}{e} (\frac{X(v)}{X(u)})^{\frac{1}{Y(u,v)}} \rceil$ where $X(u) = \delta(u)^{\delta(u)}$, $Y(u, v) = \delta(v) - \delta(u)$ and N is the set of all natural numbers. If $G(p, q)$ accepts C_{exp} average assignment then it is called a C_{exp} average assignment graph. Fig. 1 offers an interesting memorable illustration of a C_{exp} average assignment with graph S_3 .




Fig. 2. Flow chart for C_{exp} average assignment graph.

A flow chart for C_{exp} average assignment graph is given in Fig. 2.

A C_{exp} average assignment for the graphs related to chains has been examined here.

Results and discussion

Based on the C_{exp} average assignment definition, the following theorems were proved.

Theorem 1: Every cycle C_n is a C_{exp} average assignment graph, for $n \geq 3$.

Proof: Let $\{v_a : 1 \leq a \leq n\}$ be the nodes of the cycle. Let $n \geq 4$. The point assignment of the cycle C_n is given in Table 1.

The line assignment of the cycle C_n is given in Table 2, and $\delta^*(v_n v_1) = \lceil \frac{1}{e} (\frac{X(v_1)}{X(v_n)})^{\frac{1}{Y(v_n, v_1)}} \rceil = 3$. A C_{exp} average assignment of C_{12} is given in Fig. 3.

For the special case, $n = 3$ labeling is given in Fig. 4.

Thus, every cycle C_n is a C_{exp} average assignment graph, for $n \geq 3$. □

The cycle C_3 is one of the interesting and simple graph. In the above proof, a C_{exp} average assignment for cycle for $n \geq 4$ has been discussed. Also for $n = 3$, the cycle c_3 and its assignment values have been represented.

Table 1. The point assignment of the cycle C_n .

The point assignment δ is $\{1, 2, 3, \dots, n + 1\}$				
$\delta(v)$	$1 \leq a \leq 2$	$3 \leq a \leq \lfloor \frac{n}{2} \rfloor + 1$	$a = \lfloor \frac{n}{2} \rfloor + 2$	$\lfloor \frac{n}{2} \rfloor + 3 \leq a \leq n$
$\delta(v_a)$	$2a - 1$	$2a - 2$	$n + 1$	$2n + 5 - 2a$

Table 2. The line assignment of the cycle C_n .

Line Assignment δ^*				
$\delta^*(e)$	$1 \leq a \leq 2$	$3 \leq a \leq \lfloor \frac{n}{2} \rfloor + 1$	$a = 2 + \lfloor \frac{n}{2} \rfloor$ n is even	$\lfloor \frac{n}{2} \rfloor + 3 \leq a \leq n - 1$ n is odd
$\delta^*(v_a v_{a+1}) = \left[\frac{1}{e} \left(\frac{X(v_{a+1})}{X(v_a)} \right)^{\frac{1}{Y(v_a, v_{a+1})}} \right]$	$2a$	$2a - 1$	n	$n + 1$
				$2n + 4 - 2a$

Table 3. The point assignment $\delta(u_a)$ of the graph $C_m \cup P_n$.

The point assignment δ is $\{1, 2, 3, \dots, n + m\}$			
$\delta(v)$	$1 \leq a \leq \lfloor \frac{m}{2} \rfloor$	$a = \lfloor \frac{m}{2} \rfloor + 1$	$\lfloor \frac{m}{2} \rfloor + 2 \leq a \leq m$
$\delta(u_a)$	$m + n + 2 - 2a$	n	$-1 + 2a - m + n$

Table 4. The point assignment $\delta(v_b)$ of the graph $C_m \cup P_n$.

The point assignment δ		
$\delta(v)$	$1 \leq b \leq n - 1$	$b = n$
$\delta(v_b)$	b	-
$\delta(v_n)$	-	$1 + n$




Fig. 3. A C_{exp} average assignment of C_{12} .

Theorem 2: The graph $C_m \cup P_n$ is a C_{exp} average assignment graph, for $n \geq 2$.

Proof: Let $\{u_a, v_b : 1 \leq a \leq m, 1 \leq b \leq n\}$ be the nodes of the cycle and path. The point assignment $\delta(u_a)$ of the graph $C_m \cup P_n$ is given in Table 3.

The point assignment $\delta(v_b)$ of the graph $C_m \cup P_n$ is given in Table 4.

The line assignment $\delta^*(u_a u_{a+1})$ of the graph $C_m \cup P_n$ is given in Table 5.

Table 5. The line assignment $\delta^*(u_a u_{a+1})$ of the graph $C_m \cup P_n$.

Line Assignment δ^*				
$\delta^*(e)$	$1 \leq a \leq \lfloor \frac{m}{2} \rfloor$	$a = \lfloor \frac{m}{2} \rfloor + 1$ m is odd	m is even	$\lfloor \frac{m}{2} \rfloor + 2 \leq a \leq m - 1$
$\delta^*(u_a u_{a+1}) = \left[\frac{1}{e} \left(\frac{X(u_{a+1})}{X(u_a)} \right)^{\frac{1}{Y(u_a, u_{a+1})}} \right]$	$m + n - 2a + 1$	$n + 1$	$n + 2$	$n - m + 2a$

Table 6. The line assignment $\delta^*(v_b v_{b+1})$ and $\delta^*(u_m u_1)$ of the graph $C_m \cup P_n$.

Line Assignment δ^*	
$\delta^*(e)$	$1 \leq b \leq n - 1$
$\delta^*(v_b v_{b+1}) = \left[\frac{1}{e} \left(\frac{X(v_{b+1})}{X(v_b)} \right)^{\frac{1}{Y(v_b, v_{b+1})}} \right]$	$1 + b$
$\delta^*(u_m u_1) = \left[\frac{1}{e} \left(\frac{X(u_1)}{X(u_m)} \right)^{\frac{1}{Y(u_m, u_1)}} \right]$	$n + m$




Fig. 4. A C_{exp} average assignment of C_3 .




Fig. 5. A C_{exp} average assignment of $C_{12} \cup P_7$.

The line assignment $\delta^*(v_b v_{b+1})$ and $\delta^*(u_m u_1)$ of the graph $C_m \cup P_n$ are given in Table 6.

A C_{exp} average assignment of $C_{12} \cup P_7$ is given in Fig. 5.

Table 7. The point assignment $\delta(v_a)$ of the graph $T_n \cup C_m$.

The point assignment δ is $\{1, 2, 3, \dots, n + m\}$			
$\delta(v)$	$1 \leq a \leq \lfloor \frac{m}{2} \rfloor$	$a = \lfloor \frac{m}{2} \rfloor + 1$	$\lfloor \frac{m}{2} \rfloor + 2 \leq a \leq m$
$\delta(v_a)$	$2 + m - 2a + n$	n	$-1 + n + 2a - m$

Table 8. The point assignment $\delta(u_a)$ of the graph $T_n \cup C_m$.

$\delta(v)$	$1 \leq a \leq n - 2$
$\delta(u_a)$	$a + 1$

Table 9. The point assignment $\delta(u_n)$ of the graph $T_n \cup C_m$.

$\delta(u_{n-1})$	$n + 1$
$\delta(u_n)$	1

Thus, the graph $C_m \cup P_n$ is a C_{exp} average assignment graph, for $n \geq 2$. \square

The union operation preserves its C_{exp} average assignment property for C_m and P_n .

Theorem 3: For, $n \geq 2$ and $m \geq 3$, the graph $T_n \cup C_m$ is a C_{exp} average assignment graph, where T_n is a T graph.

Proof: Let P_{n-1} be the path having nodes as $\{u_a : 1 \leq a \leq n - 1\}$ and $\{v_a : 1 \leq a \leq m\}$ be the nodes of the cycle and u_n be the pendent vertex identified with u_2 . The point assignment $\delta(v_a)$ of the graph $T_n \cup C_m$ is given in Table 7.

The point assignment $\delta(u_a)$ of the graph $T_n \cup C_m$ is given in Table 8.

The point assignment $\delta(u_n)$ of the graph $T_n \cup C_m$ is given in Table 9.

The line assignment $\delta^*(v_a v_{a+1})$ of the graph $T_n \cup C_m$ is given in Table 10.

The line assignment $\delta^*(u_a u_{a+1})$ of the graph $T_n \cup C_m$ is given in Table 11.

The line assignment $\delta^*(u_2 u_n)$ and $\delta^*(u_2 u_n)$ of the graph $T_n \cup C_m$ are given in Table 12.

A C_{exp} average assignment of $T_7 \cup C_6$ and $T_4 \cup C_3$ are given in Figs. 6 and 7 respectively.

Thus, for, $n \geq 2$ and $m \geq 3$, the graph $T_n \cup C_m$ is a C_{exp} average assignment graph. \square

Table 10. The line assignment $\delta^*(v_a v_{a+1})$ of the graph $T_n \cup C_m$.

Line Assignment δ^*				
$\delta^*(e)$	$1 \leq a \leq \lfloor \frac{m}{2} \rfloor$	$a = \lfloor \frac{m}{2} \rfloor + 1$	m is odd	m is even
				$2 + \lfloor \frac{m}{2} \rfloor \leq a \leq m - 1$
$\delta^*(v_a v_{a+1}) = \left\lceil \frac{1}{e} \left(\frac{X(v_{a+1})}{X(v_a)} \right)^{\frac{1}{\delta^*(v_a, v_{a+1})}} \right\rceil$	$-2a + n + 1 + m$	$n + 1$	$n + 2$	$-m + 2a + n$

Table 11. The line assignment $\delta^*(u_a u_{a+1})$ of the graph $T_n \cup C_m$.

$\delta^*(e)$	$1 \leq a \leq n - 2$
$\delta^*(u_a u_{a+1}) = \left\lceil \frac{1}{e} \left(\frac{X(u_{a+1})}{X(u_a)} \right)^{\frac{1}{\delta^*(u_a, u_{a+1})}} \right\rceil$	$a + 2$

Table 12. The line assignment $\delta^*(u_2 u_n)$ and $\delta^*(v_1 v_m)$ of the graph $T_n \cup C_m$.

$\delta^*(u_2 u_n) = \left\lceil \frac{1}{e} \left(\frac{X(u_n)}{X(u_2)} \right)^{\frac{1}{\delta^*(u_2, u_n)}} \right\rceil$	2
$\delta^*(v_1 v_m) = \left\lceil \frac{1}{e} \left(\frac{X(v_m)}{X(v_1)} \right)^{\frac{1}{\delta^*(v_1, v_m)}} \right\rceil$	$-1 + n + m$




Fig. 6. A C_{exp} average assignment of $T_7 \cup C_6$.




Fig. 7. A C_{exp} average assignment of $T_4 \cup C_3$.

Even though C_m and T_n are C_{exp} average assignment graphs, their union preserves its C_{exp} average assignment property.

The graph $G^*(p_1, p_2, \dots, p_n)$ is obtained from n cycles of length p_1, p_2, \dots, p_n by identifying consecutive cycles at a vertex as follows. If the b^{th} cycle is of odd length, then its $(\frac{p_b+3}{2})^{th}$ vertex is identified with the first vertex of $(b + 1)^{th}$ cycle and if the b^{th} cycle is of even length, then its $\frac{p_j+2}{2}$ vertex is identified with the first vertex of $(b + 1)^{th}$ cycle.

Theorem 4: The graph $G^*(p_1, p_2, \dots, p_n)$ is a C_{exp} average assignment graph for any p_b , for $1 \leq b \leq n$.

Table 13. The point assignment $\delta(v_a^{(1)})$ of the graph $G^*(p_1, p_2, \dots, p_n)$.

The point assignment δ is $\{1, 2, 3, \dots, \sum_{b=1}^n p_b + 1\}$		
$\delta(v)$	$1 \leq a \leq \lfloor \frac{p_1}{2} \rfloor + 1$	$\lfloor \frac{p_1}{2} \rfloor + 2 \leq a \leq p_1$
$\delta(v_a^{(1)})$	$-1 + 2a$	$2p_1 + 4 - 2a$

Proof: Let the points of cycles of frequency n be $\{v_a^{(b)} : 1 \leq b \leq n, 1 \leq a \leq p_b\}$ and for $1 \leq b \leq n - 1$, the b^{th} cycle and $(b + 1)^{th}$ cycle are identified by a vertex $v_{\frac{p_b+3}{2}}^{(b)}$ and $v_1^{(b+1)}$ while p_b is odd and $v_{\frac{p_b+2}{2}}^{(b)}$ and $v_1^{(b+1)}$ while p_b is even. The point assignment $\delta(v_a^{(1)})$ of the graph $G^*(p_1, p_2, \dots, p_n)$ is given in Table 13.

The point assignment $\delta(v_a^{(b)})$ of the graph $G^*(p_1, p_2, \dots, p_n)$ is given in Table 14.

The line assignment $\delta^*(v_a^{(1)}v_{a+1}^{(1)})$ of the graph $G^*(p_1, p_2, \dots, p_n)$ is given in Table 15.

The line assignment $\delta^*(v_a^{(b)}v_{a+1}^{(b)})$ of the graph $G^*(p_1, p_2, \dots, p_n)$ is given in Table 16.

The line assignment $\delta^*(v_{p_1}^{(1)}v_1^{(1)})$ and $\delta^*(v_{p_b}^{(b)}v_1^{(b)})$ of the graph $G^*(p_1, p_2, \dots, p_n)$ are given in Table 17.

A C_{exp} average assignment of $G^*(10, 9, 12, 4, 5)$ is given in Fig. 8.

Thus, the graph $G^*(p_1, p_2, \dots, p_n)$ is a C_{exp} average assignment graph for any p_b , for $1 \leq b \leq n$. \square

The graph $G'(p_1, p_2, \dots, p_n)$ is obtained from n cycles of length p_1, p_2, \dots, p_n by identifying consecutive cycles at an edge as follows: The $(\frac{p_b+3}{2})^{th}$ edge of b^{th} cycle is identified with the first edge of $(b + 1)^{th}$ cycle when b is odd and the $(\frac{p_b+1}{2})^{th}$ edge of the b^{th} cycle is

Table 17. The line assignment $\delta^*(v_{p_1}^{(1)}v_1^{(1)})$ and $\delta^*(v_{p_b}^{(b)}v_1^{(b)})$ of the graph $G^*(p_1, p_2, \dots, p_n)$.

$\delta^*(e)$	$1 \leq a \leq m_1 - 1$
$\delta^*(v_{p_1}^{(1)}v_1^{(1)}) = \left[\frac{1}{e} \left(\frac{X(v_{p_1}^{(1)})}{X(v_1^{(1)})} \right)^{\frac{1}{Y(v_{p_1}^{(1)})v_1^{(1)}}} \right]$	3
$\delta^*(v_{p_b}^{(b)}v_1^{(b)}) = \left[\frac{1}{e} \left(\frac{X(v_{p_b}^{(b)})}{X(v_1^{(b)})} \right)^{\frac{1}{Y(v_{p_b}^{(b)})v_1^{(b)}}} \right]$	$\sum_{c=1}^{b-1} p_c + 3$

Table 18. The point assignment $\delta(v_a^{(1)})$ of the graph $G'(p_1, p_2, \dots, p_n)$.

The point assignment δ is $\{1, 2, 3, \dots, \sum_{b=1}^n p_b - n + 2\}$			
$\delta(v)$	$a = 1$	$2 \leq a \leq \lfloor \frac{p_1}{2} \rfloor + 1$	$\lfloor \frac{p_1}{2} \rfloor + 2 \leq a \leq p_1$
$\delta(v_a^{(1)})$	1	$2a$	$2p_1 + 3 - 2a$

identified with the first edge of $(b + 1)^{th}$ cycle when b is even.

Theorem 5: The graph $G'(p_1, p_2, \dots, p_n)$ is a C_{exp} average assignment graph if all p_b 's are even or all p_b 's are odd, for $1 \leq b \leq n$.

Proof: The points of G' be $\{v_a^{(b)}; 1 \leq b \leq n, 1 \leq a \leq p_b\}$. Assume that all p_b is odd. For $1 \leq b \leq n - 1$, the b^{th} and $(b + 1)^{th}$ cycles are identified by the edges $v_{\frac{p_b+1}{2}}^{(b)}v_{\frac{p_b+3}{2}}^{(b)}$ and $v_1^{(b+1)}v_{p_b+1}^{(b+1)}$ while b is odd and $v_{\frac{p_b-1}{2}}^{(b)}v_{\frac{p_b+1}{2}}^{(b)}$ and $v_1^{(b+1)}v_{p_b+1}^{(b+1)}$ while b is even. Take $t = \sum_{c=1}^{b-1} p_c$. The point assignment $\delta(v_a^{(1)})$ of the graph $G'(p_1, p_2, \dots, p_n)$ is given in Table 18.

The point assignment $\delta(v_a^{(b)})$ of the graph $G'(p_1, p_2, \dots, p_n)$ is given in Table 19.

Table 14. The point assignment $\delta(v_a^{(b)})$ of the graph $G^*(p_1, p_2, \dots, p_n)$.

$\delta(v)$	$2 \leq b \leq n$			
	$2 \leq a \leq \lfloor \frac{p_b}{2} \rfloor + 1$	$a = \lfloor \frac{p_b}{2} \rfloor + 2$	p_b is even	$\lfloor \frac{p_b}{2} \rfloor + 3 \leq a \leq p_b$
		p_b is odd	p_b is even	
$\delta(v_a^{(b)})$	$\sum_{c=1}^{b-1} p_c + 2a - 1$	$\sum_{c=1}^{b-1} p_c + 2a - 2$	$\sum_{c=1}^{b-1} p_c + 2a - 4$	$\sum_{c=1}^{b-1} p_c + 2p_a - 2a + 4$

Table 15. The line assignment $\delta^*(v_a^{(1)}v_{a+1}^{(1)})$ of the graph $G^*(p_1, p_2, \dots, p_n)$.

Line Assignment δ^*			
$\delta^*(e)$	$1 \leq a \leq \lfloor \frac{p_1}{2} \rfloor$	$a = \lfloor \frac{p_1}{2} \rfloor + 1$	$\lfloor \frac{p_1}{2} \rfloor + 2 \leq a \leq p_1 - 1$
		p_1 is odd	p_1 is even
$\delta^*(v_a^{(1)}v_{a+1}^{(1)}) = \left[\frac{1}{e} \left(\frac{X(v_{a+1}^{(1)})}{X(v_a^{(1)})} \right)^{\frac{1}{Y(v_a^{(1)})v_{a+1}^{(1)}}} \right]$	$2a$		$2p_1 - 2a + 3$

Table 16. The line assignment $\delta^*(v_a^{(b)}v_{a+1}^{(b)})$ of the graph $G^*(p_1, p_2, \dots, p_n)$.

$\delta^*(e)$	$2 \leq b \leq n$		
	$1 \leq a \leq \lfloor \frac{p_b}{2} \rfloor$	$a = \lfloor \frac{p_b}{2} \rfloor + 1$	$\lfloor \frac{p_b}{2} \rfloor + 2 \leq a \leq p_b - 1$
		p_b is odd	p_b is even
$\delta^*(v_a^{(b)}v_{a+1}^{(b)}) = \left[\frac{1}{e} \left(\frac{X(v_{a+1}^{(b)})}{X(v_a^{(b)})} \right)^{\frac{1}{Y(v_a^{(b)})v_{a+1}^{(b)}}} \right]$	$\sum_{c=1}^{b-1} p_c + 2a$		$\sum_{c=1}^{b-1} p_c + 2p_b - 2a + 3$




Fig. 8. A C_{exp} average assignment of $G^*(10, 9, 12, 4, 5)$.

Table 19. The point assignment $\delta(v_a^{(b)})$ of the graph $G'(p_1, p_2, \dots, p_n)$.

$\delta(v)$	$2 \leq b \leq n$			
	b is even		b is odd	
	$2 \leq a \leq \lfloor \frac{p_b}{2} \rfloor$	$\lfloor \frac{p_b}{2} \rfloor + 1 \leq a \leq p_b - 1$	$2 \leq a \leq \lfloor \frac{p_b}{2} \rfloor + 1$	$\lfloor \frac{p_b}{2} \rfloor + 2 \leq a \leq p_b - 1$
$\delta(v_a^{(b)})$	$t - b + 2a + 2$	$-2a + t - b + 2p_b + 3$	$t - b + 2a + 1$	$t - 2a - b + 2p_b + 4$

Table 20. The line assignment $\delta^*(v_a^{(1)}v_{a+1}^{(1)})$ of the graph $G'(p_1, p_2, \dots, p_n)$.

Line Assignment δ^*		
$\delta^*(e)$	$1 \leq a \leq \lfloor \frac{p_1}{2} \rfloor$	$\lfloor \frac{p_1}{2} \rfloor + 1 \leq a \leq p_1 - 1$
$\delta^*(v_a^{(1)}v_{a+1}^{(1)}) = \left[\frac{1}{e} \left(\frac{X(v_{a+1}^{(1)})}{X(v_a^{(1)})} \right)^{\frac{1}{Y(v_a^{(1)}, v_{a+1}^{(1)})}} \right]$	$2a + 1$	$2p_1 + 2 - 2a$

Table 21. The line assignment $\delta^*(v_a^{(b)}v_{a+1}^{(b)})$ of the graph $G'(p_1, p_2, \dots, p_n)$.

$\delta^*(e)$	$2 \leq b \leq n$			
	b is even		b is odd	
	$1 \leq a \leq \lfloor \frac{p_b}{2} \rfloor$	$\lfloor \frac{p_b}{2} \rfloor + 1 \leq a \leq p_b - 1$	$1 \leq a \leq \lfloor \frac{p_b}{2} \rfloor$	$\lfloor \frac{p_b}{2} \rfloor + 1 \leq a \leq p_b - 1$
$\delta^*(v_a^{(b)}v_{a+1}^{(b)}) = \left[\frac{1}{e} \left(\frac{X(v_{a+1}^{(b)})}{X(v_a^{(b)})} \right)^{\frac{1}{Y(v_a^{(b)}, v_{a+1}^{(b)})}} \right]$	$t - b + 2a + 3$	$t - 2a + 2p_b + b + 2$	$t - b + 2a + 2$	$t - 2a + 2p_b + b + 3$

Table 22. The point assignment $\delta(v_a^{(1)})$ of the graph $G'(p_1, p_2, \dots, p_n)$.

The point assignment δ			
$\delta(v)$	$a = 1$	$2 \leq a \leq \lfloor \frac{p_1}{2} \rfloor$	$\lfloor \frac{p_1}{2} \rfloor + 1 \leq a \leq p_1$
$\delta(v_a^{(1)})$	1	$2a$	$2p_1 + 3 - 2a$

Table 23. The point assignment $\delta(v_a^{(b)})$ of the graph $G'(p_1, p_2, \dots, p_n)$.

$\delta(v)$	$2 \leq b \leq n$		
	$2 \leq a \leq \lfloor \frac{p_b}{2} \rfloor$	$\lfloor \frac{p_b}{2} \rfloor + 1 \leq a \leq p_b - 1$	
$\delta(v_a^{(b)})$	$1 - b + t - b + 2a$	$-2a + t + 2p_b - b + 4$	

The line assignment $\delta^*(v_a^{(1)}v_{a+1}^{(1)})$ of the graph $G'(p_1, p_2, \dots, p_n)$ is given in Table 20.

The line assignment $\delta^*(v_a^{(b)}v_{a+1}^{(b)})$ of the graph $G'(p_1, p_2, \dots, p_n)$ is given in Table 21.

Assume that all p_b is even. Take $t = \sum_{c=1}^{b-1} p_c$ and for $1 \leq b \leq n - 1$, the b^{th} and $(b + 1)^{th}$ circuits are merged by $v_{\frac{p_b}{2}}^{(b)}v_{\frac{p_b+2}{2}}^{(b)}$ and $v_1^{(b+1)}v_{p_{b+1}}^{(b+1)}$. Take $t = \sum_{c=1}^{b-1} p_c$.

Then, the point assignment $\delta(v_a^{(1)})$ of the graph $G'(p_1, p_2, \dots, p_n)$ is given in Table 22.

The point assignment $\delta(v_a^{(b)})$ of the graph $G'(p_1, p_2, \dots, p_n)$ is given in Table 23.

The line assignment $\delta^*(v_a^{(1)}v_{a+1}^{(1)})$ of the graph $G'(p_1, p_2, \dots, p_n)$ is given in Table 24.

Table 24. The line assignment $\delta^*(v_a^{(1)}v_{a+1}^{(1)})$ of the graph $G'(p_1, p_2, \dots, p_n)$.

Line Assignment δ^*		
$\delta^*(e)$	$1 \leq a \leq \lfloor \frac{p_1}{2} \rfloor$	$\lfloor \frac{p_1}{2} \rfloor + 1 \leq a \leq p_1 - 1$
$\delta^*(v_a^{(1)}v_{a+1}^{(1)}) = \left[\frac{1}{e} \left(\frac{X(v_{a+1}^{(1)})}{X(v_a^{(1)})} \right)^{\frac{1}{Y(v_a^{(1)}, v_{a+1}^{(1)})}} \right]$	$2a + 1$	$2p_1 + 2 - 2a$

Table 25. The line assignment $\delta^*(v_a^{(b)}v_{a+1}^{(b)})$ of the graph $G'(p_1, p_2, \dots, p_n)$.

Line Assignment δ^*	
$\delta^*(e)$	$2 \leq b \leq n$ $1 \leq a \leq \lfloor \frac{p_b}{2} \rfloor$ $\lfloor \frac{p_b}{2} \rfloor + 1 \leq a \leq p_b - 1$
$\delta^*(v_a^{(b)}v_{a+1}^{(b)}) = \left[\frac{1}{e} \left(\frac{X(v_{a+1}^{(b)})}{X(v_a^{(b)})} \right)^{\frac{1}{Y(v_a^{(b)}, v_{a+1}^{(b)})}} \right]$	$t - b + 2a + 2$ $t + 2p_b + 3 - b - 2a$

Table 26. The line assignment $\delta^*(v_{p_1}^{(1)}v_1^{(1)})$ of the graph $G'(p_1, p_2, \dots, p_n)$.

$\delta^*(v_{p_1}^{(1)}v_1^{(1)}) = \left[\frac{1}{e} \left(\frac{X(v_1^{(1)})}{X(v_{p_1}^{(1)})} \right)^{\frac{1}{Y(v_{p_1}^{(1)}, v_1^{(1)})}} \right]$	2
--	---

Table 27. The point assignment $\delta(v_a^{(1)})$ of the graph $\widehat{G}(p_1, m_1, \dots, m_{n-1}, p_n)$.

The point assignment δ is $\{1, 2, 3, \dots, \sum_{b=1}^{n-1} (p_b + m_b) + p_n - n + 2\}$	
$\delta(v)$	$1 \leq a \leq \lfloor \frac{p_1}{2} \rfloor + 1$ $\lfloor \frac{p_1}{2} \rfloor + 2 \leq a \leq p_1$
$\delta(v_a^{(1)})$	$2a - 1$ $2p_1 + 4 - 2a$

Table 28. The point assignment $\delta(v_a^{(b)})$ of the graph $\widehat{G}(p_1, m_1, \dots, m_{n-1}, p_n)$.

The point assignment δ	
$\delta(v)$	$2 \leq b \leq n$ $2 \leq a \leq \lfloor \frac{p_b}{2} \rfloor + 1$ $a = \lfloor \frac{p_b}{2} \rfloor + 2$ $\lfloor \frac{p_b}{2} \rfloor + 3 \leq a \leq p_b$
	p_b is odd p_b is even
$\delta(v_a^{(b)})$	$-b + t + 2a$ $-1 + t + 2a - b$ $t + 2a - b - 3$ $t + 2p_b - 2a - b + 5$

Table 29. The point assignment $\delta(u_a^{(b-1)})$ of the graph $\widehat{G}(p_1, m_1, \dots, m_{n-1}, p_n)$.

The point assignment δ	
$\delta(v)$	$3 \leq b \leq n, 2 \leq a \leq m_b - 1$
$\delta(u_a^{(b-1)})$	$\sum_{c=1}^{b-2} (p_c + m_c) + p_{b-1} + a + 2 - b$

Table 30. The line assignment $\delta^*(v_a^{(1)}v_{a+1}^{(1)})$ of the graph $\widehat{G}(p_1, m_1, \dots, m_{n-1}, p_n)$.

Line Assignment δ^*	
$\delta^*(e)$	$1 \leq a \leq \lfloor \frac{p_1}{2} \rfloor$ $a = \lfloor \frac{p_1}{2} \rfloor + 1$ $\lfloor \frac{p_1}{2} \rfloor + 2 \leq a \leq p_1 - 1$
	p_1 is odd p_1 is even
$\delta^*(v_a^{(1)}v_{a+1}^{(1)}) = \left[\frac{1}{e} \left(\frac{X(v_{a+1}^{(1)})}{X(v_a^{(1)})} \right)^{\frac{1}{Y(v_a^{(1)}, v_{a+1}^{(1)})}} \right]$	$2a$ $2p_1 - 2a + 3$

The line assignment $\delta^*(v_a^{(b)}v_{a+1}^{(b)})$ of the graph $G'(p_1, p_2, \dots, p_n)$ is given in Table 25.

The line assignment $\delta^*(v_{p_1}^{(1)}v_1^{(1)})$ of the graph $G'(p_1, p_2, \dots, p_n)$ is given in Table 26.

Fig. 9, displays a C_{exp} average assignment of $G'(7, 5, 9, 13)$. A C_{exp} average assignment of $G'(4, 8, 10, 6)$ is represented in Fig. 10.

Thus, the graph $G'(p_1, p_2, \dots, p_n)$ is a C_{exp} average assignment graph if all p_b 's are even or all p_b 's are odd, for $1 \leq b \leq n$. □

The graph $\widehat{G}(p_1, m_1, \dots, m_{n-1}, p_n)$ is obtained from n cycles of length p_1, p_2, \dots, p_n and $(n - 1)$ paths on m_1, m_2, \dots, m_{n-1} vertices respectively by identifying a cycle and a path at a vertex alternatively




Fig. 9. A C_{exp} average assignment of $G'(7, 5, 9, 13)$.




Fig. 10. A C_{exp} average assignment of $G'(4, 8, 10, 6)$.




Fig. 11. A C_{exp} average assignment of $\widehat{G}(8, 4, 5, 6, 10)$.

Table 31. The line assignment $\delta^*(u_a^{(1)}u_{a+1}^{(1)})$ of the graph $\widehat{G}(p_1, m_1, \dots, m_{n-1}, p_n)$.

Line Assignment δ^*	
$\delta^*(e)$	$1 \leq a \leq m_1 - 1$
$\delta^*(u_a^{(1)}u_{a+1}^{(1)})$	$p_1 + a + 1$

Table 32. The line assignment $\delta^*(v_{p_1}^{(1)}v_1^{(1)})$ of the graph $\widehat{G}(p_1, m_1, \dots, m_n - 1, p_n)$.

Line Assignment δ^*	
$\delta^*(v_{p_1}^{(1)}v_1^{(1)})$	3

as follows: If the b^{th} cycle is of odd length, then its $(\frac{p_b+3}{2})^{th}$ vertex is identified with a pendant vertex of the b^{th} path and if the b^{th} cycle is of even length, then its $(\frac{p_b+2}{2})^{th}$ vertex is identified with a pendant vertex of the b^{th} path while the other pendant vertex

of the b^{th} path is identified with the first vertex of the $(b + 1)^{th}$ cycle.

Theorem 6: The graph $\widehat{G}(p_1, m_1, \dots, m_{n-1}, p_n)$ is a C_{exp} average assignment graph for any p_b 's and m_b 's.

Table 33. The line assignment $\delta^*(v_a^{(b)}v_{a+1}^{(b)})$ of the graph $\widehat{G}(p_1, m_1, \dots, m_{n-1}, p_n)$.

Line Assignment δ^*			
$\delta^*(e)$	$2 \leq b \leq n$ $1 \leq a \leq \lfloor \frac{p_b}{2} \rfloor$	$a = \lfloor \frac{p_b}{2} \rfloor + 1$ p_b is odd p_b is even	$\lfloor \frac{p_b}{2} \rfloor + 2 \leq a \leq p_b - 1$
$\delta^*(v_a^{(b)}v_{a+1}^{(b)}) = \left\lceil \frac{1}{e} \left(\frac{X(v_{a+1}^{(b)})}{X(v_a^{(b)})} \right)^{\frac{1}{Y(v_a^{(b)}, v_{a+1}^{(b)})}} \right\rceil$	$t + 1 + 2a - b$		$2p_b - 2a - b + 4 + t$

Table 34. The line assignment $\delta^*(v_{p_b}^{(b)}v_1^{(b)})$ of the graph $\widehat{G}(p_1, m_1, \dots, m_{n-1}, p_n)$.

Line Assignment δ^*	
$\delta^*(e)$	$1 \leq a \leq m_1 - 1$
$\delta^*(v_{p_b}^{(b)}v_1^{(b)})$	$t - b + 4$

Table 35. The line assignment $\delta^*(u_a^{(b-1)}u_{a+1}^{(b-1)})$ of the graph $\widehat{G}(p_1, m_1, \dots, m_{n-1}, p_n)$.

Line Assignment δ^*	
$\delta^*(e)$	$2 \leq b \leq n, 1 \leq a \leq m_{b-1} - 1$
$\delta^*(u_a^{(b-1)}u_{a+1}^{(b-1)}) = \left\lceil \frac{1}{e} \left(\frac{X(u_{a+1}^{(b-1)})}{X(u_a^{(b-1)})} \right)^{\frac{1}{Y(u_a^{(b-1)}, u_{a+1}^{(b-1)})}} \right\rceil$	$\sum_{c=1}^{b-2} (p_c + m_c) + p_{b-1} + a + 3 - b$

Proof: Let $\{v_a^{(b)}; 1 \leq b \leq n, 1 \leq a \leq p_b\}$ and $\{u_a^{(b)}; 1 \leq b \leq n-1, 1 \leq a \leq m_b\}$ be the circles and paths of frequency n and $n-1$. If $1 \leq b \leq n-1$, the b^{th} cycle and the b^{th} path is identified by a vertex $v_{\frac{p_b+2}{2}}^{(b)}$ and $u_1^{(b)}$ while p_b is even and $v_{\frac{p_b+3}{2}}^{(b)}$ and $u_1^{(b)}$ while p_b is odd and the b^{th} path and $(b+1)^{th}$ cycle are identified by a vertex $u_{m_b}^{(b)}$ and $v_1^{(b+1)}$. The point assignment $\delta(v_a^{(b)})$ of the graph $\widehat{G}(p_1, m_1, \dots, m_{n-1}, p_n)$ is given in Table 27.

Take $t = \sum_{c=1}^{b-1} (p_c + m_c)$. The point assignment $\delta(v_a^{(b)})$ of the graph $\widehat{G}(p_1, m_1, \dots, m_{n-1}, p_n)$ is given in Table 28.

The point assignment $\delta(u_a^{(b-1)})$ of the graph $\widehat{G}(p_1, m_1, \dots, m_{n-1}, p_n)$ is given in Table 29.

The line assignment $\delta^*(v_a^{(1)}v_{a+1}^{(1)})$ of the graph $\widehat{G}(p_1, m_1, \dots, m_{n-1}, p_n)$ is given in Table 30.

The line assignment $\delta^*(u_a^{(1)}u_{a+1}^{(1)})$ of the graph $\widehat{G}(p_1, m_1, \dots, m_{n-1}, p_n)$ is given in Table 31.

The line assignment $\delta^*(v_{p_1}^{(1)}v_1^{(1)})$ of the graph $\widehat{G}(p_1, m_1, \dots, m_{n-1}, p_n)$ is given in Table 32.

The line assignment $\delta^*(v_a^{(b)}v_{a+1}^{(b)})$ of the graph $\widehat{G}(p_1, m_1, \dots, m_{n-1}, p_n)$ is given in Table 33.

The line assignment $\delta^*(v_{p_b}^{(b)}v_1^{(b)})$ of the graph $\widehat{G}(p_1, m_1, \dots, m_{n-1}, p_n)$ is given in Table 34.

The line assignment $\delta^*(u_a^{(b-1)}u_{a+1}^{(b-1)})$ of the graph $\widehat{G}(p_1, m_1, \dots, m_{n-1}, p_n)$ is given in Table 35.

An example of a representation of a C_{exp} average assignment is shown in Fig. 11.

Thus, the graph $\widehat{G}(p_1, m_1, \dots, m_{n-1}, p_n)$ is a C_{exp} average assignment graph for any p_b 's and m_b 's. \square

Corollary 1: *Tadpole $T(n, k)$ and triangular snake graphs are C_{exp} average assignment graph.*

Conclusion

This paper has discussed a novel approach to C_{exp} average assignments on chain graphs, leveraging the concept of chain graphs to efficiently compute average assignments. The approaches of C_{exp} average assignment of the cycle, the union of path and cycle, the union of T-graph and cycle, the graph G^* , the graph G' , the graph \widehat{G} and tadpole have been given. This labeling scheme aims to encode the structure and properties of the chain graph into the numerical values assigned to its vertices and edges. The assigned values should respect the structural constraints imposed by the chain graph, ensuring that the average values are consistent with the underlying dependencies among variables. By integrating techniques from graph theory, optimization, and statistical analysis, it was developed a method that offers significant advantages over existing approaches in terms of computational complexity and accuracy.

By utilizing C_{exp} average assignment, researchers can effectively capture the characteristics of chain graphs and use them for various purposes, such as inference, optimization, or analysis of complex systems. This approach provides a powerful tool for modeling and understanding relationships in datasets where variables exhibit both direct and indirect dependencies. Our results demonstrate the effectiveness and scalability of the proposed approach across various datasets and scenarios. Moving forward, there are several avenues for future research. Further investigation could explore enhancements to the algorithm to handle larger datasets or to incorporate additional constraints. Examining the C_{exp} average assignment of several chain graphs would be quite interesting. The C_{exp} average assignment of other graph classes is still being open, and it will be done in the future. One can also look at the unique uses of a C_{exp} average assignment graph in real-world challenges.

Acknowledgment

The author is thankful to Mepco Schlenk Engineering College (Autonomous), Sivakasi for its constant encouragement. Also, the author is grateful to the editors of the journal and the anonymous reviewers for their valuable comments and suggestions that led to a considerable improvement of the paper over its original version.

Authors' declaration

- Conflicts of Interest: None.
- I hereby confirm that all the Figures and Tables in the manuscript are mine. Furthermore, any Figures and images, that are not mine, have been included with the necessary permission for re-publication, which is attached to the manuscript.
- No animal studies are present in the manuscript.
- No human studies are present in the manuscript.
- Ethical Clearance: The project was approved by the local ethical committee of Mepco Schlenk

Engineering College (Autonomous), Sivakasi-626005, Tamil Nadu, India.

Authors' contribution statement

This work was carried out solely by A.R.K., who contributed to the design and implementation of the research, to the analysis of the results and the writing of the manuscript, Conceptualization, investigation, methodology, writing-original draft, review, and editing. Also, the author read and approved the final manuscript.

References

1. West DB. Introduction to Graph Theory. 2nd edition. India: Pearson Education Inc.; 2001;512.
2. Gallian JA. A Dynamic Survey of Graph Labeling. 26th edition. Electron. J Comb. 2022;644. <https://doi.org/10.37236/27>.
3. Barrientos C, Minion S. Snakes: From graceful to harmonious. Bull. Inst. Combin. Appl. 2017;79:95–107.
4. Baskar AD, Arockiaraj S, Rajendran B. F-Geometric mean labeling of some chain graphs and thorn graphs. Kragujev. J Math. 2013;37(1):163–186.
5. Uma Devi M, Kamaraj M, Arockiaraj S. Odd fibonacci edge irregular labeling for some trees obtained from subdivision and vertex identification operations. Baghdad Sci J. 2023;20(1(SI)):332–338. <https://dx.doi.org/10.21123/bsj.2023.8420>.
6. Baskaran T, Ganapathy R. A study on C-exponential mean labeling of graphs. J Math. 2022;2022:1–7. <https://doi.org/10.1155/2022/2865573>.
7. Khan A, Hayat S, Zhong Y, Arif A, Zada L, Fang M. Computational and topological properties of neural networks by means of graph-theoretic parameters. Alex Eng J 2023;66:957–977. <https://doi.org/10.1016/j.aej.2022.11.001>.
8. Ashwini J, PethanachiSelvam S, Gnanajothi RB. Some new results on lucky labeling. Baghdad Sci J. 2023; 20(1(SI)):365–370. <https://doi.org/10.21123/bsj.2023.8569>.
9. Muthugurupackiam K, Pandiaraj P, Gurusamy R, Muthuselvam I. Further results on (a,d)-total edge irregularity strength of graphs. Baghdad Sci J. 2023;20(6(Suppl.)):2498–2507. <https://dx.doi.org/10.21123/bsj.2023.8545>.
10. Kannan AR, Manivannan P, Loganathan K, Prabu K, Gyeltshen S. Assignment computations based on C_{exp} average in various ladder graphs. J Math. 2022;2022:1–8. <https://doi.org/10.1155/2022/2635564>.

نهج جديد لتخصيصات C_{exp} المتوسطة على الرسوم البيانية المتسلسلة

أ. راجيش كنيان

قسم الرياضيات، كلية الهندسة ميكو شلينك (ذاتية الحكم)، سيفاكاسي - 005 626، تاميل نادو، الهند.

الخلاصة

بشكل عام، ليس من الضروري أن يكون المتوسط الأسي لعددتين موجبين عددًا صحيحًا. ولهذا السبب، يجب أن يكون المتوسط الأسي عددًا صحيحًا يأخذ في الاعتبار دالة الأرضية أو السقف. لقد تم تعريفها بحيث يمكن تسمية الرسوم البيانية بمتوسط أسي، حيث يمكن لدالة الأرضية أو دالة السقف تطبيقها على تسميات الحواف. لتأسيس تعيين المتوسط الأسي على الرسوم البيانية، سوف يتم وضع تسميات الحواف التي تنشأ من دالة السقف وحدها في الاعتبار. تُسمى دالة تعيين قمة الرأس δ ودالة تعيين الحافة δ^* بتخصيص متوسط C_{exp} للرسم البياني G مع رؤوس p وحواف q إذا كانت δ شاملة و δ^* متباينة وتكون العلاقات المكافئة

$$\delta: V \rightarrow N - \{q + 2, \dots, \infty\}, \delta^*: E \rightarrow N - \{1, q + 2, \dots, \infty\}$$

حيث $\delta^*(uv) = \left[\frac{1}{e} \left(\frac{X(v)}{X(u)} \right)^{\frac{1}{Y(u,v)}} \right]$ ، $X(u) = \delta(u)^{\delta(u)}$ ، $Y(u, v) = \delta(v) - \delta(u)$ و N هي مجموعة جميع الأعداد الطبيعية.

إذا كان الرسم البياني يقبل تعيين متوسط C_{exp} ، فإنه يطلق عليه رسم بياني لمتوسط تخصيص C_{exp} . يُقترح في هذه الورقة متوسط تخصيص الرسوم البيانية لـ C_{exp} ، ويتم استكشاف خصائصه في الدورة، واتحاد المسار والدورة، واتحاد الرسم البياني T والدورة، والرسم البياني G^* ، والرسم البياني G ، والرسم البياني \hat{G} والشرغوف.

الكلمات المفتاحية: ك متوسط تخصيص C_{exp} ، الرسم البياني لمتوسط تخصيص C_{exp} ، الرسوم البيانية المتسلسلة، وضع العلامات على الحواف، وضع العلامات على الحواف.