STUDY THE NUTRITIONAL PROPERTIES OF SOME INFANT FORMULA AVAILABLE IN KUFA AND NAJAF LOCAL MARKETS

Mohanad Abbood Alhamid^{*} Riyadh Shamkhi Ali^{**} Abidale Alwan AlTae^{***} ^{*}Researcher/ College of Agriculture- University of Kufa, ^{**}Assistant Professor/ College of Agriculture - University of Kufa, ^{***}Assistant Professor/ College of Food Sciences-Al-Qasim Green University

mohanada.alhamid@student.uokufa.edu.iq

ABSTRACT:

This study was conducted to evaluated nutritional properties of some infant formulas available in the local markets and their effects on the growth and nutritional status of rats and their body organs. This experiment was designed using 42 white male rats (Albino) with initial average weight of 50±5 g, they were divided into 7 groups of 6 rats per group. The experiment continuous for 28 days during which the rats were fed on infant milk formula treatments T_1 , T_2 and T_3 to age less than 6 month (Nactalia1, Dielac1 and Evolac1) respectively, and in treatments T₄, T₅, T₆ feed on infant milk formula for age more than 6 month (Celia2, Dielac2 and Sunny baby2) respectively, and the rest of rats in control treatment fed on standard provender and filter water. The results showed significant (P≤0.05) decrease in final weight rate of rats in T₃ which was 118.7 g recording the lowest weight gained 72.8 g during the experiment period, while treatment T_5 recorded the highest weight increase gained reaching 118.4 g and final weight rate (169.8 g). In control treatment was significantly ($P \le 0.05$) higher in amount food intake compared to all other treatments, while treatments T_2 and T_3 have the higher and lower amount of food intake respectively. On the other hand, it was noted decrease in relative weight of rats' kidneys in treatment T₁, T₂ and T₃ compare with the control treatment, which recorded the highest average of relative weight for rats' kidneys of 1.086 g.

Keyword: Infant formula, Nutritional properties, rats, Food efficiency. دراسة الخصائص التغذوية لبعض خلائط حليب الأطفال المتوفرة في الأسواق المحلية لمدينتي الكوفة والنجف الاشرف مهند عبود مهاوي الحميد * باحث/ كلية الزراعة جامعة الكوفة، ** أستاذ مساعد/ كلية علوم الأغذية-جامعة القاسم الخضراع

الخلاصة:

أجريت هذه الدراسة لتقييم الخواص التغذوية لبعض خلائط حليب الأطفال المتوفرة في الأسواق المحلية ومعرفة تأثيرها على نمو الجرذان وحالتها التغذوية واعضاءها الداخلية. حيث صممت التجربة باستعمال 42 حيوان من ذكور جرذان Albino البيضاء بمعدل وزن ابتدائي 50±5غم. وقسمت الى سبع مجاميع بواقع 6 جرذان لكل مجموعة. واستمرت التجربة لمدة 28 يوم تم خلالها تغذية الجرذان في المعاملات T1، T2و T3 على خلائط حليب الأطفال لعمر اقل من 6 أشهر (Dielaci،Nactalial و Evolaci) وفي المعاملات T1، T2و T3 على خلائط حليب الأطفال لعمر فوق أشهر (Dielaci،Nactalial و Evolaci) وفي المعاملات T1، T5و T6 على خلائط حليب الأطفال لعمر فوق أشهر (Dielaci،Nactalial و Sunnybaby2) على التوالي. وإبقاء تغذية جرذان معاملة السيطرة على العليقة القياسية والماء المفلتر. أظهرت النتائج انخفاض معنوي في معدل الوزن النهائي لجرذان المعاملة T1 على معدل الزيادة والماء المفلتر. أظهرت النتائج انخفاض معنوي في معدل الوزن النهائي الجرذان المعاملة 71 على معدل الزيادة والماء المفلتر. أظهرت النتائج انخفاض معنوي في معدل الوزن النهائي الجرذان المعاملة 71 العلى معدل الزيادة والماء المفلتر. أظهرت النتائج انخفاض معنوي في معدل الوزن النهائي العرذان المعاملة 71 على معدل الزيادة معدل الوزنية المكتسبة حيث بلغت 18.71 غم وبمعدل وزن نهائي 80.00 عم. عمل الوزيادة الوزنية المكتسبة حيث بلغت 18.81 غم وبمعدل وزن نهائي 19.00 غم. كما لوحظ تفوق معاملة السيطرة معنويا من حيث كمية الغذاء المتناول على جميع المعاملات عند مستوى احتمال(0.00). كمية الغذاء المتناول على جميع المعاملات عند مستوى احتمال(10.00). 2003 على الوزن النماي على معاملة السيطرة معنويا من حيث تمام معاملة السيطرة معنويا من حيث المعاملة ترة المعاملات 130 على الموطز تفوق معاملة السيطرة معنويا من حيث كمية الغذاء المتناول على جميع المعاملات عند مستوى احتمال (0.00). 2013 على معاملة السيطرة التي سجلت أعلى معدل لوزن النماي حيث المام على الجرذان في المعاملة السيطرة معنويا من حيث تموم معاملة السيطرة معام التناول في المعاملات 131، معدل فوزن الكلى النسبي حيث بلغ 1.006 غم. 2013 على معاملة السيطرة التي سجلت أعلى معدل لوزن الكلى النسبي حيث بلغ 20.00 غم.

الكلمات المفتاحية: حليب الأطفال ، جرذان ، كفاءة الغذاء ، الخواص التغذوية.

INTRODUCTION

Milk is one of the most important components of the human diet in many parts of the world and the average per capita consumption is of 180 in Iceland and Finland to less than 50 in Japan and China [1]. Milk contains all the nutrients necessary for the physiological functions of the body as well as containing fat, proteins, enzymes and bioactive peptides. It is the most natural foods beneficial to humans in all stages of life [2]. Dried powder cow's milk is one of the most widely used for alternative feeding in the world, accounting for the largest share of global production. It accounted for 83% of global milk production in 2010 [3]. Infant Formulas are used as an alternative to mother's milk (breast milk) to equip infants with nutritional needs either as а supplement Mother's to milk or individually when breastfeeding is not possible [4]. More than 80% of American children feed on infants formulas for some time during the first 12 months of their lives [5].

Most of infant formulas types are mainly made of cow milk with some modifications to substitute breast milk by adding protein in the form of basic amino acids and adjust fat and add unsaturated fatty acids, vegetable oils and some sugars are supported by some vitamins, minerals and nucleotides[6]. Infant formula must contain appropriate amounts of water, carbohydrates and protein, fats, vitamins, and minerals and regulate their composition with high rudder. Each plant must follow the guidelines established by the government authorities. For example, all major ingredients added to milk mixtures (protein, fat, carbohydrates) have minimum and maximum values to be active and these ingredients must have a safe use date [7]. Support milk mixtures only children's health and does not cause them unusual food or medical problems. Therefore, the regulation and control of the manufacturing process with high degree of

accuracy to meet national and international quality standards [8, 9].

Since Iraq depends on imports to meet its need of milk and mixtures of various companies, local markets have become replete with various kinds of infant formulas from many different origins of the world. This makes proper selection of the right formula is difficult. The careful selection of any infant formula must depend on the formula's components compared to other available formulas. The lack of such information is a pressing gap that need to be fulfilled. Thus the aim of this study was to identify the nutritional properties of some infant formulas available in local markets and whether or not they meet our young infants and children needs in terms of nutrients and growth requirements.

MATERIALS AND METHODS

Sampling: A comprehensive survey of the types of infant formula available in pharmacies and local markets has been conducted in the cities of Kufa and Najaf, three models of some infant formula were selected (Nactalia1, Dielac1 and Evolac1) for the First age of 6 months of birth and anther three formulas (Celia2, Dielac2 and Sunnybaby2) for the second stage infant age of 6 - 12 month.

EXPERIMENT ANIMALS

The experiment was conducted under laboratory conditions at the Animal House belongs to the College of Veterinary Medicine/ University of Kufa useing 42 white male rats (Albino) obtained from College of Veterinary Medicine.

The initial weight of rats were 50 ± 5 g and they were placed under controlled conditions in terms of ventilation and temperatures that ranged from 25 C° to 27 C° and lighting period of 12 hours per day. The animals were maintained in plastic cages, the cage floor was daily cleaned, sterilized and brushed with sawdust.

The animals were left for three days before starting the experiment for adapting to the new environment. The rats were randomly distributed to 7 groups with 6 rats per group, each rat was marked with different stain color for recognizing within the same group. After 3 days of adaptation, rats were treated daily for 28 days and treatments were as follows:

- C control group : The rats were fed on standard provender (Casein 20, Sucrose10, Cellulose5, Corn starch 53, Choline0.2, Mineral mixture3.5 and Vitamin mixture1, Corn oil 7)g/100g and filtered water [10].
- 2- **T1** Treatment group: The rats were fed on Nactalia1 for 4 weeks.
- 3- **T2** Treatment group: The rats were fed on Deilac1 for 4 weeks.
- 4- **T3** Treatment group: The rats were fed on Evolac1 for 4 weeks.
- 5- **T4** Treatment group: The rats were fed on Celia 2 for 4 weeks.
- 6- **T5** Treatment group: The rats were fed on Deilac2 for 4 weeks.
- 7- **T6** Treatment group: The rats were fed on Sunny baby2 for 4 weeks.

Monitoring, measuring and recording

The daily biotic activities of the experimental animals were monitored for the purpose of registering any notes and abnormal situations in the animals' movement or behavior. Animals have also been observed in case of change of eyes color, tail and hair of animals as well as recording if any case of diarrhea or any other unusual health issues.

Each animal weight was measured using a one-arm scale twice a week for further increases or decreases in body weight and the following indicators were calculated by the method used by previous study [11]

- 1. The amount of food consumed throughout the experiment period = total amount of food- remaining amount of food
- 2. Weight gain = final body weight initial body weight
- 3. Food efficiency rate = weight gain /amount of consumed food

The relative weight of the liver and kidneys:

At the end of the experiment rats feed was cut off for about 12 hours (overnight fasting) after that rats were anesthetized by muscle injection using my ketamine and Xylazine [12], Then the front and rear limbs were firmly fixed by pins and dissected posteriorly using a sharp pair of scissors and the liver and kidneys were cut off and washed in physiological solution of sodium chloride 0.9%, dried up between two towel papers with hand pressing, and the relative weight of liver and kidneys was calculated according to[13] using the equation where the relative weight of the liver = weight of the liver/weight of body *100 and the relative weight of the kidneys= weight of the kidneys/ weight of body *100.

THE STATISTICAL ANALYSIS

The experimental data were subjected to Analysis of Variance (ANOVA) using GenStat package 2009, (12th edition) version 12.1.0.3278 [14] Statistical analysis carried out for was the experiment, Completely which was Randomized Design (CRD). Means were compared for differences and Duncan's Multiple Range test was used whenever appropriate at 95% of confidence ($P \leq$ 0.05) [15].

RESULTS AND DISCUSSION

Effect of treatment with milk mixtures on weights of experimental animals:

Results of the statistical analysis shown in table (1) that there was no differences among the experimental animals in terms of initial average weight ($P \le 0.05$), while there was significant difference among averages of final weight. The lowest final weight in treatment T3 which was 118.7g compared with the control treatment C that of 151.9 g. However, T3 did not differ from the first ages formulas (T2, T1) with final average weight were 137.3 and 129.2g, respectively.

Whereas the T5 treatment had the highest body weight average at the end of experiment which was 169.8g with significant differences ($P \le 0.05$) from the first stage formulas, but not with the control (151.9 g), T4 (153.7 g) and T6 (148.5 g). All tested formulas including the control caused increase in the body weight. This mostly is due to their rich contents essential nutrients. vitamins of and minerals especially proteins that contribute in the construction of cells and body organs [16].

Weight gain after 28 days as in table (1) showed significant differences where the average of body weight increased ($P \le 0.05$), the highest increase in weight gain was in T₅ (118.4 g), while the lowest increase in weight gain was in T₃ (72.8 g). In the first stage formulas, T₂ had the highest increase of weight gain with 86.3 g,

which did not differed from the rest of the group members but at the same time weight gain of this group was less than that of the control (100.7 g). The total increase of weight gain in the second stage formulas also did not differed ($P \le 0.05$) from that yielded from the control. Rise in the overweight rates of rat groups by second-stage milk mixtures was not of a moral value compared to control treatment C at the same level of probability.

The variation in the rates of weight gained among treatments may be due to the variation in protein ratios that differed as infant formula differs. Dielac2 milk has higher protein content ratio and this resulted in higher rate of weight gain, and this is agreeing with Al-Jabari [17] study in which the weight gain was attributed to the type of infant formulas and their protein content.

Measured qualities Groups	The primary weight (g)	The weight after 28 days(g)	Weight Gain after 28 days (g)	daily increase in body weight(g/day)
C standard provender +filtered water	51.3 a	151.9 bc	100.7 bc	3.597 bc
T_1	54.6	129.2	74.6	2.664
Nactalia I	a	ab	a	a
T_2	51.0	137.3	86.3	3.081
Dielac 1	а	ab	ab	ab
Τ ₃	45.9	118.7	72.8	2.599
Evolac 1	а	а	а	а
T_4	48.4	153.7	105.3	3.762
Celia 2	а	bc	bc	bc
T_5	51.4	169.8	118.4	4.229
Dielac 2	а	С	С	С
T ₆	47.5	148.5	101.0	3.607
Sunny baby 2	а	bc	bc	bc

 Table 1. Primary (initial) and final weight and average of daily increase in body weight gained in 28 days experiment.

-data are average of 6 replications (6 rats per group)

-Means followed by different letter within column indicate significant difference ($P \le 0.05$)

Table (1) also showed differences among treatments in daily increase of body weight. T₅ was 4.229 g/day and significantly, ($P \le 0.05$) differed from all

first stage formulas $(T_1, T_2 \text{ and } T_3)$ which were 2.664, 3.081 and 2.599 g/day, respectively. T5, however, did not differed from all other treatments in the daily weight gain.

Weekly increase gained in weight of male rats

Figure (1) illustrates weekly increase gained in body weight of male rats during 4 weeks of treatments. It can be observed that weight gain during the first week in all treatments was dropped down with values less than that gained from the control. This may be due to the sudden shift in food supplement where the feed was changed from standard provender to infant milk formulas in the liquid form. The liquid decreasing the growth and body weight

decreasing the growth and body weight gain of all treatments except the control rats which were fed of the standard provender.

In the second week, the growth and weight gain for all treatments was enhanced. T_2 had highest weight gain (28.13g/week) among the first stage treatments. While, T5 had the highest body weight gain (35.6 g/week) compared to the second stage treatments. The body weight gain was continuous throughout the third week. T2 again had the highest weight gain (28.54 g/week) among the first stage formulas but was less than the weight gain of the control (33.44 g/week). However, two of the second stage formulas T5 (36.85 g/week) and T6 (35.59 g/week) resulted in higher, but not significant, weight gain compared to the control.



Means followed by different letter within column indicate significant difference ($P \le 0.05$)

Figure 1. Effect of treatment of infant formula on the weekly increase in the weight of male rats.

During fourth week, clearly decrease in weight gain for treatments C, T_2 , T_5 and T_6 which were 22.19, 15.27, 30.1 and 31.67 g/week, respectively, with a slight weight gain decrease in treatment T_3 that of 21.01g/week compared to values of the previous week. While continues weight gain was observed in treatments T4 (35.75 g/week) and T1 (28.49 g/week) compared with the previous week.

This decrease in weight gain may be related to damage of kidney tissue due to high protein ratio in second-stage milk formulas especially in Dielac2 and the control treatment ©. Al-Salihi [18] proposed that there was bleeding and congestion and kidney inflation caused by high protein in dietary supplements used to treat white rabbit males. It also reported by Fleddermann et. al., [19] that using less protein in infant milk formulas and fortified with high nutritional protein (a-Lactalbumin) will significantly reduce the overall functional load on infant's kidney.

Food intake, food efficiency rate and relative weight of internal organs

Table (2) illustrated the amount of food intake calculated on the basis of dry weight and food efficiency rate and relative weight of internal organs(liver and kidney) for rats treated by the infant formulas under this study.

There was significant differences in the amount of food intake at a level of probability ($P \le 0.05$). Rats group T₁, T₂ have highest values in term of amount of food intake, which reached 341.5 g and 337.5 g, respectively, and they were significantly different from the rest of all other treatments except the control in which the amount of food intake was 384.4g. T_3 , on the other hand, had significantly the lowest amount of food intake (251.00 g). This may be related to continuous cases of diarrhea in treated animals with this type of infant formula. resulting in anorexia or due to excessive fortification with vitamins and minerals. Al-Talib and Manki [6] stated that most infant formulas are supported by high proportions of vitamins and mineral more than recommended daily intake, especially vitamin A which causes a decrease in rat appetite and consequent weight loss [20].

			lieneg und i en	
liver and kidneys				
Measured qualities	amount of	food	relative	relative
Groups	food intake	efficiency	weight of	weight of
	(g)	rate (g)	liver %	kidney%
С	384.4	0.26	4.768	1.086
standard provender +filtered water	e	ab	а	b
T ₁	341.5	0.21	4.101	0.878
Nactalia 1	d	а	а	а
Τ2	337.5	0.26	4.769	0.898
Dielac 1	d	ab	а	а
Т3	251.0	0.29	4.839	0.849
Evolac 1	а	b	а	а
T4	277.2	0.34	4.884	0.924
Celia 2	b	с	а	ab
Т5	320.5	0.37	4.605	0.964
Dielac 2	С	с	а	ab
T6	276.2	0.36	4.519	0.955
Sunny baby 2	b	с	а	Ab

Table 2. The amount of Food intake and the rate of food efficiency and relative weight of

-data are average of 6 replications (6 rats per group)

-Means followed by different letter within column indicate significant difference ($P \le 0.05$)

Dewey et. al., [21] noted that having high amounts of iron by children between the ages of 4 and 9 months who have sufficient iron intake can lead to negative impact on the incidence of diarrhea. In the same way another studies [22, 23] referred to that intake of supported infant formula with high amounts of iron by infants with sufficient quantities may adversely affect growth, development and perception.

From table (2) there is significant difference in food efficiency rate among different treatments ($p \le 0.05$). Highest food efficiency rate were in treatments T_4 (0.34 g), T_5 (0.37 g) and T_6 (0.36 g). This may be due to the high protein content relative to the amount of energy provided by infant formulas Celia2, Dielac2 and Sunny baby2. While treatment T_1 was (0.21 g) the lowest in food efficiency rate with the lowest protein per 100 kilos of energy. Food efficiency rate in treatments T2 and T_3 were (0.26 and 0.29 g) respectively, and they did not differ from the control (0.26 g) food efficiency rate.

Also table (2) illustrated absence of significant differences in relative weight of rats' liver treated with the studied infant formulas ($p \le 0.05$). While observing the decrease of relative weight of rats' kidneys in treatments T_1 , T_2 and T_3 (first stage formulas) resulting in 0.878, 0.898 and 0.894 %, respectively, compared to that from the control treatment (1.086%).

The relative weight of rats' kidneys was much higher in treatment T4 (0.924%), T₅ (0.964%) and T₆ (0.955%)(Second stage formulas), but with no significant difference from the control treatment. The height of the relative weights of the kidneys related to high protein ratio in standard provender used in the control treatment and also in second stage formulas. This caused inflation, congestion and bleeding in the kidney tissue due to unwanted byproducts of the protein metabolism. And this agrees with Al-Jabari [10] explaining that the infant milk formulas containing higher protein ratio than recommended rate by the World Health Organization, Dielac (e.g.) milk group caused damage to the overall kidneys tissue, causing varying degrees of congestion and bleeding in the renal tissue, while the appropriate protein content (in Nactalia) did not cause any damage to the kidney tissue where the kidney tissue had normal appearance during the microscopic examination.

REFERENCES:

- Saxelin, M., Korpela, R., & Mäyrä-Mäkinen, A. 2003. Introduction: classifying functional dairy products. In *Functional Dairy Products, Volume* 1 (pp. 1-16).
- Kajal, M. F. I., Wadud, A., Islam, M. N., & Sarma, P. K. 2012. Evaluation of some chemical parameters of powder milk available in Mymensingh town. *Journal of the Bangladesh Agricultural University*, 10(1), 95-100.
- 3. FAOSTAT. 2012. FAO statistical database. Available at: <u>http://faostat.fao.org/</u>. Accessed 12 September 2012.
- 4. Harfouch, M., Mansour, A., & Mohammed, A. 2015. A study of the most important physical, chemical and microbial changes in infant milk during consumption and storage at room or refrigerator temperature. *Tishreen University Journal for Research and Scientific Studies -Biological Sciences Series*.73 (2), in Arabic: 73-87.
- Li, R., Darling, N., Maurice, E., Barker, L., & Grummer-Strawn, L. M. 2005. Breastfeeding rates in the United States by characteristics of the child, mother, or family: the 2002 National Immunization Survey. *Pediatrics*, 115(1), e31-e37.
- AL-Talib, Fatin F., & Manki, Nadia H. 2010. Study of Nutritional Adequacy Provided by some Imported Infant Milk Formula Available in the Local Market, J. of the College of Education for Women.21(3), in Arabic 730-741.
- 7. Koletzko, B., Baker, S., Cleghorn, G., Neto, U. F., Gopalan, S., Hernell, O., ... & Pzyrembel, H. 2005. Global standard for the composition of infant formula: recommendations of an **ESPGHAN** coordinated international expert group. Journal of pediatric gastroenterology and nutrition, 41(5), 584-599.
- World Health Organization. 2017. Safe Preparation, Storage and Handling of Powdered Infant Formula Guidelines. Available online:http: //www.who.int/foodsafety

/publication/micro /pit_quideline.pdf (accessed on13Feberuary 2017).

- 9. Lönnerdal, B. 2012. Preclinical assessment of infant formula. Annals of Nutrition and Metabolism, Vol.60, No.3, pp.196-199.
- 10. Al harbi, L. Bint Naif. 2008. Effect of Selected Natural and Synthetic Colors Used in Children's Candy on Liver and Kidney in Rats. Thesis, Dept. Food Scie & Nutr., Coll. of Food & Agric. Sciences. Univ. of King Saud, pp.107.
- 11. Miller, D. S. 1963. Procedure for Determination of NPU using Rats Body nitrogen Techniques, In Evaluation of Protein Quality, Nutrition Academy of National Research Sciences council Publication, Washington, D. C.
- 12. Al-Shaikh, Sharaf. A. H. 2018. Studying Some Biochemical and Nutritional Characteristics of Selenium-enriched Yoghurt. Thesis, Dept. Food Scie., Coll. of Agric., Univ. of Baghdad, in Arabic.pp.140.
- 13. A. O. A. C. (1990). Official methods of analysis of associtation of official analytical chemists. Arlington, west Virginia, USA.
- 14. GenStat. (2009). General Statistical GenSta22. Majumdar, I., Paul, P., Talib, V. H., & Guides V. 12.1, Copyright 2009, VSN International Ltd. UK.
- 15. Duncan, D. B. (1955). Multiple range and multiple F tests. Biometrics, 11(1), 1-42.
- 16. Al-Zuhairi, A. M. Th. 2000. Human Nutrition, Ministry of Higher Education & Scientific Research, Univ. of Mosul, Iraqi National Library & Archives, pp.607.
- 17. Al-Jabari Mariam K. Gh. 2016.Study of Physiological, Biochemical and Microbiological of Infant Formula Available in the Iraqi Local Markets.

Thesis, Dept. Food Scie., Coll. of Agric., Univ. of Tikrit.in Arabic, pp.118.

- 18. Al-Salihi, Ronza F. 2014. Study of Some Physiological and Histological Effects of a Number of Dietary Supplements On Male White Rabbits. Thesis, Dept. Food Scie., Coll. of Agric., Univ.of Tikrit.in Arabic.pp.112
- 19. Fleddermann, M., Demmelmair, H., Grote, V., Nikolic, T., Trisic, B., & Koletzko, B. 2014. Infant formula composition affects energetic efficiency for growth: the BeMIM study, a randomized controlled trial. Clinical nutrition, 33(4), 588-595.
- 20. Hadeel, M. Ahmad. 2012. Effect of Excessive Levels of Vitamin A on Bone Formation and some Biochemical Parameters in Young Rats. Rafidain journal of science, 23(2), 44-54.
- 21. Dewey, K. G., Domellof, M., Cohen, R. J., Rivera, L. L., Hernell, O., & Lonnerdal, B. Community and International 2002. Nutrition-Iron Supplementation Affects Growth and Morbidity of Breast-Fed Infants: Results of a Randomized Trial in Sweden and Honduras. Journal of Nutrition, 132(11), 3249-3255.
- Ranga, S. 2003. The effect of iron therapy on the growth of iron-replete and ironchildren. Journal of Tropical deplete Pediatrics, 49(2), 84-88.
- 23. Lind, T., Seswandhana, R., Persson, L. Å., & Lönnerdal, B. 2008. Iron supplementation of iron-replete Indonesian infants is associated with reduced weight-for-age. Acta Paediatrica, 97(6), 770-775.