Baghdad Science Journal

Volume 22 | Issue 5

Article 11

5-19-2025

Amplification of D18S51, D8S1179, and FES Loci of DNA Samples Isolated From Guitar String Stored at Room Temperature

Abdul Hadi Furqoni

Center for Biomedical Research, Research Organization for Health, National Research and Innovation Agency (BRIN), Cibinong Science Center, Cibinong - Bogor, West Java, Indonesia AND Faculty of Medicine, Universitas Islam Negeri Syarif HIdayatullah Jakarta, Banten, Indonesia, cocohadi01@gmail.com

Anindita Riesti Retno Arimurti

Diploma Medical Laboratory Technology, Faculty of Health Science, Universitas Muhammadiyah Surabaya, Surabaya, Indonesia, aninditariesti@um-surabaya.ac.id

Meri Meri Universitas Bakti Tunas Husada, Tasikmalaya, West Java, Indonesia, meri@universitas-bth.ac.id

Ni Luh Ayu Megasari

Postgraduate School, Universitas Airlangga, Surabaya, Indonesia AND Institute of Tropical Disease, Universitas Airlangga, Surabaya, Indonesia, ni.luh.ayu@pasca.unair.ac.id

Indah Nuraini Masjkur Institute of Tropical Disease, Universitas Airlangga, Surabaya, Indonesia, indahnuraini61@gmail.com

See next page for additional authors Follow this and additional works at: https://bsj.uobaghdad.edu.iq/home

How to Cite this Article

Furqoni, Abdul Hadi; Arimurti, Anindita Riesti Retno; Meri, Meri; Megasari, Ni Luh Ayu; Masjkur, Indah Nuraini; Huda, Qurrota A'yunil; Nuraini, Intan Sari; and Hermawati, Luluk (2025) "Amplification of D18S51, D8S1179, and FES Loci of DNA Samples Isolated From Guitar String Stored at Room Temperature," *Baghdad Science Journal*: Vol. 22: Iss. 5, Article 11. D0I: https://doi.org/10.21123/bsj.2024.10206

This Article is brought to you for free and open access by Baghdad Science Journal. It has been accepted for inclusion in Baghdad Science Journal by an authorized editor of Baghdad Science Journal.

Amplification of D18S51, D8S1179, and FES Loci of DNA Samples Isolated From Guitar String Stored at Room Temperature

Authors

Abdul Hadi Furqoni, Anindita Riesti Retno Arimurti, Meri Meri, Ni Luh Ayu Megasari, Indah Nuraini Masjkur, Qurrota A'yunil Huda, Intan Sari Nuraini, and Luluk Hermawati

This article is available in Baghdad Science Journal: https://bsj.uobaghdad.edu.iq/home/vol22/iss5/11

Scan the QR to view the full-text article on the journal website



RESEARCH ARTICLE

Amplification of D18S51, D8S1179, and FES Loci of DNA Samples Isolated from Guitar String Stored at Room Temperature

Abdul Hadi Furqoni[®] ^{1,2}, Anindita Riesti Retno Arimurti[®] ³, Meri Meri[®] ⁴, Ni Luh Ayu Megasari[®] ^{5,6,*}, Indah Nuraini Masjkur[®] ⁶, Qurrota A'yunil Huda[®] ⁶, Intan Sari Nuraini[®] ⁷, Luluk Hermawati[®] ²

¹ Center for Biomedical Research, Research Organization for Health, National Research and Innovation Agency (BRIN), Cibinong Science Center, Cibinong - Bogor, West Java, Indonesia

- ³ Diploma Medical Laboratory Technology, Faculty of Health Science, Universitas Muhammadiyah Surabaya, Surabaya, Indonesia
- ⁴ Universitas Bakti Tunas Husada, Tasikmalaya, West Java, Indonesia
- ⁵ Postgraduate School, Universitas Airlangga, Surabaya, Indonesia
- ⁶ Institute of Tropical Disease, Universitas Airlangga, Surabaya, Indonesia
- ⁷ Student of Forensic Science Master Program, Postgraduate School, Universitas Airlangga, Surabaya, Indonesia

ABSTRACT

The act of criminality is an unlawful behavior that constitutes a criminal offense. Criminals use various ways to hide evidence at the crime scene. Forensic experts often find DNA on items found at the crime scene, one of the types of evidence is a guitar. Guitar is one of the musical instruments often played by humans that can be used as a comparison. Methods: This study used 18 guitar string samples that had been used for five minutes and incubated at room temperature. The 18 samples were then divided into three groups, in which each group consisted of six samples and incubated for one, five, and 10 days. DNA identification was then carried out using a UV spectrometer for DNA quantification and the DNAzol method for DNA extraction. Results: The mean result of DNA quantification on day 1 was $152.57 \pm 48.02 \text{ ng/}\mu\text{L}$, day 5 was $138.66 \pm 47.66 \text{ ng/}\mu\text{L}$, and day 10 was $87.09 \pm 9.07 \text{ ng/}\mu\text{L}$. Polymerase chain reaction was carried out using three STR primers loci, namely D18S51 (290–366 bp), D8S1179 (203–224 bp), and FES (222–250 bp), followed by visualization using the silver nitrate method. The final results showed that all samples could be amplified using the D18S51, D8S1179, and FES STR loci.

Keywords: D18S51, D8S1179, FES, STR Loci, Guitar string, Crime

Introduction

Forensic identification is a step that is carried out to assist investigators in identifying individuals who are victims in criminal cases that have caused death, because this is an attempt to respect human rights. Therefore, it can be interpreted that the identification process is a very crucial stage that must be carried out before proceeding to the next stage, namely the examination of the human remains (autopsy). $^{\rm 1}$

Personal identification is very essential in the investigation of both criminal and civil cases because any mistake can be fatal in the judicial process. Identification can be done in three ways: visual (relatives or acquaintances seeing the human remains); in detailed (e.g., ante-mortem data that matched with the

Received 14 November 2023; revised 7 July 2024; accepted 9 July 2024. Available online 19 May 2025

* Corresponding author.

E-mail addresses: cocohadi01@gmail.com (A. H. Furqoni), aninditariesti@um-surabaya.ac.id (A. R. R. Arimurti), meri@universitas-bth.ac.id (M. Meri), ni.luh.ayu@pasca.unair.ac.id (N. L. A. Megasari), indahnuraini61@gmail.com (I. N. Masjkur), qurrotaayun.elhuda@gmail.com (Q. A'yunil Huda), isari8098@gmail.com (I. S. Nuraini), lulukhermawati.bio@gmail.com (L. Hermawati).

https://doi.org/10.21123/bsj.2024.10206

² Faculty of Medicine, Universitas Islam Negeri Syarif HIdayatullah Jakarta, Banten, Indonesia

^{2411-7986/© 2025} The Author(s). Published by College of Science for Women, University of Baghdad. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

information that was collected during the autopsy and other situational information); and scientific or objective (e.g., the examination of teeth, fingerprints, or deoxyribonucleic acid (DNA)).²

DNA is the smallest genetic structure present in all forms of life, from microorganisms to higher-level organisms such as humans, animals and plants. Each type of tissue has a different level of DNA concentration, which depends on the structure and composition of its cells. Usually, tissues that have many nucleus cells and few connective tissues tend to have high DNA concentration. Therefore the selection of DNA samples for forensic analysis is very diverse.³

The method used in forensic DNA analysis is to examine the region of DNA repetition that have a base sequence of less than 1 kb. These regions are known as microsatellites or also called Short Tandem Repeats (STR). Short Tandem Repeats are microsatellite sequences that have repetition units between 2 and 7 base pairs in length. STR analysis is a molecular biology technique commonly used to compare the number of allele repetition at a particular locus in DNA between two or more samples. To determine the STR length, the Polymerase Chain Reaction (PCR) method is used in which is based on the length of the PCR product. Short Tandem Repeats, as the most commonly used genetic marker, is distributed widely in the human genome, and has been widely used in DNA profiling, especially in terms of individual identification and paternity testing for decades.⁴

DNA identification is an attempt to compare the DNA profile of evidence with a comparison sample, with the aim of determining whether or not the DNA profile of the evidence matches with the comparison sample. The DNA identification process relies on specific inheritance patterns found in nuclear DNA and mitochondrial.⁵

In DNA identification involving somatic chromosomes, the commonly used comparators are the parents (father and mother). While in DNA identification involving the Y chromosome, the comparator is usually the biological father, paternal grandfather, or paternal brother and so forth. In the examination of mitochondrial DNA, the comparator used is a family member in one lineage derived from the mother.⁶

The most commonly used DNA profiling technique is Short Tandem Repeat (STR) because of its convenience, requiring only a minimum of 13 loci as reference. The FBI recommends the following thirteen loci, mainly known as Combined DNA Index System (CODIS) Core STR loci, as references: TH01, TPOX, CSF1P0, vWA, FGA, D3S1358, D5S818, D7S820, D13S317, D16S539, D8S1179, D18S51, D21S11, and added with amelogenin markers that are useful for analyzing gender.⁷

Touch DNA is defined as the epithelial cells left on an object held or used by the suspect.^{8–11} Touch DNA can be used for profiling analysis, and can be obtained from biological samples. Several studies have demonstrated successful DNA profiling using blood, buccal swab, hair, semen, saliva, teeth and bones samples.¹¹⁻¹³ In addition to biological samples, DNA can also be isolated from items found at the crime scene, such as clothes and gloves,¹⁴ masks,¹⁵ and rings.⁸ Other common items, such as household objects might also provide samples for DNA profiling.¹⁶ Musical instruments are among popularly bought and owned household items. Aside of violin and piano, guitar was reported as one of the most purchased musical instruments.¹⁷ However, no previous study describing profiling analysis using DNA isolated from musical instrument, including guitar. As an instrument commonly owned and played, valuable touch DNA sample might be obtained from guitar.

Indonesian Supreme Court has previously reported guitar as one of the crime pieces of evidence.¹⁸ However, examination of DNA swabs derived from guitar (guitar strings) using the STR method has not been carried out in Indonesia. This study aims to analyze DNA derived from guitar string swabs using the STR method as treated with different lengths of storage time at room temperature. The loci used in the analysis of this study include loci D18S51 D8S1179, FES. The three loci were used because they have high discrimination in the Indonesian population.¹⁹

Materials and methods

This research has passed the ethical test with number 337/HRECC.FODM/VII/2020. Samples were taken from guitar strings played by three volunteers for five minutes. The total number of samples were 18 guitar strings, which were divided into three groups consisted of six strings each. Group I consisted of the first and second strings, group II of the third and fourth strings, and group III of the fifth and sixth string (Fig. 1). Groups I to III were then left at room temperature for one, five, and 10 days, respectively. Room temperature was maintained around 20°C.²⁰ using air conditioner, due to higher and fluctuating ambient temperature in Indonesia. Wet swabs samples were then collected from each strings, followed by DNA extraction using the DNAzol method.²¹ The extracted DNA was quantified using a UV-visible spectrophotometer Evolution One (Thermo Fisher Scientific, United States of America).²²

After DNA quantification, two samples from each group with the highest and lowest concentration and purity were chosen for DNA amplification using



Fig. 1. Illustration of the guitar strings position.

polymerase chain reaction (PCR). Higher DNA yields and purity generally more suitable for PCR analysis, while low DNA yields and purity might cause amplification failure. Therefore, we selected samples with high and low DNA concentration and purity to determine whether PCR amplification could be performed in different sample condition.²³ The PCR reactions were carried out using GoTaq®Green Master Mix (Promega, United States of America) in T100TM Thermal Cycler (Bio-Rad Laboratories, United States of America). Primers used in the PCR reactions were from Promega (United States of America), and described in Table 1. PCR products were visualized using a 2% acrylamide gel run on 100V for 45 minutes, and then colored with silver staining.²⁴

Results and discussion

Our results showed that samples collected from group I (stored for one day) has the highest average DNA concentration and purity, and group III (stored for 10 days) has the lowest average DNA concentration and purity (Tables 2 and 3).

In the visualization process using an acrylamide gel, seen in Fig. 2, it is evident that all selected samples from all groups have successfully shown positive amplification of locus D18S51, despite having different DNA concentration and purity. Likewise, in Figs. 3 and 4, using the D8S1179 and FES loci, all selected samples have also successfully amplified.

This study showed variations in the average DNA concentration and purity of samples collected from guitar string stored at room temperature. The concentration and purity of DNA are affected by several factors that contribute to DNA degradation. One of these factors is endogenous factors, which originate within the cell itself. The process of cell damage begins with autolysis and then undergoes decay followed by aerobic decomposition. The other is exogenous factors, which include environmental influences such as temperature and humidity.^{26–28} Abdel Hady et al.²⁹ reported that samples stored at minus 20°C have higher levels and purity compared to samples stored at 4°C. Other studies have also shown that there are many factors that can affect DNA degradation, such as chromatin structure, transcriptional activity and cellular location. Therefore, it is recommended to perform DNA analysis not only by examining nuclear DNA but also by identifying using mitochondrial DNA.³⁰

Methods used for DNA extraction can also affect the quality of DNA obtained. The DNAZol methods is suitable for the isolation wide range of DNA molecules,

Table ⁻	. PCR	primers.	25
--------------------	-------	----------	----

Primer	Sequence	Amplicon length (base pairs (bp))
D18S51 loci	Forward Primer: 5'-AGCTTATGCAAGGACTGGAC-3'	290-366
	Reverse Primer: 5'-TCTTCCAGAAAGCAGGGTGT-3'	
D8S1179 loci	Forward Primer: 5'-TCTGGGGCAGTAGGGAACAG-3'	203-224
]	Reverse Primer: 5'-GGGGGCTGACTAGCAAGAGA-3'	
FES loci	Forward Primer: 5'-T7-GCTTGTTAATTCATGTAGGGAAGGC-3' Reverse Primer: 5'-GTAGTCCCAGCTACTTGGCTACTC-3'	222–250

Table 2. DNA concentration of samples from each gro	oup.
---	------

Treatment Group	Average DNA Concentration (ng/ μ l) \pm SD
Group I	$152.57 \pm 48,02$
Group II	$138.66 \pm 47,66$
Group III	$87.09 \pm 9,07$

Treatment Group	Average DNA Purity (ng/ μ l) \pm SD
Group I	1.58 ± 0.11
Group II	1.42 ± 0.12
Group III	1.21 ± 0.08



Fig. 2. Visualization results of guitar string DNA using locus D18S51 (290–336 bp).

including genomic DNA and DNA fragments down to 0.1 kb in length which are mainly targeted in DNA profiling. Isolated DNA is ready for molecular biology applications, without needing additional purification. Samples stored in DNAzol also shown stable DNA concentration and purity, despite being stored up to six months in room temperature.³¹

The degradation of DNA due to long storage periods can be caused by a number of factors, Hydrolysis is one of the factors of DNA degradation where hydrolytic reactions occur in the glycosidic sugar bonds of DNA which can lead to depurination, which is the loss of one nitrogen base from the DNA sequence over time.³² Environmental influences also affect the process of DNA degradation where environmental factors such as temperature, humidity, and pH can affect DNA stability. For example, high temperatures or acidic or alkaline environmental conditions can accelerate DNA degradation. Exposure to ultraviolet (UV) light can damage the structure of DNA and cause cross-linking between adjacent DNA molecules. Oxidation also converts DNA bases, especially pyrimidines, into different forms such as hydantoin, which



Fig. 3. Visualization results of guitar string DNA using locus D8S1179 (203–224 bp).



Fig. 4. Visualization results of guitar string DNA using locus FES (222–250 bp).

can disrupt DNA stability. All these factors can contribute to DNA degradation over time, and such damage can hinder DNA analysis such as PCR and STR analysis. Therefore, it is necessary to consider these factors in the storage and analysis of DNA samples.³³

Locus D18S51 is one of the loci in the chromosome 18. The locus has a specific DNA sequence which is useful in forensic DNA analysis, and often used for STR analysis to produce an individual DNA profile. This profile is useful for uniquely identifying individuals, as the length and sequence of DNA at locus D18S51 can vary between individuals, thus being used to distinguish one individual from another.³⁴

Locus D8S1179 is a specific locus on the human chromosome 8 used in forensic DNA analysis. In the context of forensic analysis, locus D8S1179 is one of the STR loci that is often tested to generate a DNA profile of an individual. STRs are sections of DNA where short sequences of multiple base pairs repeat in a sequence, and these lengths and sequences can vary between individuals. Therefore, locus D8S1179 is used to distinguish individuals in forensic identification by examining variations in the length and sequence of STRs at this locus.³⁵ The FES locus is a specific location on the genome that refers to a human gene known as "FES" or "Feline Sarcoma Oncogene". The locus is located in the chromosome 15. This gene is involved in the regulation of cell growth and development, and has been linked to various aspects of biology such as cancer and cell proliferation. In addition, certain loci are also used in forensic DNA analysis. In that context, the FES locus is one of the loci or sites tested to generate individual DNA profiles for forensic analysis.³⁶

These three loci have a high level of discrimination in personal identification using DNA. According to the study conducted by Arimurti et al.³⁷ showed that locus D18S51 and locus D8S1179 can be used for DNA identification of earphone swabs by STR method. According to the study conducted by Yudianto et al.³⁸ mentioned that the locus FES can be used to identify allele sharing patterns between siblings in the Madura tribe that are inherited by their parents.

Conclusion

From the results and discussion of the study above, it can be concluded that storage of guitar string samples at room temperature for 1, 5 and 10 days causes DNA degradation in terms of average levels and DNA purity results. The DNA isolated from guitar strings can be amplified using D18S51, D8S1179, and FES loci as detection targets; therefore, may be beneficial for forensic identification.

Acknowledgment

This research had been supported by the Institute of Tropical Disease as a Center of Excellence of the Indonesian Ministry of Education, Culture, Research, and Technology.

Authors' declaration

- Conflicts of Interest: None.
- We hereby confirm that all the Figures and Tables in the manuscript are ours. Furthermore, any Figures and images, that are not ours, have been

included with the necessary permission for republication, which is attached to the manuscript.

- Authors sign on ethical consideration's approval.
- No animal studies are present in the manuscript.
- Ethical Clearance: The project was approved by the local ethical committee at Institute of Tropical Disease, Universitas Airlangga, Surabaya, Indonesia.

Authors' contribution statement

A.H.F and N.L.A.M contributed to the conception of study. A.R.R.A, I.N.M., Q.A.H., I.S.N., A.P., and R.F. performed sample collection and experiment. A.H.F. and A.R.R.A. analyzed the data. A.H.F. and N.L.A.M. wrote the manuscript. The final version of the manuscript has been reviewed and approved by all authors.

References

- Daga SS, Ahmed U, Kumawat RK. Forensic tools and techniques of absolute human identification: Physical and molecular approach. J Forensic Sci Crim Investig. 2021;15(3):1–6.
- Turingan RS, Brown J, Kaplun L, *et al.* Identification of human remains using rapid DNA analysis. Int J Legal Med. 2020;134(3):863–872. https://doi.org/10.1007/ s00414-019-02186-y.
- Romeika JM. Recent advances in forensic DNA analysis. J Forensic Res. 2013;s12(01). https://doi.org/10.4172/2157-7145.S12-001.
- Mahat N, Alwi A, Salleh F, Ishar S, Kamaluddin M, Rashid M. Applications of X-chromosome short tandem repeats for human identification: A review. J Trop Life Sci. 2023;13(1):193– 218. https://doi.org/10.11594/jtls.13.01.19.
- Hernández C. Mitochondrial DNA in human diversity and health: From the golden age to the Omics era. Genes (Basel). 2023;14(8):1534. https://doi.org/10.3390/genes14081534.
- Poma A, Cesare P, Bonfigli A, Volpe AR, Colafarina S, Vecchiotti G, *et al.* A qPCR-duplex assay for sex determination in ancient DNA. PLoS One. 2022;17(6):e0269913. https://doi. org/10.1371/journal.pone.0269913.
- Yudianto A, Nuraini MI, Furqoni AH, Nzilibili SMM, Harjanto P. The use of touch dna analysis in forensic identification focusing on STR CODIS LOCI THO1, CSF1PO and TPOX. Indian J Forensic Med Toxicol. 2020 Jul 6;12(Suppl 1):8716. https://doi.org/10.37506/ijfmt.v14i3.10667.
- Furqoni AH, Megasari NLA, Yudianto A, Azizah F, Maulidiyanti ETS, Fahmi NF, *et al.* Amplification of vWA, FGA, and TH01 loci of DNA samples isolated from ring stored at room temperature. Malays J Med Health Sci. 2023;19(5):97–101.
- 9. Gosch A, Courts C. On DNA transfer: The lack and difficulty of systematic research and how to do it better. Forensic Sci Int Genet. 2019;40:24–36. https://doi.org/10.1016/j.fsigen. 2019.01.012.
- Comte J, Baechler S, Gervaix J, Lock E, Milon MP, Delemont O, *et al.* Touch DNA collection – Performance of four different swabs. Forensic Sci Int Genet. 2019;43:102113. https://doi. org/10.1016/j.fsigen.2019.06.014.

- 11. Li Y, Xie M, Wu J. DNA profiling in peripheral blood, buccal swabs, hair follicles and semen from a patient following allogeneic hematopoietic stem cells transplantation. Biomed Reports; 2014;2(6):804–808. doi:10.3892/br.2014.332.
- Puri P, Dubey M, Kumar N. DNA Profiling in Forensic Odontology. 1st ed. (Dash H, Shrivastava P, Lorente J, eds.). Springer Singapore; 2022. https://doi.org/10.1007/978-981-15-9364-2_63-1.
- Le TN, Muratovic D, Handt O, Henry J, Linacre A. DNA profiling from human bone cells in the absence of decalcification and DNA extraction. J Forensic Sci. 2022;67(4):1690–1696. doi:10.1111/1556-4029.15033.
- Udogadi NS, Abdullahi MK, Bukola AT, Imose OP, Esewi AD. Forensic DNA profiling: Autosomal short tandem repeat as a prominent marker in crime investigation. Malaysian J Med Sci. 2020;27(4):22–35. https://doi.org/10.21315/mjms2020. 27.4.3.
- Dash HR, Arora M, Khatoon S. Evaluation of face masks as a valuable forensic DNA evidence in the post-COVID era. Int J Legal Med. 2023;137(2):311–317. https://doi.org/10.1007/ s00414-023-02945-y.
- Tozzo P, Mazzobel E, Marcante B, Delicati A, Caenazzo L. Touch DNA sampling methods: efficacy evaluation and systematic review. Int J Mol Sci. 2022;23(24):15541. https://doi. org/10.3390/ijms232415541.
- Tang L. Primary research on the network marketing of musical instrument galleries. In: Proceedings of the 2012 2nd International Conference on Computer and Information Applications (ICCIA 2012). Atlantis Press; 2012. https://doi.org/10.2991/ iccia.2012.206.
- Indonesian Supreme Court. PUTUSAN PN MATARAM 164/PID.B/2021/PN MTR. Published 2021.
- Furqoni AH, Palupi R, Yudianto A, Sosiawan A, Tirthaningsih NW, Fore F, *et al.* Analysis of sibling pair relationships of balineses indonesia, using 12 STR loci for human identification process. J Int Dent Med Res. 2021;14(2):855–859. ISSN 1309–100X.
- Silva JATD. Room temperature in scientific protocols and experiments should be defined: A reproducibility issue. Biotechniques. 2021;70(6):306–308. https://doi.org/10.2144/btn-2020-0131.
- Firmansyah NE, Thongseesuksai T, Boonmars T, Sungkar S, Laummaunwai P. Comparison of 3 DNA extraction methods for extracting DNA from an adult Culex quinquefasciatus (Diptera: Culicidae). J Insect Sci. 2023;23(5):15. https://doi. org/10.1093/jisesa/iead080.
- Minhas-Khan A, Ghafar-Zadeh M, Shaffaf T, Forouhi S, Scime A, Magierowski S, *et al.* UV-Vis spectrophotometric analysis of DNA retrieval for DNA storage applications. Actuators. 2021;10(10):246. https://doi.org/10.3390/act10100246.
- 23. García-Alegría AM, Anduro-Corona I, Pérez-Martínez CJ, Guadalupe Corella-Madueño MA, Rascón-Durán ML, Astiazaran-Garcia H. Quantification of DNA through the NanoDrop spectrophotometer: Methodological validation using standard reference material and sprague dawley rat and human DNA. Int J Anal Chem. 2020;1–9. https://doi.org/10.1155/2020/8896738.
- Jiang Y, Zheng L, Lin L, Lin S, Xu K, Deng S, *et al.* Modification in silver staining procedure for enhanced protein staining. Teekaraman Y, ed. Biomed Res Int. 2022;1–9. https://doi.org/ 10.1155/2022/6243971.

- Hanssen EN, Lyle R, Egeland T, Gill P. Degradation in forensic trace DNA samples explored by massively parallel sequencing. Forensic Sci Int Genet. 2017;27:160–166. https://doi.org/10. 1016/j.fsigen.2017.01.002.
- Bhoyar L, Mehar P, Chavali K. An overview of DNA degradation and its implications in forensic caseworks. Egypt J Forensic Sci. 2024;14(1):15. https://doi.org/10.1186/s41935-024-00389-y.
- Naef T, Besnard A, Lehnen L, Petit EJ, van Schaik J, Puechmaille SJ. How to quantify factors degrading DNA in the environment and predict degradation for effective sampling design. Environ DNA. 2023;5(3):403–416. https://doi.org/10.1002/edn3.414.
- Zhu G, Guo YA, Ho D, Poon P, Poh ZW, Wong PM, et al. Tissue-specific cell-free DNA degradation quantifies circulating tumor DNA burden. Nat Commun. 2021;12(1):2229. https: //doi.org/10.1038/s41467-021-22463-y.
- 29. Abdel Hady RH, Thabet HZ, Ebrahem NE, Yassa HA. Thermal effects on DNA degradation in blood and seminal stains: Forensic view. Acad Forensic Pathol. 2021;11(1):7–23. https: //doi.org/10.1177/1925362121998547.
- Samehsalari S, Chandrasekar A. Muslims from South India. J Genet Resour. 2021; 7(2): 220–226. https://doi.org/10. 22080/jgr.2021.20887.1239
- Yudianto A, Erliyah T, Furqoni AH, Nuraini I, Huda QA. Effect of storage time on DNA content and purity in lip print. J Biosains Pascasarj. 2023;25(1):43–48. https://doi.org/10. 20473/jbp.v25i1.2023.43-48.
- Divya K, GK M, Umema A, SS D. Environmental factors affecting the concentration of DNA in blood and saliva stains: A review. J Forensic Sci Res. 2024;8(1):009–015. https://doi. org/10.29328/journal.jfsr.1001057.
- 33. Khoury M El. The effects of heat and explosions on forensic DNA analyses by the effects of heat and explosions on forensic DNA analyses (PhD Dissertation). University of Leicester; 2020.
- 34. Aflanie I, Prastowo W, Panghiyangani R, Yudianto A, Koesbardiati T. Genetic variation analysis and kinship relationship between Dayak Ngaju tribe and Dayak Bukit tribe through examination of core DNA of Bukit CODIS STR Locus (combine DNA index system) 13 for the purpose of forensic identification. Indian J Forensic Med Toxicol; 2020;14(3):1094–1097. https://doi.org/10.37506/ijfmt.v14i3.10519.
- Manela C, Susanti R, Tjong DH, Yudianto A. Genetic analysis 21 short tandem repeats (STR) locus in minangkabau population, West Sumatera, Indonesia. African J Infect Dis. 2022;16(2):35–41. https://doi.org/10.21010/Ajid.v16i2.4.
- Wyner N, Barash M, McNevin D. Forensic autosomal short tandem repeats and their potential association with phenotype. Front Genet. 2020;11:884. https://doi.org/10.3389/ fgene.2020.00884.
- Arimurti ARR, Yudianto A, Astuti SP. Identifikasi DNA dari swab earphone dengan teknik STR (Short Tandem Repeat) untuk Kepentingan Forensik. J Biosains Pascasarj. 2015;17(1):26–32. https://doi.org/10.20473/jbp. v17i1.2015.26-32.
- Sosiawan A, Yudianto A, Furqoni AH, Nzilibili SMM, Nuraini I. Full-sibling allelic frequency and sharing among Madurese: STR technique by 12 locus and the sex-typing amelogenin gene. Egypt J Forensic Sci. 2019;9(1):38. https://doi.org/10. 1186/s41935-019-0143-5.

تضخيم مواقع D18S51و D8S1179 وFES لعينات الحمض النووي المعزولة من أوتار الجيتار المخزنة في درجة حرارة الغرفة

عبد الهادي فرقوني ^{1,2}، أنينديتا ريستي ريتنو أريمورتي ³، ميري ميري ⁴، ني لوه أيو ميجاساري ^{5,6}، إنداه نوريني مسجد كور⁶، قروتا عيون الهدي⁶، إنتان ساري نوريني⁷، لولوك هرماواتي ²

¹مركز البحوث الطبية الحيوية، منظمة أبحاث الصحة، الوكالة الوطنية للبحث والابتكار ، مركز سيبينونج للعلوم _بسيبينونج، بوجور ، جاوة الغربية، إندونيسيا.

> ²كلية الطب، جامعة شريف هداية الله الإسلامية الحكومية جاكرتا، بانتن، إندونيسيا. ³دبلوم في تكنولوجيا المختبرات الطبية، كلية العلوم الصحية، الجامعة المحمدية في سور ابايا، سور ابايا، إندونيسيا. ⁴جامعة باكتي توناس هوسادا، تاسيكمالايا، جاوة الغربية، إندونيسيا. ⁵مدرسة الدراسات العليا، جامعة إير لانجا، سور ابايا، إندونيسيا. 7 معهد أمراض المناطق المدارية، جامعة إير لانجا، سور ابايا، إندونيسيا.

> ⁷طالبة في برنامج الماجستير في علوم الطب الشرعي، كلية الدر اسات العليا، جامعة إير لانجا، سور ابايا، إندونيسيا.

الخلاصة

الفعل الأجرامي هو سلوك غير قانوني وقد يكون جريمة جنائية ، ويستخدم المجرمون طرقا مختلفة لأخفاء الأدلة في مسرح الجريمة, ومن هذه الجريمة لكن غالبا ما يجد خبراء الطب الشرعي اثار الحمض النووي على العناصر الموجودة في مسرح الجريمة, ومن هذه الأدلة هو الجيتار. والجيتار من الألات الموسيقية التي يعزف عليها البشروالتي يمكن استخدامها للمقارنة . استخدمت الطرق في هذه الدراسة 18 عينة من أوتار الجيتار التي تم استخدامها لمدة خمس دقائق وحضنت في درجة حرارة الغرفة .تم بعد في هذه الدراسة 18 عينة من أوتار الجيتار التي تم استخدامها لمدة خمس دقائق وحضنت في درجة حرارة الغرفة .تم بعد ذلك تقسيم العينات الـ 18 إلى ثلاث مجموعات، حيث تتكون كل مجموعة من ست عينات وتم تحضينها لمدة يوم وخمسة و 10 أيام. تم بعد ذلك تقسيم العينات الـ 18 إلى ثلاث مجموعات، حيث تتكون كل مجموعة من ست عينات وتم تحضينها لمدة يوم وخمسة و 10 أيام. تم بعد ذلك قياس الحمض النووي باستخدام مطياف الأسعة فوق البنفسجية للحمض النووي المستخلص بطريقة . DNAzol و 10 أيام. تم بعد ذلك قياس الحمض النووي باستخدام مطياف الأسعة فوق البنفسجية للحمض النووي المستخلص بطريقة كان را 20 أيام. تم بعد ذلك قياس الحمض النووي باستخدام مطياف الأسعة فوق البنفسجية للحمض النووي المستخلص بطريقة . DNAzol و 20 أيام. تم دلم النووي المستخلص النووي أي ما 20 أيام. تم بعد ذلك قياس الحمض النووي في بعد الحضن لليوم الأول (20.8 ± 25.5 نانو غرام/ميكروليتر) واليوم 20 أيام (20.5 ± 20.5) و 20 أيام ما 20 أيام ميكروليتر) و اليوم 20 أي خارم ميكروليتر) و واليوم 20 أيام القارنة بتضخيم . 3 مواقع هي 1280 و 201 و 202 و 202 أيام (20.5 ± 20.5) و 202 و 202 أيام و 201 و 205 و 202 أولتو و قاعر و المرميكروليتر) . والم ما يونوي أيام ما و 20 أيام ما والما رأم ما والأول الحمض النووي المسلمان الموي الما والترك المور والتر) . واليوم 20 كانت (20 ت 20 و 20 أول و 20 أم ما ميكروليتر) . وألم ما ما قان و الماميكروليتر) . واليوم 20 أيام والو و كان معدل الميد وام ما ما وروي أول و 20 أول و 20 أول و 20 أول و 20 أم ما أول و 20 أم ما أول و 20 أول و 20 أول ما ما أول و 20 أول والنه ما ما قال والم ما أول وال والتري أول ما ما وي وال والتر أول والو وو أول ما ما أول وول والم ما أول وال والم ما وول والم مام والو والما

الكلمات المفتاحية:STR Loci، FES، D8S1179،D18S51 ، أوتار الجيتار ، الجريمة.