



Effect Of Salicylic Acid In Growth And Flowering Of Bulbs .

Fanar Hashim Al-Hashemi¹

Zohoor Fuaad Abdaljabar¹

¹Department of Horticulture and Landscape Design, Collage of Agriculture and Forestry, Mosul university, Mosul, IRAQ.

*Corresponding Author: fanar@uomosul.edu.iq.

Received: 05/01/2024

Revised: 06/02/2024

Accepted: 10/02/2024

Published: 01/3/2024

ABSTRACT

Salicylic acid is one of the internal growth regulators derived from phenols and has multiple functions. this acid was found in *Salix* spp. (willow tree) and extracted naturally since 1928 by the scientist John Buchner, it's produced from the amino acid phenylalanine. *Salix* plant contains the salicin compound by 9-11%. It plays an important role in the growth and development of the plant for physiological effects, such as increasing the plant's response to stress conditions (biotic and abiotic). Salicylic acid is also involved in many bio processes, including growth, photosynthesis, cellular metabolism, protein synthesis, stomatal closure, gas exchange, and strengthening the defence system, which protects the plant against various diseases and increases the effectiveness of antioxidants and enzymes. It also stimulates the flowering, ion absorption, and nutrient transfer within the plant. Numerous experiments have indicated that spraying or soaking many bulbs with salicylic acid gave the best results for vegetative and flowering growth, as well as giving the best diameter and size of the bulbs formed for many plants such as gladiolus, tuberose, amaranth, lilium, tulipa, narcissus and lycoris.

Keywords: Salicylic acid, Growth, Bulbs.

Copyright © 2024. This is an open-access article distributed under the Creative Commons Attribution License.

INTRODUCTION

Plants are one of the world's richest sources of natural medicine, until the 19th century when the salicylates, including salicylic acid (SA), methyl salicylates, were isolated from extracts of different plants like willow [1]. Plants have specific mechanisms to resist various environmental stresses, which include the activation of antioxidant enzymes and non-enzymatic antioxidants such as carotenoids, ascorbic and salicylic acid, SA is a phenolic compound that inhibits ethylene production, the inhibitory actions of SA most closely resembled with that of dinitro phenol, a known inhibitor of ethylene forming enzymes, and the starting compound for salicylic acid for its production inside plants is cinnamic acid [2]. It was found throughout the plant kingdom and has been traced historically. It was used by the Greek physician Hippocrates in 400 BC, for its medicinal virtues; he prescribed it as a decoction of *Salix* to treat patients [3] [4]. Since 1859 Kolbe & Schmitt discovered the Kolb-Schmitt reaction to chemically synthesize SA from phenol, allowing for it to be produced on a large commercial scale, then in 1893 Felix Hoffman was the 1st to synthesise pure derivative acetylsalicylic acid (known as aspirin) which produces less gastrointestinal irritation yet has similar medicinal properties [5]. SA was extracted naturally since 1928 by the scientist John Buchner from the *Salix* (Latin name for the willow tree, from which the name was derived), and produced from the amino acid phenylalanine via Coumaric and benzoic acid [6].

SA was classified as an endogenous plant hormone because of its physiological roles in plant such as induction of flowering, regulating ions absorption, nutrient transfer, increasing the representation of CO₂ gas, protein synthesis, hormonal balance and stomatal movement, it also plays a role in regulating the response of plants to environmental stress condition and provides protection against them, such as abiotic stress (salt, heat, drought, chilling) and stress resulting from minerals, as well as biotic stress factor especially to *Fusarium*, SA affects the biosynthesis of ethylene and has an opposite role to the growth inhibitor "Absciscic acid " [1][7].

The importance of using salicylic acid in a plant's tolerance to abiotic stresses, especially drought, is due to its multiple roles in regulating plant metabolism [8]. It protects the plant from frost, drought, and salinity by resisting the formation of active free oxygen radicals (ROS), which destroy cell membranes by oxidizing unsaturated fatty acids and changing the effectiveness of antioxidant-resistant enzymes [9].

Several researchers have determined that salicylic acid causes an increase in vegetative activity and the positive effect of SA on growth and productivity may be attributed to its direct effect on other plant hormones, as it affects the rates of auxin and cytokinin, as well as its role to increase flowering rates by increasing the number of flower buds in comparison with vegetative buds [10]. It also works to accelerate the formation of chlorophyll, carotene and anthocyanin, photosynthesis by stimulating the formation of grana plates, development of chlorophyllase and inhibiting the action of the chlorophyllase enzymes, which naturally has a positive effect on the photosynthesis, enzyme activity. It has a role in the process of thermoregulation in some plants, as well as increasing plant resistance to diseases and insects [4] [11]. In addition, SA was

participated in increasing of metabolic rates which contributes to the energy of the plant through alternative pathways accompanied by a change in the level of nucleic and amino acids within the plant, SA has the ability to bind conjugate with some amino acids such as proline and arginine, which increase the plant's effectiveness in resisting environment stresses [12].

SA compound may also contribute to the signal regulation process during the practical of gene expression of leaf senescence in Arabidopsis plants [13]. So many studies have proven that spraying plants with salicylic acid gives a signal to the plant that it's under conditions of tension and stress, so this is what prompts the plant to increase its production of antioxidant enzymes and the most important of them catalase, peroxidase and super oxide dismutase (SOD), these enzymes were the first lines of defense for the plant cell, they also limit damage resulting.

From stress and encourage the plant to regrow [14] [15]. Catalase converts hydrogen peroxide into oxygen and water, the SOD is the plant cells defender against free radicals, peroxidase contributes to reducing oxidation reactions [16]. During recent years, this compound has gained the attention of researchers due to its ability to induce systemic acquired resistance (SAR) in plants when attacked by many pathogens, as this leads to the production of defence proteins [17].

The relationship between salicylic acid and the growth and development of plants was revealed by [18] on Gladiolus, [19] on Tuberose, [20] on Red amaranth, [21] on Lilium. Salicylic acid is an aromatic carboxylic acid of a phenolic nature, a low molecular weight and is colorless, dissolves in water and polar organic Solvents, it is formed by the process of removing the carboxyl group from Trans-cinnamic acid to form benzoic acid and by the action of the enzyme (Benzoic acid 2-hydroxylase), salicylic acid was produced, and its chemical formula is $C_6H_6O_3$ [22]. The effectiveness of salicylic acid used to spray the plants depends on the type of plant, the time of addition, and the concentration used, in this regard, [23] has been noted that when treatment of *Polianthus tuberosa* with salicylic acid by foliar spraying every 20 day at concentration of 0,50, 100 and 200 ppm, it led to recording the maximum duration of flowering the second floret, most extended length of flower stalk, number of leaves per plant, the chlorophyll content, number of florets and its diameter, vase life at concentration of 200 ppm. The bulbs of *Tulipa* sp. were soaked with SA at a levels (0, 250, 500, 1000) ppm and then planted in (green house, outdoor) conditions, the results indicated that the bulbs were grown in the green house had given the first emergence, lowest number of days for flowering moreover the flower length, flower stalk length, flower life was significantly superior with those grown outdoor when treated with SA at a level 1000 ppm [24].

[25] mentioned that the lower concentration of SA 150 ppm had a longer vase life of *Narcissus tazetta* L. than 300 ppm, which had more effective in increasing fresh weight of cut flowers and water uptake is due to stress-relieving and acidifying properties of salicylic acid. [26] indicated that spraying of three cultivars of *Gladiolus hortulanus* L. with SA at 250 ppm had a significant effect of showing the highest values for the weight, diameter and volume of the corms from the red variety and more plant height, leaf area, dry weight of leaves, number and diameter of florets, length of inflorescence, duration of flowering as well as the diameter and their dry weight and total sugars accumulated in the corms, total chlorophyll content of leaves and the percentage of dry matter when sprayed with SA at four concentration (0, 50, 100, 150) ppm. It's significantly 150 ppm to give the best characteristics, compared to the two varieties of white and purple flowers. These results illustrate that SA may be involved in many processes, i.e. morphogenetic processes like the formation of flowers and tubers, photosynthesis process results, and more dry total sugars accumulated in the corms [27]. More dry matter productively, according to [28] examined the effect of SA in delaying the dissolution of plant organs, especially (leaves and flowers), which automatically helps the site of photosynthesis remain effective for a more extended period, it prevents the risk of falling flowers and immature fruits, which leads to increased yield. [29] concluded in a study evaluating the development of *Lycoris radiata* and its carbohydrate content and enzyme activity when treated with three growth regulators, including SA sprayed on the shoots at different concentration, the result showed that SA was a stimulant the growth and development of the bulbs, increasing the carbohydrate content, enzyme activity and bulb's size, in addition to the role of the β -amylase enzyme in starch metabolism in bulbs. In a study on the response of Gladiolus corms to soaking with two levels of salicylic acid (100, 150) ppm for 24 hours, there was a significant superiority at the concentration 150 ppm in the time for emergence of 50% of the corms, most significant number of branches per plant, most extended period for the inflorescences to remain in good coordination, and the largest (number, diameter, length) of the inflorescence as well as vase life [30].

Also, spraying this hormone at 200 ppm leads to a significant increase in plant height, leaves per plant, leaf area, total chlorophyll, as well as days to appearance of flower buds, days to colour visibility in buds and also increases the stalk length, days for sprouting of *Lilium asiatic* hybrid tresor [31]. [32] who reported better seedling growth with SA application on the trait of day to flowering, which was highly significant. It occurred earlier in SA 2% compared with the control, which had a few days' delay in flowering of *Crocus sativus* L.

Conclusion

Salicylic acid plays a role in improving most of the characteristics of vegetative, flowering growth and bulbs, due to its effect increasing the level of cell division within the apical tissues responsible for increasing growth, including plant height, number of branches, and leaf area. It also works to prevent the oxidation of internal hormones through its direct effect in encouraging and stimulating auxins to stimulate meristematic cell division from the beginning of root growth with increasing the number of root hairs and their percentage, thus improving the characteristics system, which reflects positively on increasing vegetative growth rate. In addition to the role of hormonal acid in the accumulation of dry matter and increasing the dry weight of the plant and the activity of β -glucosidase enzyme. SA affects the acceleration of the formation of Photosynthetic pigments

(chlorophyll) which increases the efficiency of photosynthesis and this is reflected positively in increasing the activity of biological processes.

References

- [1] Davies, P.J. (2010). plant hormones biosynthesis sign transduction action. Kluwer Academic. Publishers, Netherlands.
<http://dx.doi.org/10.1007/978-1-4020-2686-7>
- [2] Abdoallah, H and M, Hatami, Mahmood (2012). Efficiency of salicylic acid delay petal senescence and extended quality of cut spikes of gladiolus grandiflora cv 'wing's sensation'. African Journal of Agricultural Research, 7(4):540-545..
<https://shorturl.at/joBEU>
- [3] Rainsford, K. D. (Ed.). (2004). Aspirin and related drugs. CRC Press. Boca Raton, USA.
<https://shorturl.at/ajmzY>
- [4] Hedden, P., and Thomas, S. (Eds.). (2006). Plant Hormone Signaling. Annual Plant Reviews, Volume 24. Blackwell Pub.
<https://shorturl.at/AIT08>
- [5] Mahdi, J. (2010). Medicinal potential of willow: A chemical Perspective of aspirin discovery. J. of Saudi Chemical Society ,14(3), 317-322.
<https://www.sciencedirect.com/science/article/pii/S1319610310000578>
- [6] Anket, S; Gagan F, Aranit and M, Brestic L, Marco (2020). The role of salicylic acid in plants exposed to heavy metals. Molecules, 22(2):540-560.
<https://www.mdpi.com/1420-3049/25/3/540>
- [7] Eman, S; R. Amira and M, Moubarak (2018). Effect of methyl jasmonate and salicylic acid on the production of gladiolus grandifloras L. Nature and Science, 16(6) 40-47.
<https://shorturl.at/qDPV1>
- [8] Muthulakshmi, S., and Lingakumar, K. (2017). Role of salicylic acid (SA) in plants—A review. Int. J. Appl. Res, 3(3), 33-37. <https://shorturl.at/fjTZ0>
- [9] Parashar, A.; Yusuf, M. and Fariduddin, Q., & Ahmad, A. (2014). Salicylic acid enhances antioxidant system in Brassica juncea grown under different levels of manganese. International journal of biological macromolecules, 70, 551-558.
<https://www.sciencedirect.com/science/article/pii/S0141813014004863>
- [10] Kumar, P.; Dube, S. D., and Chauhan, V. S. (1999). Effect of salicylic acid on growth, development and some biochemical aspects of soybean (*Glycine max* L. Merrill). Indian Journal of Plant Physiology, 4(4), 327-330.
https://link.springer.com/chapter/10.1007/1-4020-5184-0_1
- [11] Hayat, S and A. Ahmad (2007). Salicylic acid biosynthesis metabolism and physiological role in plants. Springer Netherland.
<http://ndl.ethernet.edu.et/bitstream/123456789/70805/1/4.pdf#page=14>
- [12] Janda, M.; Ei Ruelland and M. Magical (2014). Salicylic acid signaling, Environmental and Experimental, Botany, In press.
<https://www.frontiersin.org/articles/10.3389/fagro.2021.781027/full>
- [13] Srihunsu, M.; N., Vannatime; C, Somruthai; A and paemane (2023). Comparative analysis of salicylic acid levels and gene expression in resistant, tolerant, and susceptible cassava varieties following whitefly-mediated SLCMV infection.
<https://shorturl.at/hnH89>
- [14] Noreen, S.; Ashraf, M.; Hussain, M., and Jamil, A. (2009). Exogenous application of salicylic acid enhances antioxidative capacity in salt stressed sunflower (*Helianthus annuus* L.) plants. Pak. J. Bot, 41(1), 473-479.
<https://shorturl.at/aEJWX>
- [15] Sedghi M.; H. K. Basiri and R.S. Sharifi (2013). Effect of activity salicylic acid on the antioxidant enzymes at in Sunflower. Annals of West Univ of Timisoara, Ser, Biology, 18(2):67-72.
<https://shorturl.at/jkmRT>
- [16] EL-Beltagi, H. S; A. A.; Mohamed, B. and Mekki, B. El-Din (2011). Differences in some Constituents, enzymes activity and electrophoretic characterization of different rapeseed (*Brassica napus* L.) cultivars. Tom, 118(1):45-52.
<https://shorturl.at/lzAP6>
- [17] Rai, K. K.; Pandey, N.; Rai, N.; Rai, S. K., and Pandey-Rai, S. (2021). Salicylic acid and nitric oxide: insight into the transcriptional regulation of their metabolism and regulatory functions in plants. Frontiers in Agronomy, 3, 781027.
<https://www.frontiersin.org/articles/10.3389/fagro.2021.781027/full>
- [18] Sajjad, Y.; M. J. Jaskain; M.X. Ashraf and Ahmad (2014). Response of morphological and physiological growth attributes to foliar application of plant growth regulators in Gladiolus "whit Prosperity", pak. J. Agri. Sci., 51 (1) 123-129.
<https://bit.ly/3waulo9>

- [19] Anwar; M.; H.A. Sahito, I. Hassan and A. Hussain (2014). Effect of pre harvest treatment of salicylic acid on growth and vase life of tuberose with aroma environment. *Wudpecker Journal of Agris Res*, 3(2):050-57. <https://shorturl.at/epxV6>
- [20] Khandaker, L.; Akond, A. M., and Oba, S. (2011). Foliar application of salicylic acid improved the growth, yield and leaf's bioactive compounds in red Amaranth (*Amaranthus tricolor*). *vegetable crops research bulletin* 74:77-86. <https://shorturl.at/gIR38>
- [21] Hajizadeh, H. S., and Aliloo, A. A. (2013). The effectiveness of pre-harvest salicylic acid application on physiological traits in lily (*Lilium longiflorum* L.) cut flower. *International Journal of Scientific Research in Environmental Sciences*, 1(12), 344. <https://shorturl.at/boJ78>
- [22] Lee, H.; J. Leon. and L. Raskin (1995). Biosynthesis and metabolism salicylic acid, *Proc. Natl. Acad Sci.*, 92: 4076-4079. <https://www.pnas.org/doi/abs/10.1073/pnas.92.10.4076>
- [23] Khodakhah, B.; Nabigol, A., and Salehi, B. (2014). The effect of different levels of humic acid and salicylic acid on growth characteristics and qualities of tuberose. *Advances in Environmental Biology*, 8(16), 118-123. <https://www.cabdirect.org/cabdirect/abstract/20153262115>
- [24] Cavusoglu, a.; özer-uyar, g. E., and ünal, f. (2022). An abiotic stress factor: phytotoxicity of high doses of salicylic acid on growth parameters of *Tulipa* sp. Under glasshouse and outdoor conditions. *Applied ecology and environmental research*, 20(6), 4925-4934. <https://bit.ly/3HKnG6L>
- [25] Sefa, G. Ü. N., and Ö. Burhan. (2020). Effects of salicylic acid and citric acid treatments on some parameters of *Narcissus tazetta* L. during vase life. *Journal of Agricultural Faculty of Gaziosmanpaşa University (JAFAG)*, 37(3), 202-207. <https://dergipark.org.tr/en/download/article-file/2117124>
- [26] Saja, S. L. and O. Ammar (2022). Effect of spraying with salicylic acid and calcium chloride on the yield characteristics of corm and cormlets for three cultivars of *Gladiolus X hortulanus* L. *Mesopotamia journal of Agriculture*, 50(3): 59-69. <https://www.iasj.net/iasj/download/3e1be11ad2c5ecc1>
- [27] Sewedan, E.; Osman, A. R., and Moubarak, M. (2018). Effect of methyl jasmonate and salicylic acid on the production of *Gladiolus grandifloras*, L. *Nature and Science*, 16(6), 40-47. <https://shorturl.at/tACO6>
- [28] CAO, Li; Xi, Yang; Xu, Yue and yang (2021). Transcriptional profiles underlying the effects of salicylic acid on fruit ripening and senescence in pear (*Pyrus pyrifolia* Nakai). *Journal of Integrative Agriculture*, 20(9):2424-2437. <https://bit.ly/491SuMI>
- [29] Ya-main, X.; S. Lin - fang; Xiav and X. Yi-ping (2013). Effect of three plant growth regulators on the bulblets development of *Lycoris radiata*. *Journal of Nuclear Agricultural Sciences*, 27 (9): 1409-1415. <https://www.hnxb.org.cn/EN/10.11869/hnxb.2013.09.1409>
- [30] Pawar, A.; Chopde, N., and Nikam, B. (2018). Thiourea and salicylic acid influences growth, yield and quality of gladiolus. *Journal of Pharmacognosy and Phytochemistry*, 7(5), 970-972. <https://shorturl.at/iFHLT>
- [31] Pahare, P.; Scholar, R., and Beura, S. (2022). Impact of salicylic acid and humic acid on vegetative and flower bud production of *Lilium asiatic* hybrid Tresor. ~ 2185 ~ *The Pharma Innovation Journal*, 11(7), 2185–2190. www.thepharmajournal.com
- [32] Rastegari, S.; Naser Alavi, S. M., and Mohayeji, M. (2022). Effect of Salicylic Acid and Pre-Cold Treatment on Flower Induction in Saffron. *Scientifica*, 2022. <https://doi.org/10.1155/2022/6108161>

تأثير حامض السالسيليك في نمو وازهار الأبصال (مقالة مراجعة).

فنان هاشم يوسف الهاشمي¹ زهور فؤاد عبد الجبار¹

قسم البستنة وهندسة الحدائق، كلية الزراعة والغابات، جامعة الموصل، الموصل، العراق¹.

الخلاصة

حامض السالسيليك هو أحد منظمات النمو الداخلية المشتق من الفينولات ذات الوظائف المتعددة، يتواجد هذا الحامض في شجرة الصفصاف *Salix spp*. واستخلص طبيعياً منذ عام 1928 على يد العالم جون بوشنر وتم إنتاجه من الحامض الأميني فينيل ألانين. يحتوي نبات الصفصاف على مركب السالسين بنسبة 9-11%. يلعب حامض السالسيليك دوراً هاماً في نمو وتطور النبات لتأثيره الفسيولوجي في زيادة استجابة النبات لظروف الإجهاد (الاحيائية واللاحيائية). كما يشارك حامض السالسيليك في العديد من العمليات الحيوية منها النمو والتمثيل الضوئي والتمثيل الغذائي الخلوي وتخليق البروتينات، وإغلاق الثغور وتبادل الغازات، كما أنه يقوي نظام الدفاع ويحمي النبات من الأمراض المختلفة ويزيد من فعالية مضادات الأكسدة والإنزيمات. ويعمل أيضاً على تحفيز التزهير وامتصاص الأيونات ونقل العناصر الغذائية داخل النبات. لقد أشارت التجارب إلى أن رش أو نقع العديد من الأبصال بحامض السالسيليك قد أعطى أفضل النتائج للنمو الخضري والزهري، كما أعطى أفضل قطر وحجم للأبصال المتكونة للعديد من النباتات مثل الكلاديولس ومسك الروم والقطفة الدمية والليليم والتوليب والرنجس وزنبق العنكبوت الأحمر.

الكلمات المفتاحية : حامض السالسيليك، النمو، الأبصال.