

Synthesis and Characterization of some Substituted 1, 2, 5-oxadiazine Derived from 9H- carbazole

Maha M. Tawfiq

Salim J. Mohammed

Department of Chemistry/ College of Science/ University of Mosul

p-ISSN: 1608-9391

e-ISSN: 2664-2786

Article information

Received: 25/ 11/ 2022

Revised: 28/ 1/ 2023

Accepted: 8/ 2/ 2023

DOI: 10.33899/rjs.2023.181265

corresponding author:

Maha M. Tawfiq

Chemimaha96@gmail.com

ABSTRACT

Hydrazones possess unit an azomethine (-NHN=CH) group and are considered derivatives of aldehydes and ketones compounds in which the oxygen atom has been replaced by the NNH_2 . In this paper, we study synthesis of the novel of some new 1, 2,5-oxadiazine derivatives were prepared starting with carbazole (1) as a synthon, which on reaction with chloroacetyl chloride in absolute ethanol yielded ethyl 1-(9H-carbazole-9-yl)2-chloroethanone (2). This compound (2) on reaction with hydrazine hydrate giving 1-(9H-carbazol-9-yl)-2-hydrazinylethanone (3) followed treatment the later compound with either substituted benzaldehyde or substituted acetophenone will give substituted hydrazones (4a-e) and (5a-d) respectively. These hydrazone compounds on were allowed to react with an acetic acid anhydride to give the corresponding 1, 2, 5-oxadiazine compounds (6a-e) and (7a-d). The purity of substituted hydrazones and the corresponding 1,2,5-oxadiazine compounds were tested by thin-layer chromatography (TLC). All newly synthesized compounds in this study were confirmed by physical and spectral (FT-IR, nuclear magnetic resonance (^1H NMR and ^{13}C NMR) analysis.

Keywords: Carbazole, Hydrazones, Schiff bases, Oxadiazine, Acetic Anhydride.

INTRODUCTION

Carbazole has a tricyclic structure, consisting of two benzene rings located on either side of a five-membered nitrogen-containing ring (Zhang *et al.*, 2010). The structure of the compounds is based on that of indole, but in which a second benzene ring is fused to the arcuate ring at position 2-3 of indole (Fattorusso and Tagliatela, 2008). Many researches have been conducted on carbazole and its derivatives. The carbazole molecule consists of two six-membered benzene rings fused on either side by a five-membered nitrogen-containing ring (Nandy *et al.*, 2014). Thus, hydrazones are types of Schiff bases and very important chemical compounds because of their ease of obtaining and preparing them directly, as well as to their stability towards hydrolysis. Hydrazones were prepared from the condensation of substituted aldehydes or ketones with hydrazine derivatives and have the general formula $R-NH-N=C-R_1$ (Xin Su and Ivan Aprahamian, 2014). Hydrazones and their derivatives are also very important compounds in organic and medicinal chemistry due to their biological activity which includes: anticancer (Yang *et al.*, 2021), antiviral (Alam *et al.*, 2014) (Popiolek *et al.*, 2020), antibacterial (Al-Daher and Omer, 2019), antifungal (Siemann *et al.*, 2002; Rollas *et al.*, 2007; Aslan *et al.*, 2012; Backes *et al.*, 2014; Alsaeedi *et al.*, 2015; Ghiya *et al.*, 2016; Segretti *et al.*, 2016; Bhat *et al.*, 2018; Zhou *et al.*, 2019; Shaaban *et al.*, 2020; Mohammed, 2022), antidepressant properties (Abid *et al.*, 2017) and in the treatment of many diseases such as Alzheimer's disease and infections (Murtaza *et al.*, 2016 and Fernandes *et al.*, 2017). Also, some hydrazones are used as ligands in preparing some complexes (Al-Daher and Mohammed, 2018).

Oxadiazine are the molecules with oxygen and two nitrogen's resting in different positions of the cyclic structure. six-membered heterocyclic rings having three heteroatoms are very less in nature, but they possess various biological applications. Derivatives of 1,2,5-oxadiazine exhibited various biological applications such as anti-inflammation (Sava *et al.*, 1985 and Xia *et al.*, 2008) against malaria (Eissa *et al.*, 2019) and tuberculosis (Holdiness, 1987; Zheng *et al.*, 2009; Ajani *et al.*, 2010) anti-cancer (Mohareb *et al.*, 2010) plant growth regulator, anti-convulsive activity, antibacterial, nematocidal, cardiovascular, anti-HIV. Oxadiazine derivatives are used as an intermediate to synthesized some of the β -lactam antibiotics or prodrugs (Shet *et al.*, 2010; Mohareb and Schatz, 2011; Shubakara *et al.*, 2014; Gurlananappa and Kariappa, 2017).

EXPERIMENTAL

Chemicals and solvents used during the research are commercially available, Aldrich and Fluka companies. Melting points are recorded in open capillaries tubes by using Stuart SMP30 melting point apparatus. Infrared spectra were recorded by using (Shimadzu FT-IR-ATR) (Japan) infrared spectrophotometer. Measurements were performed in Salahaddin University, Erbil. 1H -NMR spectra were recorded by using (Bruker 400MHZ) with tetramethyl silane as internal standard in ((DMSO- d_6)).

Synthesis of 1-(9H-carbazole-9-yl) chloroethanone (2) (Al-Majidi, 2013)

Dissolved (0.002 moles) of carbazole (1) in dry acetone (40 ml) then (0.002 moles) of chloroacetylchloride was added. The mixture was refluxed for (6hrs.), and then solvent was evaporating under reduced pressure. The resulting solid was dried, to give the precipitate a clear color (brown), m.p. (138-140 °C), yield (75%).


Synthesis of 1-(9H- carbazole- 9- yl) -2- hydrazinylethanone (3) (Al-Majidi and Al-Quaz, 2010)

Dissolved (0.001 moles) of 1-(9H-carbazol-9-yl)-2-chloroethan-1-one (2) in absolute ethanol (25 ml), with (20 ml) of hydrazine hydrate (85%) was added. The mixture was refluxed for (5hrs.), and then the solvent was evaporated to half volume. The formed precipitate was isolated by filtration, dried to give pure compound with a Light brown clear color, m.p. (231-232 °C), yield (80%).

Synthesis of benzylidene hydrazinyl and phenylethylidene hydrazinyl (4a-e) (5a-d) (Al-Kaissy *et al.*, 2013)

A mixture of (0.001moles) of hydrazinylethanone (3) and (0.001moles) of substituted benzaldehyde or acetophenone were dissolved in (30ml) of absolute ethanol. A few drops of glacial acetic acid were added. The mixture was refluxed for about (4hrs.), and then the solvent was evaporating under reduced pressure. The resulting solid was dried, to give precipitate of hydrazones (4a-e) (5a-d) respectively. The physical data are recorded in (Table 1).

Table 1: Physical Properties for compounds (4a-e) (5a-d)




Compd. No.	R	X	m.p °C	Yield %	Color
4a	H	P-Br	243-246	51	Light brown
4b	H	m-NO ₂	173-179	42	Gold
4c	H	o-Cl	186-188	58	Olive
4d	H	3,4-OH	140-143	94	Light green
4e	H	2,3-OCH ₃	136-138	83	Brown
5a	CH ₃	p-NH ₂	112-113	97	Dark brown
5b	CH ₃	p-CH ₃	197-199	91	Brown
5c	CH ₃	o-OH	211-214	57	Light brown
5d	CH ₃	m-OCH ₃	204-206	55	Light brown

Synthesis of substituted 1,2,5-oxadiazine compounds (6a-e) (7a-d) (Joshi *et al.*, 2008)

Dissolved (0.001moles) of hydrazones (4a-e) (5a-d) in (15ml) of acetic anhydride. The mixture was heated in refluxed system for (4hrs.), at 100 °C. The resulting mixture poured on crushed ice with stirring. The formed precipitate was isolated by filtration, washed with water, dried, to give precipitate of 1,2,5-oxadiazine compounds (6a-e) (7a-d) respectively. The physical data are recorded in (Table 2).


Table 2: Physical Properties for compounds (6a-e) (7a-d)




Compd. No.	R	X	m.p °C	Yield %	Color
6a	H	P-Br	57-59	62	Grey
6b	H	m-NO ₂	85-87	54	Gold
6c	H	o-Cl	197-199	57	Light brown
6d	H	3,4-OH	112-114	42	Light brown
6e	H	2,3-OCH ₃	191-193	44	Dark brown
7a	CH ₃	P-NH ₂	61-63	43	Olive
7b	CH ₃	p-CH ₃	205-207	51	Light brown
7c	CH ₃	o-OH	72-74	52	Dark grey
7d	CH ₃	m-OCH ₃	55-57	94	Bright grey

RESULTS AND DISCUSSION

In this research, reaction of carbazole (1) with chloro acetylchloride to give an ester namely 1-(9H-carbazole-9-yl) chloroethanone (2) followed on reaction with hydrazine hydrate afforded the corresponding hydrazide (3) as shown in the scheme (1).



Treatment of hydrazide (M3) with substituted benzaldehyde or acetophenone gave the corresponding hydrazones (M4a-e) (M5a-d) respectively, which on reaction with acetic anhydride giving the corresponding 1, 2, 5-oxadiazine compounds (6a-e) and (7a-d) as shown in scheme (2).



Scheme 2: Synthesis of hydrazones and 1,2,5-oxadiazine compounds

The structures of the target hydrazones (4a-e) (5a-d) were elucidated using (FT-IR, ¹H-NMR, ¹³C-NMR). The FT-IR spectra for compounds (4a-e) (5a-d) showed the following stretching bands; (1620 -1681 cm⁻¹) due to the (C=N) bond for hydrazones, (1740-1810 cm⁻¹) for (C=O) group, (3008-3418 cm⁻¹) for the (NH) bond (NH) group. The ¹H-NMR spectra for compounds (4a, 4e, 5a) in (DMSO-d₆) representation for these hydrazones showed the peaks for (CH=N, NH, and aromatic protons) and other groups in these compounds is in full agreement with the proposed structures which confirmed in (Table 3). While ¹³C-NMR for compounds (4a, 4e, 5a) in (DMSO-d₆) representation for these hydrazones showed the signal for carbon atoms in these compounds as shown in (Table 4).

Table 3: FT-IR data of compounds (4a-e) (5a-d)

Compd. No.	C=N	C=O	NH	Other
4a	1622	1790	3049	C-Br (570)
4b	1640	1780	3049	NO ₂ (1519) asym (1340) sym
4c	1681	1775	3417	C-Cl(756)
4d	1653	1790	3051	OH(1394)
4e	1625	1792	3008	OCH ₃ (1068)
5a	1647	1730	3226	-----
5b	1620	1740	3418	-----
5c	1645	1755	3049	OH (1327)
5d	1643	1750	3149	-----

Table 4: The ^1H -NMR and ^{13}C -NMR spectral data of compounds (4a,4e,5a)




Compd. No.	Structure	^1H -NMR, (ppm), DMSO-d_6	^{13}C -NMR, (ppm), DMSO-d_6
4a		3.468(s,2H, CH_2), 7.138-8.716(m,13H,ArH+CH), 11.254(s,1H,NH).	168.3(C=O), 51.4(CH_2NH), 144.3(N=CH), 115.6, 116.3, 118.1, 119.2, 120.3, 121.4, 131.7(ArC).
4e		11.245(s,1H,NH), 8.871(s,1H,CH), 7.135-8.871[m, 11H, ArH], 3.903(s,6H,2OCH ₃).	165.4(C=O), 50.5(CH_2NH), 143.3(N=CH), 55.1-60.3(OCH ₃), 111.2, 113.5, 116.9, 119.4, 119.8, 132.6, 153.5(ArC).
5a		1.049(s,3H, CH_3), 3.461(bs,2H, CH_2), 5.538(bs,2H,NH ₂), 7.137-8.121(m,12H,ArH), 11.254, [s,3H,NH],	196(C=O), 51.8(CH_2NH), 167.8(N=C), 17.2(C-CH ₃), 151.8(C-NH ₂), 11.3, 116.5, 118.8, 119.6, 120.4, 121.3, 122.6, 130.0(ArC).

The structure of compounds (6a-e) (7a-d) have been identified based on their FT-IR, ^1H -NMR, and ^{13}C -NMR. The FT-IR spectra were characterized by the presence at the range ($1224\text{-}1301\text{cm}^{-1}$) due to (C-N) bands, the range ($1645\text{-}1695\text{ cm}^{-1}$) due to (C=N) bands also at the range ($1700\text{-}1790\text{ cm}^{-1}$) refer to (C=O) amide groups besides at the range ($1004\text{-}1076\text{ cm}^{-1}$) due to (C-O-C) bands. The assignment of the vibration ν (cm^{-1}) of the IR absorption bands spectra was illustrated in (Table 5). Whereas the chemical shifts that appeared in the ^1H -NMR show signals due to exhibition of protons of (N-CH) as singlet at the range (5.0 -5.2 ppm) and at the range (2.04-3.19 ppm) of (N-C=O-CH₃) in addition the other obtained values for compounds (6a-e) (7a-d) are listed in (Table 6). In the ^{13}C -NMR spectra of compounds (6a-e) (7a-d) showed different signals for all carbon atoms which mentioned in (Table 6). The range σ (90-97ppm) due to (N-CH), σ (152.7-159.3 ppm) assigned to (C=N-N). The carbons atom of carbonyl group was appeared at the range σ (166.5-169.7 ppm) while the signals of the carbon of aromatic ring (Ar-C) all signal appears.

Table 5: FT-IR data of compounds (6a-e) (7a-d)

Compd. No.	C-N	C=N	C=O	C=C	C-O-C	Others
6a	1238	1685	1762	1595	1024	C-Cl (794)
6b	1236	1680	1721	1595	1020	NO ₂ (1521) asym (1342) sym
6c	1240	1688	1750	1595	1022	C-Cl (725)
6d	1236	1695	1747	1590	1076	OH (1375)
6e	1298	1691	1757	1593	1004	OCH ₃ (1068)
7a	1301	1681	1770	1595	1014	NH(3061) asym (928)sym
7b	1238	1668	1750	1597	1028	—
7c	1238	1645	1700	1590	1025	OH (1390)
7d	1224	1683	1790	1594	1045	OCH ₃ (1045)

Table 6: The ^1H -NMR and ^{13}C -NMR spectral data of compounds (6a, 6e and 7a)

Com pd. No.	Structure	^1H -NMR, (ppm), DMSO- d_6	^{13}C -NMR, (ppm), DMSO- d_6
6a		1.917(s,3H,CH ₃),5.23(bs,1H,CH),6.28(bs,2H,CH ₂),7.409-8.836 (m, 12H, -ArH),	92.5(CH),164.7(C=N-N), 168.59(N-C=O), 110.4,111.3,113.4,116.2,117.8,118.3,119.2,123.4,127.3 ,130(Ar-C).
6e		1.915(s,3H,CH ₃),3.805(s,6H,2OCH ₃),7.38-8.40 [m, 12H, ArH,], 5.03 [s,1H,CH]CH=N) .	94(CH), 166.4(C=N-N), 168.8(N-C=O), 117-130(Ar-C), 23.4(CH ₃),19.1(Ar-CH ₃)
7a		1.914(s,3H,CH ₃),2.089(s,3H,CH ₃),4.56(bs,2H,CH ₂),7.412-8.282(m,12H,ArH).	90(N-CH), 154.7(C=N-N), 167(N-C=O), 115-127(Ar-C), 22.9(CH ₃).

CONCLUSION

We conclude that in this research we studied the synthesis of some new 1,2,5-oxadiazine derivatives that were prepared starting with carbazole, which when treated with chloroacetyl chloride gave the ester (2). When reacted with hydrazine hydrate, the hydrazide(3) compound was prepared, which was reacted with either substituted benzaldehyde or substituted acetophenone, giving us hydrazone-hydrazide. When these compounds are reacted with acetic acid anhydride, the final products are obtained, represented by the compounds 1, 2, 5-oxadiazine, corresponding . The purity of all compounds prepared in the study was tested by thin layer chromatography (TLC), and finally these compounds were identified by physical and spectroscopic analysis.

ACKNOWLEDGEMENTS

The authors are thankful to Department of Chemistry, College of Science, University of Mosul for Cooperation to complete this work.

REFERENCES

- Abid, S.M.A.; Younus, H.A.; Al-Rashida, M.; Arshad, Z.; Maryum, T.; Gilani, M.A.; Iqbal, J. (2017). Sulfonyl hydrazones derived from 3-formylchromone as non-selective inhibitors of MAO-A and MAO-B: Synthesis, molecular modelling and in-silico ADME Evaluation. *Bioorg. Chem.*, **75**, 291-302. <https://doi.org/10.1016/j.bioorg.2017.10.001>
- Ajani, O.O.; Obafemi, C.A.; Nwinyi, O.C.; Akinpelu, D.A. (2010). Microwave assisted synthesis and antimicrobial activity of 2-quinoxalinone-3-hydrazone derivatives. *Bioorg. Medic. Chem.*, **18**(1), 214-221. <https://doi.org/10.1016/j.bmc.2009.10.064>
- Al-Daher, A.M.; Mohammed, A.H. (2018). Preparation, characterization and study of ethyl pyruvate aroyl hydrazone metal complexes. *Rafi. J. Sci.*, **27**(2), 103-110.
- Al-Kaissy, W.W.N.; Tuama, S.H.F.; Al-Majidi, S.M.H. (2013). Synthesis, characterization and evaluation of antimicrobial activity of some new acetylenic amine and 2- oxoazetidine of carbazole. *Am. J. Sci. Ind. Res.*, **4**(4), 389-398. <https://doi.org/10.33899/rjs.2018.145395>

- Al-Majidi, S.M.H. (2013). Synthesis, characterization and evaluation of antimicrobial activity of several new n-substituted carbazole. *J. Al-Nahrain University*, **16**(4), 67-79. <https://doi.org/10.22401/jnus.16.4.08>
- Al-Majidi, S.M.H.; Al-Quaz, A.M.N. (2010). Synthesis of some new-substituted-1,2,3,4-tetrahydrocarbazole derivatives and study their biological activity. *J. Al-Nahrain University*, **13**(1), 26-39. <https://doi.org/10.22401/jnus.13.1.04>
- Alsaedi, H.S.; Aljaber, N.A.; Ara, I. (2015). Synthesis and investigation of antimicrobial activity of some nifuroxazide analogues. *Asian J. Chem.*, **27**(10), 3639. <https://doi.org/10.14233/ajchem.2015.18896>
- Aslan, H.G.; Özcan, S.; Karacan, N. (2012). The antibacterial activity of some sulfonamides and sulfonyl hydrazones, and 2D-QSAR study of a series of sulfonyl hydrazones. *Spectroch. Acta Part A: Molec. and Biomole. Spectrosc.*, **98**, 329-336. <https://doi.org/10.1016/j.saa.2012.08.043>
- Backes, G.L.; Neumann, D.M.; Jursic, B.S. (2014). Synthesis and antifungal activity of substituted salicylaldehyde hydrazones, hydrazides and sulfohydrazides. *Bioorg. Medic. Chem.*, **22**(17), 4629-4636. <https://doi.org/10.1016/j.bmc.2014.07.022>
- Bashir, M.; Bano, A.; Ijaz, A.S.; Chaudhary, B.A. (2015). Recent developments and biological activities of N-substituted carbazole derivatives: a review. *Molecules*, **20**(8), 13496-13517. <https://doi.org/10.3390/molecules200813496>
- Bhat, M.; Poojary, B.; Kumar, S.M.; Hussain, M.M.; Pai, N.; Revanasiddappa, B.C.; Kullaiah, B. (2018). Structural, crystallographic, Hirshfeld surface, thermal and antimicrobial evaluation of new sulfonyl hydrazones. *J. Molec. Struct.*, **1159**, 55-66. <https://doi.org/10.1016/j.molstruc.2018.01.041>
- Eissa, F.M.; Selim, M.; Sayed, A.; Sharkawi, M. (2019). Green synthesis and antibacterial activity of pyrido [2, 3-e] 1, 3, 4-oxadiazine derivatives. *American J. Heterocyclic Chem.*, **5**(3), 71-75. <https://doi.org/10.11648/j.ajhc.20190503.13>
- Fattorusso, E.; Taglialatela-Scafati, O. (2008). "Modern Alkaloids". Wiley-Vch Verlag Gmbh and Co. Kga, 8th ed., 481p. <https://doi.org/10.1002/9783527628155>
- Fernandes, T.B.; Cunha, M.R.; Sakata, R.P.; Candido, T.M.; Baby, A.R.; Tavares, M.T.; Parise-Filho, R. (2017). Synthesis, molecular modeling, and evaluation of novel sulfonylhydrazones as acetylcholinesterase inhibitors for alzheimer's disease. *Archiv. der Pharm.*, **350**(11), 1700163. <https://doi.org/10.1002/ardp.201700163>
- Ghiya, S.; Joshi, Y.C. (2016). Synthesis and antimicrobial evaluation of hydrazones derived from 4-methylbenzenesulfonohydrazide in aqueous medium. *Med. Chem. Research*, **25**(5), 970-976. <https://doi.org/10.1007/s00044-016-1543-9>
- Gurlananjappa, P.A.; Kariyappa, A.K. (2017). An easy procedure for synthesis of 1, 3, 4-oxadiazines: A potential antimicrobial agents. *Asian J. Chem.*, **29**(8), 1687-1689. <https://doi.org/10.14233/ajchem.2017.20540>
- Holdiness, M.R. (1987). A review of blood dyscrasias induced by the antituberculosis drugs. *Tubercle*, **68**(4), 301-309. [https://doi.org/10.1016/0041-3879\(87\)90072-9](https://doi.org/10.1016/0041-3879(87)90072-9)
- Joshi, S.D.; Vagdevi, H.M.; Vaidya V.P.; Gadaginamath, G.S. (2008). Synthesis of new 4-pyrrol-1-yl benzoic acid hydrazide analogs and some derived oxadiazole, triazole and pyrrole ring systems: A novel class of potential antibacterial and antitubercular agents. *Eur. J. Med. Chem.*, **43**, 1989- 1996. <https://doi.org/10.1016/j.ejmech.2007.11.016>
- Mohammed, A.B. (2022). Synthesis and antibacterial evaluation of novel hydrazones and bis-hydrazones containing 1,2,3-triazole moiety. *J. Taibah University for Sci.*, **16**(1), 1157-1164. <https://doi.org/10.1080/16583655.2022.2151297>

- Mohareb, R.M.; Ibrahim, R.A.; Moustafa, H.E. (2010). Hydrazide-Hydrazones in the synthesis of 1, 3, 4-oxadiazine, 1, 2, 4-triazine and pyrazole derivatives with antitumor activities. *The Open Organic Chem. J.*, **4**(1). <https://doi.org/10.2174/1874095201004010008>
- Mohareb, R.M.; Schatz, J. (2011). Anti-tumor and anti-leishmanial evaluations of 1, 3, 4-oxadiazine, pyran derivatives derived from cross-coupling reactions of β -bromo-6H-1, 3, 4-oxadiazine derivatives. *Bioorg. Med. Chem.*, **19**(8), 2707-2713. <https://doi.org/10.1016/j.bmc.2011.02.051>
- Murtaza, S.; Shamim, S.; Kousar, N.; Tahir, M.N.; Sirajuddin, M.; Rana, U.A. (2016). Synthesis, biological investigation, calf thymus DNA binding and docking studies of the sulfonyl hydrazides and their derivatives. *J. Molec. Struct.*, **1107**, 99-108. <https://doi.org/10.1016/j.molstruc.2015.11.046>
- Nandy, B.C.; Gupta, A.K.; Mittal, A.; Vyas, V. (2014). Carbazole: It's biological activity. *J. Biomed. Pharm. Res.* **3**, 42-48 <https://doi.org/10.1002/chin.199648240>
- Omer, Dh.A.; Al-Daher, A.M. (2019). New tridentate hydrazone metal complexes derived from 2-hydroxy-4-methoxyacetophenone and some acid hydrazides: Synthesis, characterization and antibacterial activity evaluation. *Raf. J. Sci.*, **28**(2), *Special Issue for the Third Scientific Conference of Chemistry*, 100-111. <https://doi.org/10.33899/rjs.2019.159969>
- Pereira, T.M.; Kümmerle, A.E. (2020). Hydrazone-Based small-molecule chemosensory. In *Computat. Biol. Chem.* Intech Open. <https://doi.org/10.5772/intechopen.92144>
- Popiołek, Ł. (2020). 1, Monika Gawrońska-Grzywacz 2, Anna Berecka-Rycerz 3, Kinga Paruch 1, Iwona Piątkowska-Chmiel 2, Dorota Natarska-Chomicka 2, Mariola Herbet 2, Anna Gumieniczek 3, Jarosław Dudka 2, Monika Wujec. New benzenesulphono hydrazide derivatives as potential antitumor agents. *Oncology letters*, **20**(5), 1-1. <https://doi.org/10.3892/ol.2020.12047>
- Rollas, S.; Güniz Küçüküzgel, Ş. (2007). Biological activities of hydrazone derivatives. *Molecules*, **12**(8), 1910-1939. <https://doi.org/10.3390/12081910>
- Sava, G.; Perissin, L.; Lassiani, L.; Zabucchi, G. (1985). Anti-inflammatory action of hydrosoluble dimethyl-triazenes on the carrageenin-induced edema in guinea pigs. *Chem. Biolog. Interact.*, **53**, 37-43. [https://doi.org/10.1016/s0009-2797\(85\)80082-x](https://doi.org/10.1016/s0009-2797(85)80082-x)
- Segretti, N.D.; Serafim, R.A.; Segretti, M.C.; Miyata, M.; Coelho, F.R.; Augusto, O.; Ferreira, E. I. (2016). New antibacterial agents: Hybrid bioisoster derivatives as potential E. coli FabH inhibitors. *Bioorg. Med. Chem. Lett.*, **26**(16), 3988-3993. <https://doi.org/10.1016/j.bmcl.2016.06.089>
- Shaaban, M.M.; Ragab, H.M.; Akaji, K.; McGeary, R.P.; Bekhit, A.E.A.; Hussein, W.M.; Bekhit, A.A. (2020). Design, synthesis, biological evaluation and in silico studies of certain aryl sulfonyl hydrazones conjugated with 1, 3-diaryl pyrazoles as potent metallo- β -lactamase inhibitors. *Bioorg. Chem.*, **105**, 104386. <https://doi.org/10.1016/j.bioorg.2020.104386>
- Shet, L.S.; Shelar, A.R.; Manvi, F.V. (2010). Synthesis of new chiral and nonchiral pyrido [3, 2-e],[1, 3, 4] oxadiazine derivatives. *E-J. Chem.*, **7**(1), 149-156. <https://doi.org/10.1155/2010/791796>
- Shubakara, K.; Umesha, K.B.; Srikantamurthy, N.; Chethan, J. (2014). Antioxidant and DNA damage inhibition activities of 4-Aryl-N-(4-aryl-thiazol-2-yl)-5, 6-dihydro-4H-1, 3, 4-oxadiazine-2-carboxamides. *J. Chem. Sci.*, **126**(6), 1913-1921. <https://doi.org/10.1007/s12039-014-0638-4>
- Siemann, S.; Evanoff, D.P.; Marrone, L.; Clarke, A.J.; Viswanatha, T.; Dmitrienko, G. I. (2002). N-arylsulfonyl hydrazones as inhibitors of IMP-1 metallo- β -lactamase. *Antimicrob. Agents Chemoth.*, **46**(8), 2450-2457. <https://doi.org/10.1128/aac.46.8.2450-2457.2002>
- Verma, G.; Marella, A.; Shaquiquzzaman, M.; Akhtar, M.; Ali, M.R.; Alam, M.M. (2014). A review exploring biological activities of hydrazones. *J. Pharm. Bioall. Sci.*, **6**(2), 69. <https://doi.org/10.4103/0975-7406.129170>

- Xia, Y.; Fan, C.D.; Zhao, B.X.; Zhao, J.; Shin, D.S.; Miao, J.Y. (2008). Synthesis and structure–activity relationships of novel 1-arylmethyl-3-aryl-1H-pyrazole-5-carbohydrazide hydrazone derivatives as potential agents against A549 lung cancer cells. *European J. Med. Chem.*, **43**(11), 2347-2353. <https://doi.org/10.1016/j.ejmech.2008.01.021>
- Xin Su; Ivan Aprahamian (2014). Hydrazone-based switches, metallo-assemblies and sensors. *Chem. Soc. Reviews*, **6**, 1791-1998. <https://doi.org/10.1039/c3cs60385g>
- Yang, K.; Yang, J.Q.; Luo, S.H.; Mei, W.J.; Lin, J.Y.; Zhan, J.Q.; Wang, Z.Y. (2021). Synthesis of N-2 (5H)-furanonyl sulfonyl hydrazone derivatives and their biological evaluation in vitro and in vivo activity against MCF-7 breast cancer cells. *Bioorg. Chem.*, **107**, 104518. <https://doi.org/10.1016/j.bioorg.2020.104518>
- Zhang F.F.; Gan L.L.; Zhou G.H. (2010). Synthesis, anti-bacterial and antifungal activities of some carbazole derivatives. *Bioorg. Med. Chem.*, **20**, 1881-1884. <https://doi.org/10.1016/j.bmcl.2010.01.159>
- Zheng, L.W.; Wu, L.L.; Zhao, B.X.; Dong, W.L. (2009). Miao. *YJ Bioorg. Med. Chem*, **17**, 1957-1962. <https://doi.org/10.1016/j.bmc.2009.01.037>
- Zhou, Z.; Wang, X.; Zhou, T. (2019). Synthesis and antibacterial activity of benzenesulfonylhydrazone derivatives of methyl dehydroabietate. *Russian J. General Chem.*, **89**(4), 819-823. <https://doi.org/10.1134/s1070363219040297>

تحضير وتشخيص بعض معوضات 5,2,1-أوكساديازين المشتقة من H9 كاربازول

سالم جاسم محمد

مها محمد توفيق

قسم الكيمياء/ كلية العلوم/ جامعة الموصل

الملخص

تمتلك الهيدرازونات مجموعة الأزوميثين $\text{CH} = \text{NHN} -$ وتعتبر من مشتقات الألديدات والكيتونات والتي تم فيها استبدال ذرة الأكسجين بـ NNH_2 . في هذه البحث، تم دراسة تحضير بعض مركبات 5,2,1-أوكساديازين الجديدة وتم تحضير المشتقات ابتداءً من الكاربازول كمادة أولية، والذي عند التفاعل مع كلورو كلوريد الاسيتيل في الإيثانول المطلق كمذيب ينتج (2) ethyl 1- (9H-carbazole-9-yl) 2-chloroethanone. هذا المركب (2) عند التفاعل مع الهيدرازين المائي سيعطي (3) (H-carbazol-9-yl) -2-hydrazinylethanone. يتبع ذلك معاملة الهيدرازيد الناتج إما معوضات البنزالديهايد أو معوضات الأسيتوفينون البديل سوف ينتج مشتقات الهيدرازونات (5a-d) (4a-e) على التوالي. ثم إجراء التفاعل لمركبات الهيدرازون هذه مع أنهيدريد حمض الخليك لإعطاء مركبات 1,2,5-أوكساديازين المقابلة (6a-e) and (7a-d) تم التأكد من نقاوة هذه المركبات ومتابعة سير التفاعلات بواسطة كروماتوغرافيا الطبقة الرقيقة (TLC)، وتم تشخيص المركبات المصنعة حديثاً في هذه الدراسة عن طريق التحليل الفيزيائي والطيفي المتضمن الأشعة تحت الحمراء (FT-IR)، الرنين المغناطيسي النووي (^1H NMR and ^{13}C NMR) لبعض هذه المركبات.

الكلمات الدالة: الكاربازول، الهيدرازونات، قواعد شيف، الأوكساديازين، أنهيدريد حامض الخليك.