



## Effects of Topsin M fungicide, nanoparticles with soil biobacteria on growth and their residues on eggplant fruits

Subreen Muhsen Hasan<sup>1</sup>      Salwan Abdul Zahra Jabbar Allobawi<sup>2</sup>  
<sup>1</sup> Directorate of Agriculture in Muthanna Governorate / Ministry of Agriculture  
<sup>2</sup> College of Agriculture / University of Kufa  
E- mail : Sabrynm560@gmail.com

Received on 28/12 /2024 Accepted on / 10/2/2025 Published on /30 /6/2025

### Abstract

The study was conducted in the plastic house at the research station which belongs to the College of Agriculture / University of Kufa during the winter season of 2024. The experimental soil was contaminated with the pathogenic fungus *Sclerotinia sclerotiorum* which loaded on millet seeds before transplanted at a rate of 5 g / 1 kg, and the bio-resistance bacteria *Pseudomonas fluorescens* was added at a rate of 5 ml / 1 kg and mixed well with the soil to ensure its spread. Furthermore, the fungicide was added at the dose which recommended by the producing company in its common and Nano form, at the following treatments : T1( Control treatment) , T2 (Biobacteria), T3 (Nano-pesticide 0.1) , T4 (Nano-pesticide 0.05) , T5 (common pesticide 0.1), T6(common fungicide 0.05), T7( Bacteria + Nano 0.1), T8( Bacteria + Nano 0.05), T9 (Bacteria + common fungicide 0.1) and T10( Bacteria + common pesticide 0.05). Vegetative growth, yield and fungicide residues in fruits were measured. The experiment was carried out using a randomized completely block design (R.C.B.D.) with three replications. Means were compared using the L.S.D test at a probability level of 0.05%. The results revealed that the effect of common and Nano fungicide in reducing the negative effects on vegetative growth and plant yield as following: T7( Bacteria + Nano 0.1) was superior with highest plant height, fresh and dry weight of vegetative growth, roots dry weight, number of fruits and plant yield which reached (19.86 cm, 255.15 g, 103.56 g, 12.63 g, 10.17 fruits plant-1 and 547.05 g plant-1) respectively .

**Keywords:** Age, Cumulus Cells, morphological, , bovine oocytes, IVM **Key words:** Topsin M , eggplant ,Nano pesticide, fungicide .

## Introduction

Gray and white mold *S. sclerotiorum* are the most common types of fungal diseases that infect the eggplant. These fungi cause significant economic damages in generally due to its infection for more than 408 genera and 75 families such as Solanaceae, Crucifaceae, Umbellifaceae, Chenopdiaceae families during the harvest season. [1]. In addition to eggplant, *S. sclerotiorum* also infect a wide range of crops including sunflowers, soybeans, canola, chickpeas, pistachios, as well as lentils and peas [2]. According to (Kamran et al., 2017). The infection rate exceeded 85% depending on the planting time and the environmental conditions. In Iraq, there are no accurate statistics on the losses caused by this disease. It must be combated through chemical pesticides used in this field. However, chemical pesticides are considered one of the most important methods of controlling pests [3]. Despite their side effects on crops, water pollution, living organism, air and soil due to the accumulation of toxic substances within the various nutritional components which in turn reflects its negative impact on animals and humans health [4]. According to [5, 6] a positive relationship were found between fungicides and cancer, respiratory diseases as well as hormonal imbalance within the organism depending on the residual toxic accumulation of these pesticides

Nanotechnology has played an important role in reducing the damage of chemical pesticides by adapting materials at the level of atomic and molecular arrangement, with the aim of bringing about a radical change in current food and agricultural systems. What is expected during the next two decades is to exceed the effects of nanotechnology on agricultural systems and products, with lower concentrations as compare with the traditional materials [7]. The modern trend in pest control is

to increase the effectiveness of low concentrations of chemical pesticides by reducing fungicide particles from common sizes to Nano sizes, thus increasing their penetration speed into the bodies of the targeted pests and increasing their effectiveness as a result of increasing their surface area. Based on the above and the importance of white mold disease on eggplant plants and the increase in its damage in recent years.

The study aimed to measure the residues of the fungicide in its nano and common forms on growth characteristics and yield of eggplant fruits.

## Material and methods

A field experiment was conducted in the plastic house at the research station in College of Agriculture / University of Kufa during the winter season of 2024. A mixed soil was used for cultivation, as the plastic pots were treated with the pathogenic fungus *Sclerotinia sclerotiorum* loaded on millet seeds at a rate of 5 g / 1 kg and mixed well to ensure the spread of the fungal vaccine. The bio-resistance bacteria *Pseudomonas fluorescens* were added at a rate of 5 ml / 1 kg and mixed well with the soil to ensure its spread. Then, the next day, the seeding process was carried out and the fungicide was added at the recommended dose from the producing company in its common and Nano form at a rate of three time periods (10, 20, 30) minutes and at a distance of 4 cm from the source. The plastic pot treatments were as follows: T1( Control treatment), T2( Biobacteria), T3(Nano pesticide dose 0.1), T4( Nano pesticide 0.05), T5( common fungicide dose 0.1), T6(

Classic pesticide 0.05), T7( Bacteria + Nano fungicide e 0.1), T8( Bacteria + Nano pesticide 0.05), T9( Bacteria + common fungicide 0.1) and T10( Bacteria + Regular pesticide 0.05).Vegetative growth, yield and fruit residues were measured. A randomized complete block design with GenStat was used to analyze the results and the means were compared at 0.05 probability level.

## Results and Discussion

Plant height (cm):

From the results shown in Table 1, it is clear that there were significant differences between the different treatments and soil type on plant

height characteristic. However, T10 ,T7, T9 were superior in giving the highest shoots plant height reached (19.893, 19.868 and 19.443 cm) and no significant differences were found among them . Likewise, it was found that contaminated soil affected significantly on shoots plant height with average (19.665 cm) for control soil(P2) comparing with 14.465 cm for P1 (contaminated soil) , also noted that there were significant differences between Contamination with pathogenic fungi treatments. The P2 treatment (19.665 cm) was significantly superior compare than P1 (14.465 cm. For the interactions between the treatments and type of soil, the results in Table 1 showed that (T7P2) treatment was significantly superior to give (23.56 cm) shoots height comparing with (11.13 cm)in (T1P1) treatment.

Table (1): Effect of different treatments , soil types and their interactions on eggplant shoots height (cm)

Treatments	Contaminated soil		Mean of treatments
	Control (P2)	Contaminated (P1)	
T1 (Control treatment)	16.1	11.13	13.615
T2 (Biobacteria)	17.43	12.9	15.165
T3 ( Nano fungicide dose 0.1)	19.667	15.31	17.488
T4 (Nano fungicide (0.05)	18.16	12.86	15.51
T5 (common fungicide e dose (0.1)	19.63	14.76	17.195
T6 (common fungicide (0.05)	17.98	12.4	15.19
T7 (Bacteria + Nano (0.1)	23.56	16.177	19.868
T8 (Bacteria + Nano (0.05)	18.63	15.93	17.28
T9 (Bacteria + common fungicide (0.1)	23.027	15.86	19.443
T10 (Bacteria + common fungicide (0.05)	22.467	17.32	19.893
Mean of Contaminated soil	19.665	14.465	
<b>L.S.D. 0.05</b>	<b>P=0.2534</b>	<b>TP=0.8014</b>	<b>T= 0.5667</b>

### Fresh weight of vegetative growth parameter(g):

For the vegetative fresh weight parameter, data in table (2), revealed that T7 and T10 treatments affected significantly with giving highest rate (255.15, 251.16 g) respectively , whereas less average was found at T1 (66.38g).Concerning to the contaminated soil, P2 treatment was significantly superior with (234.69 g) comparing with P1 (124.94 g ). The interaction between the different treatments and contaminated soil showed that, P2T7 treatment recorded (339.11 g), while the lowest value (66.38g) was recorded in the P1T1 treatment.

Table (2): Effect of different treatments, soil types and their interactions and their interactions on eggplant shoots vegetative fresh weight (g)

Contaminated soil	Contaminated soil		Mean of treatments
	Control (P2)	Contaminated (P1)	
<b>T1 (Control treatment)</b>	153.01	66.38	109.7
<b>T2 (Biobacteria)</b>	184.23	80.16	132.2
T3 ( Nano fungicide dose 0.1)	242	155.03	198.52
T4 (Nano fungicide (0.05))	203.18	83.92	143.55
T5 (common fungicide e dose (0.1))	218.5	92.86	155.68
T6 (common fungicide (0.05))	166.51	70.3	118.41
T7 (Bacteria + Nano (0.1))	339.11	171.19	255.15
T8 (Bacteria + Nano (0.05))	222.6	169.5	196.05
T9 (Bacteria + Common fungicide (0.1))	316.89	158.7	237.8
T10 (Bacteria + Common fungicide (0.05))	300.91	201.4	251.16
Mean of Contaminated soil	234.69	124.94	
<b>L.S.D. 0.05</b>	<b>P= 1.099</b>	<b>TP= 3.474</b>	<b>T= 2.457</b>

**Dry weight of vegetative growth parameter(g):**

The results from Table 3 showed that there were significant differences among the treatments and T7 giving the highest rate (103.56 g) comparing with lowest rate (51.59g) for T1 treatment. Further, it also noted that control (P2) treatment was significantly superior with (92.74 g) dry weight compare

than P1 (55.62 g). For the interactions between treatments and contaminated soil, data in table 3 showed that, highest value was recorded in the T7P2 treatment, reached to (135.19 g), while the lowest value (32.33) was recorded in the T1 P1 treatment.

Table (3): Effect of different treatments , *S. sclerotiorum* fungus and their interactions on eggplant shoots vegetative dry weight (g)

Contaminated soil	Contaminated soil		Mean of treatments
	Control (P2)	Contaminated (P1)	
<b>T1 (Control treatment)</b>	70.86	32.33	51.59
<b>T2 (Biobacteria)</b>	73.1	45	59.05
T3 ( Nano fungicide dose 0.1)	108.6	59	83.8
T4 (Nano fungicide (0.05))	83.6	46.16	64.88
T5 (common fungicide e dose (0.1))	84.53	52.1	68.31
T6 (common fungicide (0.05))	36.4	34.68	35.54
T7 (Bacteria + Nano (0.1))	135.19	71.93	103.56
T8 (Bacteria + Nano (0.05))	99.98	67.8	83.89
T9 (Bacteria + Common fungicide (0.1))	124.58	60.94	92.76
T10 (Bacteria + Common fungicide (0.05))	110.55	86.26	98.4
Mean of Contaminated soil	92.74	55.62	
<b>L.S.D. 0.05</b>	<b>P=1.082</b>	<b>TP=3.421</b>	<b>T= 2.419</b>

### Roots fresh and dry weight (g) :

The results in table 4 and 5 clarified that there were significant differences among the treatments under the study in terms of roots fresh and dry weight parameters. However, (T3, T4, T5, T6, T7 and T8) affected significantly on roots fresh weight with average of (23.68, 31.32, 22.09, 27.81, 23.49 and 25.41 g) respectively. while T7 was superior in giving highest average of dry weight (12.632 g). Concerning to contaminated soil,

control soil (P2) produced highest fresh and dry weight (27.34 and 9.941 g) as compare to (17.81 and 7.495 g) in contaminated soil (P1) respectively. For the interaction between the treatments and contaminated soil, data in table 4 showed that most of interaction the T7P2(35.7g) treatments affected significantly on eggplant roots fresh and significant differences were found among these treatments for roots fresh weight. On the contrary, the highest rate for dry weight was recorded (13.433 g) at interactions T7P2 (table 5).

Table (4): Effect of different treatments , *S. sclerotiorum* fungus and their interactions on eggplant roots fresh weight (g)

Contaminated soil	Contaminated soil		Mean of treatments
	Control (P2)	Contaminated (P1)	
<b>T1 (Control treatment)</b>	20.27	10.07	15.17
<b>T2 (Biobacteria)</b>	23.63	12.3	17.96
T3 ( Nano fungicide dose 0.1)	30.82	16.53	23.68
T4 (Nano fungicide (0.05))	26.95	35.7	31.32
T5 (common fungicide e dose (0.1))	26.28	17.9	22.09
T6 (common fungicide (0.05))	22.59	12.37	17.48
T7 (Bacteria + Nano (0.1))	35.7	19.93	27.81
T8 (Bacteria + Nano (0.05))	29.45	17.53	23.49
T9 (Bacteria + Common fungicide (0.1))	32.18	18.63	25.41
T10 (Bacteria + Common fungicide (0.05))	25.58	17.1	21.34
Mean of Contaminated soil	27.34	17.81	
<b>L.S.D. 0.05</b>	<b>P=3.699</b>	<b>TP=11.696</b>	<b>T=8.270</b>

Table (5): Effect of different treatments , *S. sclerotiorum* fungus and their interactions on eggplant roots dry weight (g)

Contaminated soil	Contaminated soil		Mean of treatments
	Control (P2)	Contaminated (P1)	
<b>T1 (Control treatment)</b>	7.63	4.307	5.968
<b>T2 (Biobacteria)</b>	8.44	5.507	6.973
T3 ( Nano fungicide dose 0.1)	11.97	10.103	11.037
T4 (Nano fungicide (0.05))	9.16	6.797	7.978
T5 (common fungicide e dose (0.1))	9.043	6.843	7.943
T6 (common fungicide (0.05))	8.103	5.103	6.603
T7 (Bacteria + Nano (0.1))	13.433	11.83	12.632
T8 (Bacteria + Nano (0.05))	10.67	7.51	9.09
T9 (Bacteria + Common fungicide (0.1))	12.53	11.003	11.767
T10 (Bacteria + Common fungicide (0.05))	8.433	5.943	7.188

Mean of Contaminated soil	9.941	7.495	
<b>L.S.D. 0.05</b>	<b>P=0.1960</b>	<b>TP=0.6198</b>	<b>T= 0.4382</b>

### Fruit numbers :

For number of eggplant fruits, the results in Table 6 showed that there were significant differences among the treatments in. However, T7 and T9 achieved ,highest number of fruits (10.17 and 10) respectively and no significant differences was found between both treatments. For soil type, contaminated soil (P1) was

affected positively on number of fruits with average (7.40) comparing with (5.0) for control soil (P2). Furthermore, the interaction between the treatments and soil type revealed that T7P1 was superior with highest number of fruits 13.67 while T1P2 treatment recorded least number 4,00 of eggplant fruits .

Table (6): Effect of different treatments, *S. sclerotiorum* fungus and their interactions on the number of eggplant fruits

Contaminated soil	Contaminated soil		Mean of treatments
	Control (P2)	Contaminated (P1)	
<b>T1 (Control treatment)</b>	3	3.33	3.17
<b>T2 (Biobacteria)</b>	3.67	4.67	4.17
T3 ( Nano fungicide dose 0.1)	6	6.67	6.33
T4 (Nano fungicide (0.05)	4.67	9	6.83
T5 (common fungicide e dose (0.1)	4	6	5
T6 (common fungicide (0.05)	4.67	5	4.83
T7 (Bacteria + Nano (0.1)	6.67	13.67	10.17
T8 (Bacteria + Nano (0.05)	5	6.67	5.83
T9 (Bacteria + Common fungicide (0.1)	7.67	12.33	10
T10 (Bacteria + Common fungicide (0.05)	4.67	6.67	5.67
Mean of Contaminated soil	5.00	7.40	
<b>L.S.D. 0.05</b>	<b>P= 0.412</b>	<b>TP=1.302</b>	<b>T= 0.920</b>

### The yield of eggplant fruits :

The results inTable (7) showed that there are significant differences among different treatments in related to the yield of eggplant fruits. However, T7 produced highest rate (547.05 g ) and significantly superior as compare to the other treatments. It also noted that there were significant differences between

Contaminated and non contaminated soil whereas P2 was superior with (390.77 g) compare with (249.64 g ) for P1 treatment. For the interaction between different treatments and soil type, T7 P2 recorded highest value reached (683. 7 g), while the lowest value (71.0 g) was recorded in the T1 P1 treatment.

**Table (7): Effect of different treatments , *S. sclerotiorum* fungus and their interactions on eggplant yield**

Contaminated soil	Contaminated soil		Mean of treatments
	Control (P2)	Contaminated (P1)	
T1 (Control treatment)	175.8	71	123.4
T2 (Biobacteria)	300.5	185.9	243.2
T3 ( Nano fungicide dose 0.1)	493.6	311.4	402.5
T4 (Nano fungicide (0.05))	350.3	188.5	269.4
T5 (common fungicide e dose (0.1))	375.4	290.5	332.95
T6 (common fungicide (0.05))	275.2	142.61	208.9
T7 (Bacteria + Nano (0.1))	683.7	410.4	547.05
T8 (Bacteria + Nano (0.05))	406.6	307.9	357.25
T9 (Bacteria + Common fungicide (0.1))	540	400.12	470.06
T10 (Bacteria + Common fungicide (0.05))	306.6	188.05	247.32
Mean of Contaminated soil	390.77	249.64	
L.S.D. 0.05	P =3.048	TP =6.816	T=9.64

### Fungicide residues:

Results in table 8 showed that there were significant effects between the fungicide doses and time of infection characteristic. However, Nano fungicide at the dose 0.1 produced highest rate (372.77 mg Kg<sup>-1</sup>), which significantly superior to the rest of treatments. Table 8 also revealed that there were significant differences between different times and 1 hour

Table (8): fungicide residues found in eggplant fruits infected with the pathogenic fungus *S. sclerotiorum*.

gave (436.05 mg Kg<sup>-1</sup>) which significantly superior to the rest. For the interaction, data showed that treated with nano fungicide at 0.1 concentrations for 1 hour was significantly superior with (710.8 mg Kg<sup>-1</sup>) while no residues were found for common fungicide at 0.1 or 0.05 when interaction 5 day or 0.05 with 3 day.

Fungicide	Dose	Infection time			Mean of fungicide
		5 day	3 day	One hour	
Common fungicide	0.1	0.0	88.9	524.9	204.6
	0.05	0.0	0.0	123.6	40.87
Nano fungicide	0.1	85.9	321.6	710.8	372.77
	0.05	0.0	122.6	385.9	169.50
Mean of time		21.48	133.28	436.05	
L.S.D. 0.05	Time=1.974		fungicide=2.279	Interaction =3.947	

### Discussion :

From the results above, it is clear that, the decrease in vegetative growth resulting from treatment with common and Nano pesticides alone may be due to the possibility that many chemical pesticides kill the organism by interfering with the process of cellular

respiration and energy production in the mitochondria or by changing the bio-encapsulation of some compounds which is necessary for the organism's life, such as nucleic acids (RNA, DNA) as well as proteins, and as a result, they affect on the glycolysis process

which occurs in the cytoplasm and the Krebs cycle and oxidative phosphorylation in the mitochondria [8 ,9]. The fungi that were severely affected by pesticides may be due to the inhibition process, in addition to the fact that some pesticides inhibit cutinase and Phosphatase enzyme, or affect plant growth by inhibiting (DNA) synthesis, cell division or by inhibiting important enzymes in the mitochondria [10].

As a result of what was mentioned above in this research, it is believed that the treatment is superior due to the effect of the biobacteria in reducing the effect of the pathogenic fungus that causes white mold disease on eggplant (source), and thus allowed the plant to grow in a more normal way, which increased the plant hormones responsible for growth, including auxins, which in turn increased the plant height fresh and dry weight of vegetative growth and root system. [11 , 12 , 13] , that's led to increase the number

#### **Conclusion:**

Through research, it has been proven that it is necessary to add biobacteria to reduce the residual effect of the pesticide, thus reducing the damage to the plant and thus increasing the yield and reducing the residues .

#### **References**

- 1- **Suleman, p ; Abdullah, m. t. & ali, n. y. (2008)** . Biological control of *Sclerotinia sclerotiorum* (Lib.) de Bary with *Trichoderma harzianum* and *Bacillus amyloliquefaciens*, *Crop Protection*, 27: 1354-1359.
- 2- **Bolton, M. D. ; Thomma, B. P. H. J. & Nelson, B. D. (2006)**. *Sclerotinia sclerotiorum* (Lib.) de Bary: biology and molecular traits of a cosmopolitan pathogen, *MOLECULAR PLANT PATHOLOGY*, 7(1): 1 – 16.
- 3- **Al-Mallah, N. M. (2015)**. Chemical and biological pesticides for germs and plant pathogens, Ibn Al-Atheer Printing and Publishing House, Mosul, Iraq.Pp 328.
- 4- **Al-Homsi, . and N. Zidane. (2016)**. Modern techniques in preparing and purifying pesticides. Dar Alkotb Alilmiyah.Pp 304 .
- 5- **Piel, C., Pouchieu, C., Carles, C., Beziat, B., Boulanger, M., Bureau, M., ... & Baldi, I. (2019)**. Agricultural



exposures to carbamate herbicides and fungicides and central nervous system tumour incidence in the cohort AGRICAN. *Environment international*, 130, 104876.

- 6- **Juntarawijit, C.(2018).** Association between diabetes and pesticides: A case-control study among Thai farmers. *Environ. Health Prev. Med.* . 23, 1–10.
- 7- **Al-Rubaie, Hussein Fadhel and Abdul Jassim Mohsen Al-Jubouri. (2014).** Use of nanotechnology in agriculture and food. Ministry of Science and Technology, P.O. Box 765, Baghdad - University of Nahrain - Biotechnology Research Center. Baghdad.
- 8- **Araoud, M., Douki, W., Rhim, A., Najjar, M. F., & Gazzah, N. (2007).** Multiresidue analysis of pesticides in fruits and vegetables by gas chromatography-mass spectrometry. *Journal of Environmental Science and Health Part B*, 42(2), 179-187. *armaceutical biology*, 55(1), 1047-1053.
- 9- **Azpiazu, C., Bosch, J., Bortolotti, L., Medrzycki, P., Teper, D., Molowny-Horas, R., & Sgolastra, F. (2021).** Toxicity of the insecticide sulfoxaflor alone and in combination with the fungicide fluxapyroxad in three bee species. *Scientific reports*, 11(1), 6821.
- 10- **Al-Adel, K. M. (2006).** Pesticides. Directorate of Dar Al-Kutub for Printing and Publishing. University of Baghdad. Ministry of Higher Education and Scientific Research. Republic of Iraq.
- 11- **Issa, F.H.; M .S.Omar and F.H.Al-Sahaf (2001).** Effect of pesticides treatments on decay and sprouting percentage of two microtubers size of two potato cultivars. *The Iraqi Journal of Agricultural*.32(3):59-64.
- 12- **Issa, F.H.S.O.Sadik and T.A. Zahwan. ( 2008).** The effect of growth regulator treatments on Minitubers to produce Tuberlet from potato *Solanum tuberosum* *Journal Of Tikrit University for Agricultural Science*.9(1):96-103.
- 13- **Kazem, A.T.;F.H.Issa and A. Abdulla .(2021).** Effect of Nano NPK Fertilizer on Growth and Early Yield of Eggplant . *IOP Conf. Series: Earth and Environmental Science* . Pp 4.

- 14- Raheem,A. and Issa ,F.H.(2018).**  
Evaluating productivity of an eggplant as influenced by Mycorrhiza fungi and other selected fertilizing materials. Merit Research Journal of Agricultural Science and Soil Sciences. 6(9) pp. 118-122.
- 15- Abd El-Ghany, T.M., & Masmali, I.A. (2016).** Fungal Biodegradation of Organophosphorus Insecticides and their Impact on Soil Microbial Population. Journal of Plant Pathology & Microbiology. 7(5) ; 2157-7471.
- 16- Colombo, J.C., Cabello, M. and Arambarri, A. (1996).**  
Biodegradation of aliphatic and hydrocarbons by natural soil microflora and pure cultures of imperfect and lignolytic fungi. Env. Poll., 94: 355-362.1., 61: 289-298.