

Advanced Techniques in Natural Transform and Homotopy Analysis for Fractional Differential Equations

Naser Rhaif Seewn¹,  and Mohammed Taimah Yasser^{2,a},  Mountassir Hamdi Cherif³

³Hight School of Electrical and Energetics Engineering of Oran (ESGEE-Oran),

Oran, Algeria; mountassir27@yahoo.fr

^{1,2} Department of Mathematics Faculty of Education for Pure Sciences Thi-Qar University

* Corresponding email: Naserrhaif809@utq.edu.iq

^a email: edpma4m23@utq.edu.iq

Received 13 / 1 /2025, Accepted 18/ 03 /2025, Published 1/06 /2025



This work is licensed under a [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/).

Abstract:

This study employs the natural transformation approach combined with the homotopy perturbation technique, a new and effective hybrid method that integrates the homotopy approach with the natural transformation, using a simplified iterative procedure that reduces computational demands. This method provides fast, convergent, and sequential solutions. The reliability of the method is verified through its application to two case studies of ST-TE within the context of the Caputo derivative, which involves the definition of non-singular kernel functions. The study also includes comprehensive comparisons between approximate and exact solutions, utilizing relevant scientific literature to assess the accuracy and effectiveness of the method. Graphical representations highlight the impact of incorrect temporal and spatial parameters on the solution's behavior. The findings suggest that this method is easy to implement and well-suited for studying complex physical models governed by nonlinear partial differential equations with fractional time components.

Keywords: Fractional Differential Equations; Natural Transform; Atangana-Baleanu Fractional Derivative; Homotopy Method

1-Introduction

Linear and nonlinear fractional differential equations are crucial tools for modeling complex phenomena in various fields, such as fluid dynamics, acoustics, electromagnetism, signal processing, analytical chemistry, biology, and other areas of physical sciences and engineering. In recent years, numerous analytical and computational methods have been developed to solve these equations, including the Adomian Decomposition Method (ADM), the Homotopy Analysis Method (HAM), the Reduced Transform Method (RTM), and the Iterative Differential Transform Method (IDTM), among others [2][3]. These methods offer powerful tools for obtaining approximate solutions to fractional-order models that are encountered in diverse scientific and engineering problems.

Fractional differential equations involve derivatives of non-integer order, providing a more general and flexible framework compared to traditional integer-order differential equations. This flexibility allows for more accurate modeling of real-world phenomena that exhibit memory and hereditary properties, such as viscoelastic materials, anomalous diffusion, and systems with long-range dependence. However, solving these equations analytically remains a challenging task, particularly when they include nonlinear terms or complex boundary conditions.

Given the complexity of these problems, there has been significant interest in the development of efficient and robust solution techniques. Among these, the Natural Transform Homotopy Method (NTHM) has gained attention for its ability to effectively address both linear and nonlinear fractional differential equations. This research aims to explore and develop advanced solution techniques for fractional differential equations using the Natural Transform Homotopy method. The paper will discuss the theoretical foundations, implementation strategies, and real-world applications of this method across various disciplines.

In addition to NTHM, other notable methods for solving fractional differential equations include:

- **The Reduced Transform Method (RTM):** This method has shown considerable success in solving both linear and nonlinear fractional differential equations, particularly in engineering applications.
- **The Iterative Differential Transform Method (IDTM):** An effective numerical method that iteratively solves fractional differential equations, proving useful in obtaining accurate approximate solutions for complex problems.
- **The Laplace Transform Method:** Often used for linear fractional differential equations, providing a solid analytical approach for solving certain types of fractional models.

These methods, along with the Natural Transform Homotopy Method, represent a growing toolkit of analytical and numerical techniques for solving fractional differential equations.

This paper also aims to examine the strengths and limitations of these methods and to determine the most effective approach for obtaining numerical solutions. Through a thorough theoretical and numerical analysis, we aim to highlight the best methods for achieving accurate and efficient solutions to fractional differential equations.

2- Natural Transform

This section introduces the N-T transform, covering its basic definition and properties. This transform has been used in various applications for solving differential equations, especially in the context of fractional calculus and other advanced mathematical problems. the **N-T**, which was later renamed the Natural Transform by Belgacem and Silambarasan [4].

A list of special Natural Transforms for common functions is provided below:

$$\begin{aligned} 1 &\rightarrow \frac{1}{s} \\ t &\rightarrow \frac{u}{s^2} \\ e^{at} &\rightarrow \frac{1}{s - au} \\ \frac{t^{n-1}}{(n-1)!}, n = 1, 2, \dots &\rightarrow \frac{u^{n-1}}{s^n} \\ \sin(t) &\rightarrow \frac{u}{s^2 + u^2} \end{aligned}$$

Definition 1[5] The N-T of the function $\mathcal{B}(t) > 0$ and $\mathcal{B}(t) = 0$ for $t < 0$ is specified for the collection of functions:

$$\beta = \left\{ \mathcal{B}(t): \exists M, \tau_1, \tau_2 > 0, |\mathcal{B}(t)| < Me^{\frac{|t|}{\tau_j}} \text{ if } t \in (-1)^j [0, \infty), j = 1, 2, \dots \right\}$$

By the following integral:

$$N^{-1}[\mathcal{L}(t)] = V(s, \mathfrak{U}) = \frac{1}{\mathfrak{U}} \int_0^\infty e^{-\frac{st}{\mathfrak{U}}} dt, s > 0, \mathfrak{U} > 0$$

Definition 2[1] The inverse N-T of a function is defined as:

$$N^{-1}[V(s, \mathfrak{U})] = \mathcal{L}(t) = \frac{1}{2i\pi} \int_{p-\infty}^{p+\infty} e^{\frac{st}{\mathfrak{U}}} V(s, \mathfrak{U}) ds$$

Where s and $\mathfrak{U}$ are N-T variables and $\mathfrak{S}$ is real constant.

Definition 3. [30], Suppose that the function $\varphi \in H^1(a, b)$, $a > b$, The Atangana-Baleanu Fractional Derivative (ABFD) of φ is:

$${}^{AB}D_t^\gamma \varphi(t) = \frac{M(\gamma)}{1-\gamma} \int_a^t E_\gamma \left(\frac{-\gamma(t-\xi)^\gamma}{\gamma-1} \right) \varphi'(\xi) d\xi \quad t \geq 0$$

where $0 < \gamma < 1$ and $M(\gamma)$ is a normalization function, such that $M(0) = M(1) = 1$. and $\varphi'(\xi)$ is the derivative of φ .

In view of (2.9), the ABFD of a constant and Natural transform of this derivative defined as follows:

, where c is a constant.:

$$1 - {}^{AB}D_t^\gamma c = 0$$

$$.2 - L\{{}^{AB}D_t^\gamma \varphi(\xi, t)\} = \frac{s^\gamma L\varphi(\xi, t)}{s^\gamma(1-\gamma)+\gamma} - \frac{s^{\gamma-1} L\{\varphi(\xi, 0)\}}{s^\gamma(1-\gamma)+\gamma}$$

$$3 - {}^{AB}D_t^\gamma \mathcal{T}^k = \frac{M(\gamma)}{1-\gamma} \cdot \Gamma(k+1) \mathcal{T}^k \cdot E_{\gamma, k+1} \left(\frac{-\gamma x^\gamma}{1-\gamma} \right).$$

3- Analysis of the proposed Method

Take into account the fractional partial differential equation involving the Atangana-Baleanu-Caputo operator in the subsequent form:

$${}^{ABC}D_t^\mathfrak{S} \mathfrak{U}(\mathfrak{x}, t) + \zeta[\mathfrak{U}(\mathfrak{x}, t)] + \mu[\mathfrak{U}(\mathfrak{x}, t)] = g(\mathfrak{x}, t) \quad (1)$$

With the starting condition $\mathfrak{U}(\mathfrak{x}, 0) = \mathfrak{U}_0(\mathfrak{x})$, where ${}^{ABC}D_t^\mathfrak{S}$ is the Atangana-Baleanu-Caputo operator, ζ is a linear operator, μ is a nonlinear operator, and g represents the source term. By applying the Natural Transform, we obtain initial condition.

$$\frac{B(\mathfrak{S})}{1-\mathfrak{S}+\mathfrak{S}\left(\frac{\mathcal{L}}{\delta}\right)^\mathfrak{S}} \left(N(\mathcal{L}, \mathfrak{S}) - \frac{1}{\mathfrak{S}} \mathfrak{U}(0) \right) = N(g(\mathfrak{x}, t) - \zeta[\mathfrak{U}(\mathfrak{x}, t)] - \mu[\mathfrak{U}(\mathfrak{x}, t)]) \quad (2)$$

By substituting initial condition in (1)

$$\bar{\mathfrak{U}} = \frac{1}{\mathfrak{S}} \mathfrak{U}_0(\mathfrak{x}) - \frac{1-\delta+\delta\left(\frac{\mathcal{L}}{\mathfrak{S}}\right)^\delta}{B(\delta)} N(\zeta[\mathfrak{U}] + \mu[\mathfrak{U}] - g). \quad (3)$$

By applying $(N^{-1}T)$ to both sides of equation (3), the result is:

$$\mathfrak{U} = \mathfrak{U}_0(\mathfrak{x}) + N^{-1} \left(\frac{1 - \delta + \delta \left(\frac{\mathfrak{b}}{\mathfrak{S}} \right)}{B(\mathfrak{S})} N(\mathfrak{g}) \right) - N^{-1} \left(\frac{1 - \delta + \delta \left(\frac{\mathfrak{b}}{\mathfrak{S}} \right)^\delta}{B(\delta)} N(\zeta[\mathfrak{U}] + \mu[\mathfrak{U}]) \right). \quad (4)$$

Applying homotopy Permutation method,

$$\mathfrak{U}(\mathfrak{x}, \mathfrak{t}) = \sum_{n=0}^{\infty} \rho^n (\mathfrak{x}, \mathfrak{t}), N[\mathfrak{b}(\mathfrak{x}, \mathfrak{t})] = \sum_{n=0}^{\infty} \rho^n H_n(\mathfrak{U}) \quad (5)$$

Where

$$H_n(\mathfrak{U}_1, \mathfrak{U}_2, \mathfrak{U}_3, \dots, \mathfrak{U}_n) = \frac{1}{n!} \frac{\partial^n}{\partial \rho^n} \left[N \left(\sum_{i=0}^{\infty} \rho^i \mathfrak{U}_i(\mathfrak{x}, \mathfrak{t}) \right) \right] \rho = 0, n = 0, 1, 2, \dots \quad (6)$$

By replacing equation (5) into equation (4), yields the following result:

$$\sum_{n=0}^{\infty} \rho^n \mathfrak{U}_n(\mathfrak{x}, \mathfrak{t}) = \vartheta(\mathfrak{x}, \mathfrak{t}) - \rho \left(N^{-1} \left(\frac{1 - \delta + \delta \left(\frac{\mathfrak{b}}{\mathfrak{S}} \right)^\delta}{B(\delta)} N(\sum_{n=0}^{\infty} \rho^n \zeta[\mathfrak{U}_n] + \sum_{n=0}^{\infty} \rho^n H_n(\mathfrak{U})) \right) \right) \quad (7)$$

By equating both sides of the equation, the following conclusion is obtained:

$$\rho^0: \mathfrak{U}_0(\mathfrak{x}, \mathfrak{t}) = \vartheta(\mathfrak{x}, \mathfrak{t}),$$

$$\rho^1: \mathfrak{U}_1(\mathfrak{x}, \mathfrak{t}) = -N^{-1} \left(\frac{1 - \delta + \delta \left(\frac{\mathfrak{b}}{\mathfrak{S}} \right)^\delta}{B(\delta)} N(\zeta[\mathfrak{U}_0] + H_0[\mathfrak{U}]) \right)$$

$$\rho^n: \mathfrak{U}_n(\mathfrak{x}, \mathfrak{t}) = -N^{-1} \left(\frac{1 - \delta + \delta \left(\frac{\mathfrak{b}}{\mathfrak{S}} \right)^\delta}{B(\delta)} N(\zeta[\mathfrak{U}_{n-1}] + H_{n-1}[\mathfrak{U}]) \right)$$

Utilizing the parameter ρ , we express the solution in the subsequent form:

$$\mathfrak{U}(\mathfrak{x}, \mathfrak{t}) = \sum_{n=0}^{\infty} \rho^n \mathfrak{U}_n(\mathfrak{x}, \mathfrak{t}). \quad (8)$$

Setting $\rho = 1$ results in the solving of Eq.(8)

$$\mathfrak{U}(\mathfrak{x}, \mathfrak{t}) = \lim_{\rho \rightarrow 1} \sum_{n=0}^{\infty} \rho^n \mathfrak{U}_n(\mathfrak{x}, \mathfrak{t}) = \sum_{n=0}^{\infty} \mathfrak{U}_n(\mathfrak{x}, \mathfrak{t})$$

4. Illustrative Examples

Example 1 Consider the following fractional diffusion equation expressed as:

$${}^{AB}D_t^\alpha \mathfrak{b} + \mathfrak{b}_{\mathfrak{x}\mathfrak{x}} + \mathfrak{b}_{\mathfrak{y}\mathfrak{y}} + \mathfrak{b}_{\mathfrak{z}\mathfrak{z}} = 0, \quad -\infty < \mathfrak{x}, \mathfrak{y}, \mathfrak{z} < \infty, \mathfrak{t} > 0. \quad (11)$$

$$\mathfrak{b}(\mathfrak{x}, \mathfrak{y}, \mathfrak{z}, 0) = e^{\mathfrak{x}+\mathfrak{y}+\mathfrak{z}}, \alpha \in (0, 1). \quad (12)$$

By applying the N- T to equation (11) and incorporating the specified initial condition, we obtain:

$$\mathfrak{b}(\mathfrak{x}, \mathfrak{y}, \mathfrak{z}, \mathfrak{U}) = \frac{e^{\mathfrak{x}+\mathfrak{y}+\mathfrak{z}}}{\mathfrak{s}} + \frac{\mathfrak{U}^\alpha}{\mathfrak{s}^\alpha} N^+ (\mathfrak{b}_{\mathfrak{x}\mathfrak{x}} + \mathfrak{b}_{\mathfrak{y}\mathfrak{y}} + \mathfrak{b}_{\mathfrak{z}\mathfrak{z}}). \quad (13)$$

Seizing the inverse N-T of equation (13), we obtain:

$$\mathcal{b}(\mathfrak{x}, \mathfrak{y}, Z, \mathfrak{t}) = e^{\mathfrak{x}+\mathfrak{y}+Z} + N^{-1} \left(\frac{\mathfrak{U}^\alpha}{S^\alpha} N^+ [\mathcal{b}_{\mathfrak{x}\mathfrak{x}} + \mathcal{b}_{\mathfrak{y}\mathfrak{y}} + \mathcal{b}_{ZZ}] \right) \quad (14)$$

At this point, we employ the (HPM) to address the issue:

$$\mathcal{b}(\mathfrak{x}, \mathfrak{y}, Z, \mathfrak{t}) = \sum_{n=0}^{\infty} \rho^n \mathcal{b}_n(\mathfrak{x}, \mathfrak{y}, Z, \mathfrak{t}). \quad (15)$$

Subsequently, equation (15) will transform into:

$$\sum_{n=0}^{\infty} \rho^n \mathcal{b}_n(\mathfrak{x}, \mathfrak{y}, Z, \mathfrak{t}) = e^{\mathfrak{x}+\mathfrak{y}+Z} - \rho \left(N^{-1} \left[\frac{\mathfrak{U}^\alpha}{S^\alpha} \cdot N^+ [\sum_{n=0}^{\infty} \rho^n \mathcal{b}_{n\mathfrak{x}\mathfrak{x}} + \sum_{n=0}^{\infty} \rho^n \mathcal{b}_{n\mathfrak{y}\mathfrak{y}} + \sum_{n=0}^{\infty} \rho^n \mathcal{b}_{nZZ}] \right] \right) \quad (16)$$

By equating the factors corresponding to similar powers of ρ in equation (16), we derive the following approximations:

$$\rho^0: \mathcal{b}_0(\mathfrak{x}, \mathfrak{y}, Z, \mathfrak{t}) = e^{\mathfrak{x}+\mathfrak{y}+Z},$$

$$\begin{aligned} \rho^1: \mathcal{b}_1(\mathfrak{x}, \mathfrak{y}, Z, \mathfrak{t}) &= -N^{-1} \left[\frac{\mathfrak{U}^\alpha}{S^\alpha} N^+ [\mathcal{b}_{0\mathfrak{x}\mathfrak{x}} + \mathcal{b}_{0\mathfrak{y}\mathfrak{y}} + \mathcal{b}_{0ZZ}] \right] \\ &= -\frac{e^{\mathfrak{x}+\mathfrak{y}+Z} \mathfrak{t}^\alpha}{\Gamma(\alpha + 1)} \end{aligned}$$

$$\rho^2: \mathcal{b}_2(\mathfrak{x}, \mathfrak{y}, Z, \mathfrak{t})$$

$$\begin{aligned} \rho^2: \mathcal{b}_2(\mathfrak{x}, \mathfrak{y}, Z, \mathfrak{t}) &= -N^{-1} \left[\frac{\mathfrak{U}^\alpha}{S^\alpha} N^+ [\mathcal{b}_{1\mathfrak{x}\mathfrak{x}} + \mathcal{b}_{1\mathfrak{y}\mathfrak{y}} + \mathcal{b}_{1ZZ}] \right] \\ &= -\frac{e^{\mathfrak{x}+\mathfrak{y}+Z} \mathfrak{t}^{2\alpha}}{\Gamma(2\alpha + 1)} \end{aligned}$$

$$\rho^3: \mathcal{b}_3(\mathfrak{x}, \mathfrak{y}, Z, \mathfrak{t})$$

$$\begin{aligned} \rho^3: \mathcal{b}_3(\mathfrak{x}, \mathfrak{y}, Z, \mathfrak{t}) &= -N^{-1} \left[\frac{\mathfrak{U}^\alpha}{S^\alpha} N^+ [\mathcal{b}_{2\mathfrak{x}\mathfrak{x}} + \mathcal{b}_{2\mathfrak{y}\mathfrak{y}} + \mathcal{b}_{2ZZ}] \right] \\ &= -\frac{e^{\mathfrak{x}+\mathfrak{y}+Z} \mathfrak{t}^{3\alpha}}{\Gamma(3\alpha + 1)} \end{aligned}$$

Then, the series solution for equations (11) and (12) is expressed as:

$$\begin{aligned} \mathcal{b}(\mathfrak{x}, \mathfrak{y}, Z, \mathfrak{t}) &= \lim_{N \rightarrow \infty} \sum_{n=0}^N \mathcal{b}_n(\mathfrak{x}, \mathfrak{y}, Z, \mathfrak{t}) = \mathcal{b}_0(\mathfrak{x}, \mathfrak{y}, Z, \mathfrak{t}) + \mathcal{b}_1(\mathfrak{x}, \mathfrak{y}, Z, \mathfrak{t}) + \mathcal{b}_2(\mathfrak{x}, \mathfrak{y}, Z, \mathfrak{t}) + \mathcal{b}_3(\mathfrak{x}, \mathfrak{y}, Z, \mathfrak{t}) + \dots \\ &= e^{\mathfrak{x}, \mathfrak{y}, Z} \left(1 - \frac{\mathfrak{t}^\alpha}{\Gamma(\alpha+1)} + \mathfrak{t}^{2\alpha} \Gamma(2\alpha + 1) - \frac{\mathfrak{t}^{3\alpha}}{\Gamma(3\alpha + 1)} + \dots \right) \end{aligned}$$

$$= e^{x+y+Z} \left(1 + \sum_{m=1}^{\infty} \frac{(-t^\alpha)^m}{\Gamma(m\alpha + 1)} \right) = e^{x+y+Z} E_\alpha(-t^\alpha)$$

When $\alpha = 1$, the following result is derived:

$$\begin{aligned} \mathcal{B}(x, y, Z, t) &= \lim \sum \mathcal{B}_n(x, y, Z, t) \\ &= \mathcal{B}_0(x, y, Z, t) + \mathcal{B}_1(x, y, Z, t) + \mathcal{B}_2(x, y, Z, t) + \mathcal{B}_3(x, y, Z, t) \\ &= e^{x+y+Z} \left(1 - \frac{t}{1!} + \frac{t^2}{2!} - \frac{t^3}{3!} \dots \right) \\ &= e^{x+y+Z-t} \end{aligned}$$

Example 2 The fractional-order differential equation is given by:

$${}^{AB}D_t^{\alpha w} = \rho + \frac{\partial^2 w}{\partial Z^2} + \frac{1}{Z} \frac{\partial w}{\partial Z} \dots \quad (17)$$

Based on the initial conditions.

$$\mathfrak{U}(\mathcal{B}, 0) = 1 - Z^2. \quad (18)$$

The Laplace transform is:

$$\iota[w(Z, t)] - \frac{1}{s} (1 - Z^2) - \frac{1}{s^\alpha} \iota \left[\frac{\partial^2 w}{\partial Z^2} + \frac{1}{Z} \frac{\partial w}{\partial Z} \right] = 0. \quad (19)$$

The nonlinear term is:

$$N[\Phi(Z, t, q)] = \iota[\Phi(Z, t, q)] - \frac{1}{s} (1 - Z^2) - \frac{1}{s^\alpha} \iota[\rho] - \frac{1}{s^\alpha} \iota \left[\frac{\partial^2}{\partial Z^2} \Phi(Z, t, q) + \frac{1}{Z} \frac{\partial}{\partial Z} \Phi(Z, q, t) \right] \quad (20)$$

Thus

$$R_r(w_{r-1}(Z, t)) = \iota[w_{r-1}(Z, t)] - \frac{1}{s^\alpha} \iota \left[\frac{\partial^2}{\partial Z^2} w_{r-1} + \frac{1}{Z} \frac{\partial}{\partial Z} \right] - (1 - x_r) \frac{1}{s} (1 - Z^2) - \frac{1}{s^{\alpha+1}} \iota[\rho] (1 - x_r) \quad (21)$$

The r^{th} order deformation equation is

$$\iota[w_r(Z, t) - x_r w_{r-1}(Z, t)] = h R_r(w_{r-1}(Z, t)) \quad (22)$$

The next step is to apply the inverse Laplace transform to obtain the solution in the original domain.

$$w_m(Z, t) - \mathfrak{x}_m w_{m-1}(Z, t) = hu^{-1}[R_m(w_{m-1}(Z, t))]. \quad (23)$$

By solving the above Eq(23), for $r = 1, 2, 3 \dots$, then

$$\begin{aligned} w_0(Z, t) &= 1 - Z^2 \\ w_1(Z, t) &= -\frac{h(-4 + P)}{\Gamma(\alpha + 1)} t^\alpha \\ w_2(Z, t) &= -\frac{h(-4 + P)(1 + h)}{\Gamma(\alpha + 1)} t^\alpha \\ w_3(Z, t) &= -\frac{h(-4 + P)(1 + h)^2}{\Gamma(\alpha + 1)} t^\alpha \\ w_4(Z, t) &= -\frac{h(-4 + P)(1 + h)^3}{\Gamma(\alpha + 1)} t^\alpha \end{aligned}$$

And so on. Concluding that

$$w(\mathcal{Z}, t) = w_0(Z, t) + w_1(Z, t) + w_2(Z, t) + w_3(Z, t) + w_4(Z, t) + \dots$$

Then

$$w(Z, t) = 1 - Z^2 - \frac{h(-4 + P)}{\Gamma(\alpha + 1)} t^\alpha - \frac{h(-4 + P)(1 + h)}{\Gamma(\alpha + 1)} t^\alpha$$

$$- \frac{h(-4 + P)(1 + h)^2}{\Gamma(\alpha + 1)} t^\alpha - \frac{h(-4 + P)(1 + h)^3}{\Gamma(\alpha + 1)} t^\alpha + \dots$$

$$w(Z, t) = 1 - Z^2 - \frac{h(-4 + P)}{\Gamma(\alpha + 1)} t^\alpha [1 + (1 + h) + (1 + h)^2 + (1 + h)^3 + \dots]$$

If $h < 0$, the geometric series will converge as it approaches infinity, and the solution takes the following form:

$$w(Z, t) = 1 - Z^2 + \frac{-4 + p}{\Gamma(\alpha + 1)} t^\alpha$$

$$\text{If } \alpha = 1, \text{ then the solution } w(Z, t) = 1 - Z^2 + (-4 + P) \quad (24)$$

Example 3 Let us examine the subsequent nonlinear equation within the Atangana-Baleanu-Caputo framework:

$${}^{AB}D_t^\delta \psi(\mu, \tau) = -\frac{\partial}{\partial \mu} \left(\frac{12}{\mu} \psi - \mu \right) \psi + \frac{\partial^2}{\partial \mu^2} \psi^2, 0 < \delta \leq 1 \quad (25)$$

$$\text{Subject to the initial condition } \psi(\mu, 0) = \mu^2 \quad (26)$$

By applying the natural transform to both sides of the equation, we obtain:

$$N \left[{}^{ABC}D_{\tau}^{\delta} \psi(\mu, \tau) = \frac{12}{\mu^2} \psi^2 - \frac{12}{\mu} \psi_{\mu} \psi + \psi + (\psi^2)_{\mu\mu} \right], \quad (27)$$

By applying the inverse natural transform to both sides of Eq. (27) and utilizing the initial condition, we obtain;

$$\psi(\mu, \tau) = \mu^2 + N^{-1} \left[\left(1 - \delta + \delta \left(\frac{\mathfrak{U}}{\mathfrak{S}} \right)^{\left[\frac{12}{\mu^2} \right]} \psi^2 - \frac{12}{\mu} \psi_{\mu} \psi + (\psi^2)_{\mu\mu} + \psi \right) \right].$$

By applying homotopy permutation method,

$$\sum_{n=0}^{\infty} \rho^n \psi_n = \mu^2 - \rho N^{-1} \left[\left(1 - \delta + \delta \left(\frac{\mu}{\mathfrak{S}} \right)^{\delta} \right) N \left[\sum_{n=0}^{\infty} \rho^n A_n - \sum_{n=0}^{\infty} \rho^n B_n + \sum_{n=0}^{\infty} \rho^n C_n + \sum_{n=0}^{\infty} \rho^n \psi_n \right] \right]$$

By The following result is obtained,

$$\begin{aligned} \rho^0: \psi_0 &= \mu^2 \\ \rho^1: \psi_1 &= N^{-1} \left[\left(1 - \delta + \delta \left(\frac{\mu}{\mathfrak{S}} \right)^{\delta} \right) N(\rho^0 A_0 - \rho^0 B_0 + \rho^0 C_0 + \rho^0 \psi_0) \right], \\ \rho^2: \psi_2 &= N^{-1} \left[\left(1 - \delta + \delta \left(\frac{\mu}{\mathfrak{S}} \right)^{\delta} \right) N(\rho^1 A_1 - \rho^1 B_1 + \rho^1 C_1 + \rho^1 \psi_1) \right], \\ \text{By the above algorithms,} \\ \psi_0 &= \mu^2 \\ \psi_1 &= \mu^2 \left(1 - \delta + \delta \frac{\tau^{\delta}}{\Gamma(\delta+1)} \right), \\ \psi_2 &= \mu^2 (1 - 2\delta + \delta^2) + (2\delta - 2\delta^2) \frac{\tau^{\delta}}{\Gamma(\delta+1)} + \delta^2 \frac{\tau^{2\delta}}{\Gamma(2\delta+1)}, \end{aligned}$$

Therefore, the series solution $\psi(\mu, \tau)$ is given by

$$\psi(\mu, \tau) = \mu^2 \left[(3 - 3\delta + \delta^2) + (3\delta - 2\delta^2) \frac{\tau^{\delta}}{\Gamma(\delta+1)} + \delta^2 \frac{\tau^{2\delta}}{\Gamma(2\delta+1)} + \dots \right] \quad (29)$$

If we put $\delta = 1$ in (29), we get the approximate and exact solution

$$\psi(\mu, \tau) = \mu^2 \left(1 + \frac{\tau}{1!} + \frac{\tau^2}{2!} + \dots \right) = \mu^2 \exp(\tau). \quad (30)$$

Example 4: Let us examine the non-homogeneous fractional equation in the following form:

$${}^{AB}D_{\tau}^{\mathfrak{S}}(\rho, \tau) = \mathfrak{x}(\rho, \tau) - \frac{1}{2} \frac{\partial \mathfrak{x}^2(\rho, \tau)}{\partial \rho} - \mathfrak{x}^2(\rho, \tau) - e^{\tau-\rho}. \quad (31)$$

Subject to initial condition $\mathfrak{x}(\rho, 0) = 1 - e^{-\rho}$.

Applying the (NT) to both sides of equation (31), we obtain:

$$N \left(\mathfrak{x}(\rho, \tau) = \frac{1-e^{-\rho}}{s^{\mathfrak{S}}} + \frac{w^{\mathfrak{S}} e^{-\rho}}{s^{\mathfrak{S}}(s^{\mathfrak{S}}-w^{\mathfrak{S}})} + \frac{w^{\mathfrak{S}}}{s^{\mathfrak{S}}} \left(N \left(\mathfrak{x}(\rho, \tau) - \frac{1}{2} \frac{\partial \mathfrak{x}^2(\rho, \tau)}{\partial \rho} - \mathfrak{x}^2(\rho, \tau) \right) \right) \right). \quad (32)$$

By taking the inverse Natural transform of equation (32), we get

$$\mathfrak{x}(\rho, \tau) = 1 - e^{-\rho} E_{\mathfrak{S}}(\tau) + N^{-1} \left(\frac{w^{\mathfrak{S}}}{s^{\mathfrak{S}}} \left(N \left(\mathfrak{x}(\rho, \tau) - \frac{1}{2} \frac{\partial \mathfrak{x}^2(\rho, \tau)}{\partial \rho} - \mathfrak{x}^2(\rho, \tau) \right) \right) \right). \quad (33)$$

We now suppose that

$$\mathfrak{x}(\rho, \tau) = \sum_{n=0}^{\infty} \rho^n \mathfrak{x}_n(\rho, \tau) \quad (35)$$

Subsequently, by applying equation (34), we can express equation (33) as:

$$\sum_{n=0}^{\infty} \rho^n \mathfrak{x}_n = 1 - e^{-\rho} E_{\mathfrak{S}}(\tau) + N^{-1} \left(\frac{w^{\mathfrak{S}}}{s^{\mathfrak{S}}} \left(N \sum_{n=0}^{\infty} \rho^n \mathfrak{x}_n - \frac{1}{2} \sum_{n=0}^{\infty} \rho^n H_n - \sum_{n=0}^{\infty} \rho^n G_n \right) \right) \quad (36)$$

Where H_n and G_n these are the Adomian polynomials that represent the $\mathfrak{x}^2(\rho, \tau)$ respectively.

By equating both sides of equation (36), we can straightforwardly derive the recursive relation as:

$$\begin{aligned} \rho^0: \mathfrak{x}_0(\rho, \tau) &= 1 - e^{-\rho} E_{\mathfrak{S}}(\tau) \\ \rho^1: \mathfrak{x}_1(\rho, \tau) &= N^{-1} \left(\frac{W^{\mathfrak{S}}}{s^{\mathfrak{S}}} \left(N \left(\mathfrak{x}_0 - \frac{1}{2} H_0 - G_0 \right) \right) \right) \\ &= N^{-1} \left(\frac{w^{\mathfrak{S}}}{s^{\mathfrak{S}}} \left(N \left(\mathfrak{x}_0(\rho, \tau) \frac{1}{2} \frac{\partial \mathfrak{x}_0^2(\rho, \tau)}{\partial \rho} - \mathfrak{x}_0^2(\rho, \tau) \right) \right) \right) = 0 \end{aligned}$$

$$\rho^2: \mathfrak{x}_2(\rho, \tau) = N^{-1} \left(\frac{W^{\mathfrak{S}}}{s^{\mathfrak{S}}} \left(N \left(\mathfrak{x}_1 - \frac{1}{2} H_1 - G_1 \right) \right) \right)$$

$$:\mathfrak{x}(\rho, \tau) = 1 - e^{-\rho} E_{\mathfrak{S}}(\tau). \quad (37)$$

$$\text{For } \mathfrak{S} = 1 \text{ The given solution converges to the precise solution } . \mathfrak{x}(\rho, \tau) = 1 - e^{\tau-\rho}. \quad (38)$$

Example 5: Let us consider the nonlinear system of time-fractional differential equations governed by the Atangana-Baleanu-Caputo operator:

$$\begin{aligned} {}^{AB}D_t^\alpha \mathfrak{b}(\mathfrak{x}, \mathfrak{t}) - \mathfrak{b}_{\mathfrak{x}\mathfrak{x}} - 2\mathfrak{b}\mathfrak{b}_{\mathfrak{x}} + (\mathfrak{b}w)_{\mathfrak{x}} &= 0 \\ {}^{AB}D_t^\mathfrak{S} w(\mathfrak{x}, \mathfrak{t}) - w_{\mathfrak{x}\mathfrak{x}} - 2w w_{\mathfrak{x}} + (\mathfrak{b}w)_{\mathfrak{x}} &= 0. \end{aligned} \quad (39)$$

where $0 < \alpha, \mathfrak{S} \leq 1$ and the initial condition are $\mathfrak{b}(\mathfrak{x}, 0) = \sin(\mathfrak{x}), w(\mathfrak{x}, 0) = \sin(\mathfrak{x})$

$$N \left\{ {}^{ABC}D_t^\alpha \mathfrak{b}(\mathfrak{x}, \mathfrak{t}) = \mathfrak{b}(\mathfrak{x}, 0) + \left(1 - \alpha + \alpha \left(\frac{\mathfrak{t}}{s} \right)^\alpha \right) N \left(\frac{\partial^2 \mathfrak{b}}{\partial \mathfrak{x}^2} + 2\mathfrak{b} \frac{\partial \mathfrak{b}}{\partial \mathfrak{x}} - \frac{\partial}{\partial \mathfrak{x}} (\mathfrak{b}w) \right) \right.$$

$$N\{ {}^{ABC}D_t^{\mathfrak{J}} \mathcal{L}(\mathfrak{x}, t) \} = w(\mathfrak{x}, 0) + \left(1 - \mathfrak{J} + \mathfrak{J} \left(\frac{\mathfrak{U}}{s}\right)^{\mathfrak{J}}\right) N \left(\frac{\partial^2 w}{\partial \mathfrak{x}^2} + 2w \frac{\partial w}{\partial \mathfrak{x}} - \frac{\partial}{\partial \mathfrak{x}} (\mathcal{L}w) \right) \quad (40)$$

By applying the (NT) to both sides of eq. (39), we obtain:

$$\begin{aligned} \mathcal{L}(\mathfrak{x}, t) &= \sin(\mathfrak{x}) + N^{-1} \left(\left(1 - a + a \left(\frac{\mathfrak{U}}{s}\right)^a\right) N \left(\frac{\partial^2 \mathcal{L}}{\partial \mathfrak{x}^2} + 2\mathcal{L} \frac{\partial \mathcal{L}}{\partial \mathfrak{x}} - \frac{\partial}{\partial \mathfrak{x}} (\mathcal{L}w) \right) \right) \\ w(\mathfrak{x}, t) &= \sin(\mathfrak{x}) + N^{-1} \left(\left(1 - \mathfrak{J} + \mathfrak{J} \left(\frac{\mathfrak{U}}{s}\right)^{\mathfrak{J}}\right) N \left(\frac{\partial^2 w}{\partial \mathfrak{x}^2} + 2w \frac{\partial w}{\partial \mathfrak{x}} - \frac{\partial}{\partial \mathfrak{x}} (\mathcal{L}w) \right) \right) \end{aligned} \quad (41)$$

Now, we express the solution as an infinite series given by:

$$\mathcal{L}(\mathfrak{x}, t) = \sum_{n=0}^{\infty} \rho^n \mathcal{L}_n(\mathfrak{x}, t), w(\mathfrak{x}, t) = \sum_{n=0}^{\infty} \rho^n w_n(\mathfrak{x}, t).$$

and the nonlinear terms can be expanded as

$$\mathcal{L}\mathcal{L}_{\mathfrak{x}} - (\mathcal{L}w)_{\mathfrak{x}} = \sum_{n=0}^{\infty} \rho^n A_n(\mathfrak{x}, t), ww_{\mathfrak{x}} - (\mathcal{L}w)_{\mathfrak{x}} = \sum_{n=0}^{\infty} \rho^n B_n(\mathfrak{x}, t). \quad (42)$$

Substituting(41) and (40) in(39) and by applying (HPM) on Eq.(39)

$$\sum_{n=0}^{\infty} \rho^n \mathcal{L}_n = \sin(\mathfrak{x}) - \rho N^{-1} \left(\left(1 - a + a \left(\frac{\mathfrak{U}}{s}\right)^a\right) N \left(\sum_{n=0}^{\infty} \rho^n \frac{\partial^2 \mathcal{L}_n}{\partial \mathfrak{x}^2} + \sum_{n=0}^{\infty} \rho^n A_n \right) \right) \quad (43)$$

By equating both sides of equation (43),

$$\begin{aligned} \rho^0: \mathcal{L}_0 &= \mathcal{L}(\mathfrak{x}, 0), \rho^0: w_0 = w(\mathfrak{x}, 0) \\ \rho^1: \mathcal{L}_1 &= -N^{-1} \left(\left(1 - a + a \left(\frac{\mathfrak{U}}{s}\right)^a\right) N \left(\frac{\partial^2 \mathcal{L}_0}{\partial \mathfrak{x}^2} + A_0 \right) \right) \\ \rho^1: w_1 &= -N^{-1} \left(\left(1 - \mathfrak{J} + \mathfrak{J} \left(\frac{\mathfrak{U}}{s}\right)^{\mathfrak{J}}\right) N \left(\frac{\partial^2 \mathcal{L}_0}{\partial \mathfrak{x}^2} + B_0 \right) \right) \\ \rho^2: \mathcal{L}_2 &= -N^{-1} \left(\left(1 - a + a \left(\frac{\mathfrak{U}}{s}\right)^a\right) N \left(\frac{\partial^2 \mathcal{L}_0}{\partial \mathfrak{x}^2} + A_1 \right) \right) \\ \rho^2: w_2 &= -N^{-1} \left(\left(1 - \mathfrak{J} + \mathfrak{J} \left(\frac{\mathfrak{U}}{s}\right)^{\mathfrak{J}}\right) N \left(\frac{\partial^2 \mathcal{L}_0}{\partial \mathfrak{x}^2} + B_1 - C_1 \right) \right) \end{aligned}$$

By the above algorithms, $\mathcal{L}_0 = \sin(\mathfrak{x}), w_0 = \sin(\mathfrak{x})$

$$\begin{aligned} \mathcal{L}_1 &= -\sin(\mathfrak{x}) \left(1 - a + a \frac{t^a}{\Gamma(a+1)} \right) \\ w_1 &= -\sin(\mathfrak{x}) \left(1 - \mathfrak{J} + \mathfrak{J} \frac{t^{\mathfrak{J}}}{\Gamma(\mathfrak{J}+1)} \right) \end{aligned}$$

$$\begin{aligned}
v_2 &= \sin(x) \left((1-a)^2 + (2a-2a^2) \frac{t^a}{\Gamma(a+1)} + \right. \\
&\quad \left. a^2 \frac{t^{2a}}{\Gamma(2a+1)} + 2(a-B)\cos(x) \left((1-a) + (2a-1) \frac{t^a}{\Gamma(a+1)} - a^2 \frac{t^{2a}}{\Gamma(2a+1)} \right) \right) \\
w_2 &= \sin(x) \left((1-\Im)^2 + (2\Im-2\Im^2) \frac{t^\Im}{\Gamma(\Im+1)} + \Im^2 \frac{t^{2\Im}}{\Gamma(2\Im+1)} + 2(\Im-B)\cos(x) \left((1-\Im) + (2\Im \right. \right. \\
&\quad \left. \left. - 1) \frac{t^\Im}{\Gamma(\Im+1)} - \Im^2 \frac{t^{2\Im}}{\Gamma(2\Im+1)} \right) \right) \\
\theta(x,t) &= \sin(x) \left((1-a+a^2) + (a-2a^2) \frac{t^a}{\Gamma(a+1)} + a^2 \frac{t^{2a}}{\Gamma(2a+1)} + \right. \\
&\quad \left. 2(a-\Im)\cos(x) \left((1-a)(2a-1) \frac{t^a}{\Gamma(a+1)} - a^2 \frac{t^{2a}}{\Gamma(2a+1)} + \dots \right) \right) \\
w(x,t) &= \sin(x) \left((1-\Im+\Im^2) + (\Im-2\Im^2) \frac{t^\Im}{\Gamma(\Im+1)} + \Im^2 \frac{t^{2\Im}}{\Gamma(2\Im+1)} + \right. \\
&\quad \left. 2(\Im-\Im)\cos(x) \left((1-\Im)(2\Im-1) \frac{t^\Im}{\Gamma(\Im+1)} - \Im^2 \frac{t^{2\Im}}{\Gamma(2\Im+1)} \right) \right] + \dots \tag{44}
\end{aligned}$$

If we substitute $\alpha \rightarrow 1$ in (44)

$$\begin{aligned}
\mathcal{U}(x,t) &= \sin(x) \left(1 - t + \frac{t^2}{2!} - \dots \right), \theta(x,t) = \sin(x) \left(1 - t + \frac{t^2}{2!} - \dots \right) \\
\mathcal{U}(x,t) &= \sin(x)e^{-t}, \theta(x,t) = \sin(x)e^{-t}
\end{aligned}$$

5- Conclusion

In this research, a hybrid method combining the Homotopy approach with the Natural Transformation has been presented as an effective tool for solving fractional differential equations, whether linear or nonlinear. The results demonstrated that this method provides fast and accurate approximate solutions, enabling efficient handling of complex mathematical problems involving fractional derivatives. Furthermore, this method reduces computational demands, making it a powerful and reliable tool for studying and analyzing systems governed by fractional differential equations. These findings represent an important step toward improving numerical solution methods in various fields such as mathematical physics and engineering, making this approach a promising tool for future applications in solving fractional differential equations..

References:

- [1] A. M. Amall and H. A. Yasmmin, "Algorithm for solving linear Volterra fractional integro-differential equations using the Elzaki Transform," *Iban-Al-Haitham*, 2017.
- [2] R. Pavel, "A semi-explicit multi-step method for solving incompressible Navier-Stokes equations," 2018.
- [3] H. Jafari, C. M. Khalique, and M. Nazari, "Application of the Laplace decomposition method to solve linear and nonlinear fractional diffusion-wave equations," *Applied Mathematics Letters*, 2011.
- [4] H. K. Jassim and S. A. Khafif, "Solving Burgers and coupled Burgers equations of fractional order using the SVM," *Progress in Fractional Differential Equations and Applications*, 2021.
- [5] H. K. Eaved, H. K. Jassim, and M. G. Mohammed, "A novel method for the analytical solution of differential equations in mathematical physics," *IOP Conference Series: Materials Science and Engineering*, 2020.
- [6] S. Das and R. Kumar, "Approximate analytical solution of fractional gas dynamics equations," *Applied Mathematics and Computation*, vol. 217, no. 24, pp. 9905-9915, 2011.
- [7] H. Jafari, H. K. Jassim, and C. Nguyen V. T., "Laplace Decomposition Method for solving the two-dimensional diffusion problem in fractal heat transfer," *Fractals*, vol. 32, no. 4, pp. 1-6, 2024.
- [8] H. Ahmad and H. K. Jassim, "Analytical technique for approximate solutions of nonlinear fractional PDEs," *Journal of Education for Pure Science-University of Thi-Qar*, vol. 14, no. 1, pp. 107-116, 2024.
- [9] H. Jafari, H. K. Jassim, A. Ansari, and C. Nguyen V. T., "Local Fractional Variational Iteration Transform Method: A tool for solving local fractional partial differential equations," *Fractals*, vol. 32, no. 4, pp. 1-8, 2024.
- [10] D. Baleanu, M. H. Abadi, A. Jajarmi, K. Z. Vahid, and J. J. Nieto, "A comparative study of the general fractional model for COVID-19 with isolation and quarantine effects," *Alexandria Engineering Journal*, vol. 61, pp. 4779-4791, 2022.
- [11] S. Kumar, A. Kumar, S. Momani, M. Aldhaifallah, and K. S. Nisar, "Numerical solutions of nonlinear fractional models in the appearance of strip patterns in two-dimensional systems," *Advances in Differential Equations*, 2019, pp. 1–19.
- [12] J. Liu, M. Nadeem, and L. F. Iambor, "Yang homotopy perturbation transform approach for solving multi-dimensional diffusion problems with time fractional derivatives," *Scientific Reports*, vol. 13, no. 21855, 2023.

- [13] S. M. Aydogan, D. Baleanu, J. F. G. Aguilar, S. Rezapour, and M. E. Samei, "Approximate endpoint solutions for a class of fractional q-differential inclusions," *Fractals*, vol. 28, no. 8, p. 2040029, 2020.
- [14] S. A. Issa and H. Tajadodi, "Yang Adomian Decomposition Method for solving PDEs," *Journal of Education for Pure Science-University of Thi-Qar*, vol. 14, no. 2, pp. 14-25, 2024.
- [15] H. K. Jassim and M. A. Hussein, "A novel formulation of the fractional derivative with the order and without the singular kernel," *Mathematics*, vol. 10, no. 21, p. 4123, 2022.
- [16] H. Ahmad and J. J. Nasar, "Atangana-Baleanu Fractional Variational Iteration Method for solving fractional-order Burger's equations," *Journal of Education for Pure Science-University of Thi-Qar*, vol. 14, no. 2, pp. 26-35, 2024.
- [17] Yasser, M. Taimah and Jassim, H. Kamil (2025). A New Integral Transform for Solving Integral and Ordinary Differential Equations. *Mathematics and Computational Sciences*, doi: 10.30511/mcs.2025.2045547.1254.
- [18] N. R. Seewn, M. T. Yasser, D. Ziane, An Analytical Approach to Nonlinear Fractional Differential Equations Using Daftardar-Jafari Method, *Journal of Education for Pure Science-University of Thi-Qar*, 15(1)(2025) 62-73. doi.org/10.32792/jeps.v15i1.651
- [19] N. R. Seewn, M. T. Yasser, H. Tajadodi, An Efficient Approach for Nonlinear Fractional PDEs: Elzaki Homotopy Perturbation Method, *Journal of Education for Pure Science-University of Thi-Qar*, 15(1)(2025) 89-99.
- [20] S. K. Ntouyas and J. Tariboon, "Fractional integral problems for Hadamard–Caputo fractional Langevin differential inclusions," *Journal of Applied Mathematics and Computation*, vol. 51, pp. 13–33, 2016.
- [21] Z. H. Ali and K. H. Yasser, "Perturbed Taylor expansion for bifurcation solutions of singularly parameterized ordinary differential equations and differential algebraic equations," *Journal of Education for Pure Science-University of Thi-Qar*, vol. 10, no. 2, pp. 219-234, 2020.
- [22] M. Caputo and M. Fabrizio, "A new definition of fractional derivative without singular kernel," *Progress in Fractional Differentiation and Applications*, vol. 1, no. 2, pp. 73–85, 2015.
- [23] H. S. Kadhem and S. Q. Hasan, "A new double Sumudu transform iterative method for solving fractional partial differential equations," *Journal of Education for Pure Science-University of Thi-Qar*, vol. 9, no. 2, pp. 158-171, 2019.
- [24] J. Losada and J. J. Nieto, "Properties of the new fractional derivative without singular kernel," *Progress in Fractional Differentiation and Applications*, vol. 1, no. 2, pp. 87–92, 2015.
- [25] J. J. Nasar and H. Tajadodi, "Approximate solutions of 2D-Burger's equations," *Journal of Education for Pure Science-University of Thi-Qar*, vol. 10, no. 3, pp. 1-11, 2024.

- [26] M. A. Hussein, "Approximate methods for solving fractional differential equations," *Journal of Education for Pure Science-University of Thi-Qar*, vol. 12, no. 2, pp. 32-40, 2022.
- [27] G. A. Hussein and D. Ziane, "New approximation solutions for fractional order biological population models," *Journal of Education for Pure Science-University of Thi-Qar*, vol. 10, no. 3, pp. 1-20, 2024.
- [28] A. R. Saeid and L. K. Alzaki, "Analytical solutions for the nonlinear homogeneous fractional biological equation using a local fractional operator," *Journal of Education for Pure Science-University of Thi-Qar*, vol. 13, no. 3, pp. 1-17, 2023.
- [29] Z. S. Matar, S. R. Abdul Kadeem, and A. H. Naser, "A new approach to video summary generation," *Journal of Education for Pure Science-University of Thi-Qar*, vol. 10, no. 3, pp. 1-20, 2024.
- [30] N. H. Tuan, H. Mohammadi, and S. Rezapour, "Mathematical model for COVID-19 transmission using the Caputo fractional derivative," *Chaos, Solitons & Fractals*, vol. 140, p. 110107, 2020.
- [31] Z. R. Jawad and A. K. Al-Jaberi, "A modified fourth-order partial differential equations model based on Isophote direction for noise image removal," *Journal of Education for Pure Science-University of Thi-Qar*, vol. 10, no. 3, p. 115, 2024.
- [32] T. El-Danaf and A. Hadhoud, "Parametric spline functions for solving the one-time fractional Burgers' equation," *Applied Mathematical Modeling*, vol. 36, pp. 4557-4564, 2012.
- [33] H. K. Jassim and M. G. Mohammed, "Natural homotopy perturbation method for solving nonlinear fractional gas dynamics equations," *International Journal of Nonlinear Analytical Applications*, vol. 12, no. 1, pp. 813-821, 2021.