

Semi-Analytical Solutions of Nonlinear Time-Fractional Fornberg-Whitham Equation

Mohammed Taimah Yasser ¹, , Naser Rhaif Seewn ², , Mountassir Hamdi Cherif ³

³Hight School of Electrical and Energetics Engineering of Oran (ESGEE-Oran),

Oran, Algeria; mountassir27@yahoo.fr

^{1,2} Department of Mathematics Faculty of Education for Pure Sciences Thi-Qar University

* Corresponding email: edpma4m23@utq.edu.iq

Received 15/ 2 /2025,

Accepted 22/ 4 /202X,

Published 1/ 6 /2025



This work is licensed under a [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/).

Abstract:

In this paper, we present an approximate solution for the Fornberg-Whitham equation using fractional operators, specifically the Atangana-Baleanu derivative, and apply the Yang Adomian Decomposition Method (YADM). The results indicate high efficiency and show a significant agreement with the exact solution.

Keywords: Adomian Decomposition Method; Yang transform; Atangana-Baleanu derivative.

1-Introduction

In recent decades, fractional calculus (FC) has been applied to numerous phenomena across natural sciences, engineering, fluid dynamics, life sciences, and other practical sciences. FC provides a robust framework for effectively describing these phenomena using fractional mathematical tools. Fractional derivatives (FDs) are especially valuable in capturing the hereditary and memory characteristics of various processes and materials. Applications of FDs appear in multiple engineering and scientific challenges, including diffusion and reaction processes, frequency-dependent signal processing, system modeling, damping behaviors in materials, and the relaxation and creep of viscoelastic substances. [1-3].

The examination of nonlinear wave equations and their answers holds vital significance across many academic disciplines. One of the most compelling solutions for nonlinear fractional partial differential equations (FPDEs) is the concept of traveling waves. Nonlinear FPDEs are frequently associated with complex physical and mechanical processes, making the exact solutions of these equations, particularly traveling wave solutions, highly desirable. Certain FPDEs, such as the Korteweg-de Vries and Camassa-Holm equations, exhibit various traveling

wave solutions and are instrumental in modeling nonlinear, Omnidirectional dispersing waves in shallow depths. [4] Additionally, the Fornberg–Whitham equation (FWE) is notably significant in mathematical physics, with applications spanning multiple scientific disciplines.

The Fornberg–Whitham equation (FWE) [5],[6] is formulated as follows:

$$D_t^\vartheta u - u_{xxt} + u_x + uu_x = 3u_x u_{xx} + uu_{xxx}$$

This equation describes the qualitative behavior of wave breaking in nonlinear dispersive waves. The FWE is known to allow peakon solutions, providing a numerical framework for understanding limiting wave heights and the occurrence of wave breaks. In 1978, Fornberg and Whitham introduced a peaked solution given by:

$$u(x, t) = Ce^{\frac{x}{2} - \frac{2t}{3}}$$

where C is a constant. The analysis of FWEs has been approached through various numerical and analytical methods, including the Laplace decomposition technique [7], Lie symmetry [8], variational iteration technique [9], differential transformation technique [10], new iterative technique [11], homotopy-perturbation technique [12], and homotopy analysis transform technique [13], In addition to the numerous scientific studies in which solutions and methods for solving differential equations of both fractional and classical orders have been proposed [14-17], [19-32]

The Adomian Decomposition Method (ADM), pioneered by George Adomian, has emerged as a robust analytical framework for solving nonlinear differential equations. Unlike iterative or perturbation-based techniques, ADM systematically decomposes nonlinear terms into computable Adomian polynomials, enabling precise handling of complex nonlinearities without restrictive linearization assumptions [29–31]. This method has been extensively applied to fractional-order systems, particularly under operators like Caputo, Riemann–Liouville, and the modern Atangana–Baleanu derivative, which preserves memory effects critical for modeling hereditary phenomena [33,34].

A key advancement in ADM’s application to fractional equations lies in its integration with integral transforms. In this study, the Yang transform is combined with ADM to streamline the resolution of the time-fractional Fornberg–Whitham equation (FWE). While traditional mesh-based methods [18] provide discrete approximations, the Yang-ADM synergy generates continuous, rapidly convergent series solutions, avoiding the numerical overhead of spatial discretization. Recent innovations in ADM, such as optimized algorithms for Adomian polynomial generation [34], have further enhanced its computational efficiency, addressing historical critiques about polynomial complexity.

This manuscript advances the analytical resolution of the fractional FWE using the Atangana–Baleanu fractional **operator**, a non-singular kernel derivative that accurately models memory-dependent wave-breaking dynamics. While prior studies employed methods like differential transformation [10] or homotopy techniques [12,13], their reliance on linearization or small parameters limits generality. In contrast, ADM’s direct decomposition framework circumvents these constraints, yielding semi-analytical solutions in series form with minimal computational effort. The method’s convergence, rigorously established for fractional PDEs [35,36], ensures reliability, while its avoidance of discretization errors aligns with the need for precise physical insights into wave behavior.

Study Contributions:

1. **Yang-ADM Hybridization:** Unifies the Yang transform’s operational simplicity with ADM’s nonlinear decomposition, enabling efficient handling of Atangana–Baleanu fractional terms.
2. **Streamlined Polynomial Computation:** Leverages recent algorithmic improvements [38] to accelerate Adomian polynomial generation, enhancing practicality.

3. **Memory-Effect Preservation:** Utilizes the Atangana-Baleanu operator to maintain the FWE's intrinsic memory properties, critical for modeling wave-breaking.

This work underscores ADM's adaptability to modern fractional calculus challenges, demonstrating its superiority over iterative and mesh-dependent methods in balancing accuracy, efficiency, and analytical transparency. By achieving semi-analytical solutions with clear convergence, the study provides a template for solving broader classes of FPDEs prevalent in engineering and physics.

2-Preliminaries

Definition.1. The Caputo fractional derivative of function $f(x)$, $x > 0$ is defined by [32]

$$D_x^\alpha f(t) = \begin{cases} \frac{1}{\Gamma(n-\alpha)} \int_0^x (x-t)^{n-\alpha-1} f^{(n)}(t) dt, & n-1 < \alpha \leq n \in \mathbb{N} \\ \frac{d^n}{dx^n} f(x), & \alpha = n \in \mathbb{N} \end{cases} \quad (1)$$

Note. 1. From **Definition 1**, the following **outcome** is **derived**:

$$D_t^\alpha t^\beta = \begin{cases} \frac{\Gamma(\beta+1)}{\Gamma(\beta-\alpha+1)} t^{\beta-\alpha}, & n-1 < \alpha \leq n, \quad \beta > n-1, \beta \in \mathbb{R} \\ 0 & n-1 < \alpha \leq n, \quad n > \beta, \beta \in \mathbb{N} \end{cases} \quad (2)$$

Definition.2. The Atangana-Baleanu fractional derivative is expressed as[33][34]:

$${}^{AB}_a D_t^\vartheta u(t) = \frac{\mathcal{M}(\vartheta)}{1-\vartheta} \int_a^t \mathbb{E}_\vartheta \left(\frac{-\vartheta(t-x)^\vartheta}{1-\vartheta} \right) u'(x) dx, \quad (3)$$

Where $0 < \vartheta < 1$ and $\mathcal{M}(\vartheta)$ is a normalization function, such that $\mathcal{M}(0) = \mathcal{M}(1) = 1$.

Definition.3. The Atangana-Baleanu Fractional Integral (ABFI) of order ϑ is defined as follows[34]:

$${}^{AB}_a I_t^\vartheta u(t) = \frac{1-\vartheta}{\mathcal{M}(\vartheta)} u(t) + \frac{\vartheta}{\mathcal{M}(\vartheta)\Gamma(\vartheta)} \int_a^t (t-x)^{\vartheta-1} u(x) dx, \quad (4)$$

Where $0 < \vartheta < 1$ and $\mathcal{M}(\vartheta)$ is a normalization function, such that $\mathcal{M}(0) = \mathcal{M}(1) = 1$.

Definition.4. The Yang transform YT is stated as[35]

$$Y\{u(t)\} = \int_0^\infty e^{-\frac{t}{\nu}} u(t) dt, t > 0, \quad (5)$$

with ν representing the transform variable.

Some of properties of it:

1. $Y\{1\} = \nu$
2. $Y\{t\} = \nu^2$
3. $Y\{t^n\} = \nu^{n+1} n!$
4. $Y\{t^\vartheta\} = \nu^{\vartheta+1} \Gamma(\vartheta+1), \vartheta \in \mathbb{R}$

Theorem. The Yang transform of fractional order derivative is defined by

$$a. \quad Y\left\{{}^C D_x^\vartheta u(t)\right\} = \frac{Y\{u(t)\}}{\nu^\vartheta} - \sum_{k=0}^{n-1} \frac{u^{(k)}(0)}{\nu^{\alpha-k-1}}, \quad n-1 < \vartheta \leq n \quad (6)$$

$$b. \quad Y\left\{{}^{AB}_a D_t^\vartheta u(t)\right\} = \frac{\mathcal{M}(\vartheta)}{1-\vartheta+\vartheta\nu^\vartheta} (Y\{u(t)\} - \nu u(0)), \quad 0 < \vartheta \leq 1, \quad (7)$$

Proof. The proof of (a) as in [36]. To prove (b) We take the transformation into Eq(3) and then utilize the convolution property to obtain the desired result after simplification. Thus, the proof is completed

Definition 5. The Mittag-Leffler function with two Parameters is defined by [37], [38]

$$\mathbb{E}_{\vartheta, \rho}(Z) = \sum_{n=0}^{\infty} \frac{Z^n}{\Gamma(n\vartheta + \rho)}, \quad \alpha, \rho, Z \in \mathbb{C}, \quad \operatorname{Re}(\vartheta) > 0, \quad \operatorname{Re}(\rho) > 0$$

Note.2. From definition5. The following result is obtained:

$$(1) \quad \mathbb{E}_{2,1}(x^2) = \cos h(x).$$

$$(2) \quad \mathbb{E}_{2,2}(x^2) = \frac{\sin h(x)}{x}.$$

$$(3) \quad \mathbb{E}_{2,3}(x^2) = \frac{1}{x^2} [-1 + \cos h(x)].$$

3-Fornberg-Whitham equation with Atangana-Baleanu fractional derivative

Consider the fractional Fornberg-Whitham equation

$${}^{AB}\mathcal{D}_t^\vartheta u - u_{xxt} + u_x + uu_x = 3u_x u_{xx} + uu_{xxx}, \quad 0 < \vartheta \leq 1, \quad (8)$$

With initial condition

$$u(x, 0) = g(x), \quad (9)$$

Through the application of the Yang transform to Equation (8), we get:

$$Y\{{}^{AB}\mathcal{D}_t^\vartheta u\} = Y\{u_{xxt} - u_x - uu_x + 3u_x u_{xx} + uu_{xxx}\},$$

Or equivalently,

$$\frac{[Y\{u\} - v u(x, 0)]}{1 - \vartheta + \vartheta v^\vartheta} = Y\{u_{xxt} - u_x - uu_x + 3u_x u_{xx} + uu_{xxx}\},$$

Rearrange,

$$Y\{u\} = v u(x, 0) + (1 - \vartheta + \vartheta v^\vartheta) Y\{u_{xxt} - u_x - uu_x + 3u_x u_{xx} + uu_{xxx}\},$$

Taking inverse of Yang Transform, we obtain

$$u(x, t) = u(x, 0) + Y^{-1}[(1 - \vartheta + \vartheta v^\vartheta) Y\{u_{xxt} - u_x - uu_x + 3u_x u_{xx} + uu_{xxx}\}], \quad (10)$$

Let

$$u(x, t) = \sum_{n=0}^{\infty} u_n, \quad (11)$$

$$uu_x = \sum_{n=0}^{\infty} A_n, \quad (12)$$

$$u_x u_{xx} = \sum_{n=0}^{\infty} B_n, \quad (13)$$

$$uu_{xxx} = \sum_{n=0}^{\infty} C_n, \quad (14)$$

Where

$$A_0 = u_0 u_{0x}, \quad B_0 = u_{0x} u_{0xx}, \quad C_0 = u_0 u_{0xxx},$$

$$A_1 = u_0 u_{1x} + u_1 u_{0x}, \quad B_0 = u_{0x} u_{1xx} + u_{1x} u_{0xx}, \quad C_1 = u_0 u_{1xxx} + u_1 u_{0xxx},$$

$$A_2 = u_0 u_{2x} + u_1 u_{1x} + u_2 u_{0x}, \quad B_2 = u_{0x} u_{2xx} + u_{1x} u_{1xx} + u_{2x} u_{0xx},$$

$$C_2 = u_0 u_{2xxx} + u_1 u_{1xxx} + u_2 u_{0xxx},$$

⋮

By combining Equation (11), (12) , (13) and (14) into Equation (10) yields the following result

$$\sum_{n=0}^{\infty} u_n = u(x, 0) + Y^{-1} \left[(1 - \vartheta + \vartheta v^{\vartheta}) Y \left\{ \sum_{n=0}^{\infty} u_{nxx} - \sum_{n=0}^{\infty} u_{nx} - \sum_{n=0}^{\infty} A_n + 3 \sum_{n=0}^{\infty} B_n + \sum_{n=0}^{\infty} C_n \right\} \right], \quad (15)$$

We present recursive relations as a continuation of the FADM,

$$u_0 = (x, 0),$$

$$u_{n+1} = Y^{-1} \left[(1 - \vartheta + \vartheta v^{\vartheta}) Y \{ u_{nxx} - u_{nx} - A_n + 3B_n + C_n \} \right], \quad n \geq 0, \quad (16)$$

Consequently, the solution for Equation (8) is

$$u(x, t) = u_0 + u_1 + u_2 + \cdots \quad (17)$$

4-Application

Example.1. Consider the fractional Fornberg-Whitham equation

$${}^{AB}\mathfrak{D}_t^{\vartheta} u - u_{xxt} + u_x + uu_x = 3u_x u_{xx} + uu_{xxx}, \quad (18)$$

With initial condition

$$u(x, 0) = e^{\frac{x}{2}}.$$

using eq.(16), we obtain

$$u_0 = e^{\frac{x}{2}},$$

$$u_1 = - \left[1 - \vartheta + \frac{\vartheta t^{\vartheta}}{\Gamma(\vartheta + 1)} \right] \frac{1}{2} e^{\frac{x}{2}},$$

$$u_2 = - \left[(1 - \vartheta) \frac{\vartheta t^{\vartheta-1}}{\Gamma(\vartheta)} + \frac{\vartheta^2 t^{2\vartheta-1}}{\Gamma(2\vartheta)} \right] \frac{1}{8} e^{\frac{x}{2}} + \left[(1 - \vartheta)^2 + 2(1 - \vartheta) \frac{\vartheta t^{\vartheta}}{\Gamma(\vartheta + 1)} + \frac{\vartheta^2 t^{2\vartheta}}{\Gamma(2\vartheta + 1)} \right] \frac{1}{4} e^{\frac{x}{2}}.$$

$$u_3 = - \left[(1-\vartheta)^2 \vartheta \frac{t^{\vartheta-2}}{\Gamma(\vartheta-1)} + 2(1-\vartheta) \frac{\vartheta^2 t^{2\vartheta-2}}{\Gamma(2\vartheta-1)} + \frac{\vartheta^3 t^{3\vartheta-2}}{\Gamma(3\vartheta-1)} \right] \frac{1}{32} e^{\frac{x}{2}} \\ + \left[3(1-\vartheta)^2 \vartheta \frac{t^{\vartheta-1}}{\Gamma(\vartheta)} + 5(1-\vartheta) \frac{\vartheta^2 t^{2\vartheta-1}}{\Gamma(2\vartheta)} + 2 \frac{\vartheta^3 t^{3\vartheta-1}}{\Gamma(3\vartheta)} \right] \frac{1}{16} e^{\frac{x}{2}} \\ - \left[(1-\vartheta)^3 + 3(1-\vartheta)^2 \vartheta \frac{t^{\vartheta}}{\Gamma(\vartheta+1)} + 3(1-\vartheta) \frac{\vartheta^2 t^{2\vartheta}}{\Gamma(2\vartheta+1)} + \frac{\vartheta^3 t^{3\vartheta}}{\Gamma(3\vartheta+1)} \right] \frac{1}{8} e^{\frac{x}{2}}.$$

Then ,

$$u(x, t) = e^{\frac{x}{2}} - \left[1 - \vartheta + \frac{\vartheta t^{\vartheta}}{\Gamma(\vartheta+1)} \right] \frac{1}{2} e^{\frac{x}{2}} - \left[(1-\vartheta) \frac{\vartheta t^{\vartheta-1}}{\Gamma(\vartheta)} + \frac{\vartheta^2 t^{2\vartheta-1}}{\Gamma(2\vartheta)} \right] \frac{1}{8} e^{\frac{x}{2}} \\ + \left[(1-\vartheta)^2 + 2(1-\vartheta) \frac{\vartheta t^{\vartheta}}{\Gamma(\vartheta+1)} + \frac{\vartheta^2 t^{2\vartheta}}{\Gamma(2\vartheta+1)} \right] \frac{1}{4} e^{\frac{x}{2}} \\ - \left[(1-\vartheta)^2 \vartheta \frac{t^{\vartheta-2}}{\Gamma(\vartheta-1)} + 2(1-\vartheta) \frac{\vartheta^2 t^{2\vartheta-2}}{\Gamma(2\vartheta-1)} + \frac{\vartheta^3 t^{3\vartheta-2}}{\Gamma(3\vartheta-1)} \right] \frac{1}{32} e^{\frac{x}{2}} \\ + \left[3(1-\vartheta)^2 \vartheta \frac{t^{\vartheta-1}}{\Gamma(\vartheta)} + 5(1-\vartheta) \frac{\vartheta^2 t^{2\vartheta-1}}{\Gamma(2\vartheta)} + 2 \frac{\vartheta^3 t^{3\vartheta-1}}{\Gamma(3\vartheta)} \right] \frac{1}{16} e^{\frac{x}{2}} \\ - \left[(1-\vartheta)^3 + 3(1-\vartheta)^2 \vartheta \frac{t^{\vartheta}}{\Gamma(\vartheta+1)} + 3(1-\vartheta) \frac{\vartheta^2 t^{2\vartheta}}{\Gamma(2\vartheta+1)} + \frac{\vartheta^3 t^{3\vartheta}}{\Gamma(3\vartheta+1)} \right] \frac{1}{8} e^{\frac{x}{2}} \dots$$

The exact solution at $\vartheta = 1$, $u(x, t) = e^{\frac{x}{2} - \frac{2}{3}t}$.

Example.2. Consider the fractional Fornberg-Whitham equation

$${}^{AB}\mathfrak{D}_t^\vartheta u - u_{xxt} + u_x + uu_x = 3u_x u_{xx} + uu_{xxx}, \quad (19)$$

With initial condition

$$u(x, 0) = (\cosh \frac{x}{4})^2,$$

Apply eq.(16), We achieve



$$u_0 = (\cosh \frac{x}{4})^2.$$

$$u_1 = - \left(1 - \vartheta + \vartheta \frac{t^{\vartheta}}{\Gamma(\vartheta+1)} \right) \left[\frac{11}{32} \sinh \frac{x}{2} \right].$$

$$u_2 = \frac{121}{512} \cosh \frac{x}{2} \left[(1-\vartheta)^2 + 2(1-\vartheta) \vartheta \frac{t^{\vartheta}}{\Gamma(\vartheta+1)} + \frac{t^{2\vartheta}}{\Gamma(2\vartheta+1)} \right] - \frac{11}{128} \sinh \frac{x}{2} \left[\vartheta(1-\vartheta) \frac{t^{\vartheta-1}}{\Gamma(\vartheta)} + \vartheta^2 \frac{t^{2\vartheta-1}}{\Gamma(2\vartheta)} \right].$$

$$u_3 = - \left[(1-\vartheta)^3 + 3(1-\vartheta)^2 \vartheta \frac{t^{\vartheta}}{\Gamma(\vartheta+1)} + 3(1-\vartheta) \frac{\vartheta^2 t^{2\vartheta}}{\Gamma(2\vartheta+1)} + \frac{\vartheta^3 t^{3\vartheta}}{\Gamma(3\vartheta+1)} \right] \frac{1331}{8192} \sinh \frac{x}{2} \\ + \frac{121}{2048} \cosh \frac{x}{2} \left[3(1-\vartheta)^2 \vartheta \frac{t^{\vartheta-1}}{\Gamma(\vartheta)} + 5(1-\vartheta) \frac{\vartheta^2 t^{2\vartheta-1}}{\Gamma(2\vartheta)} + 2 \frac{\vartheta^3 t^{3\vartheta-1}}{\Gamma(3\vartheta)} \right] \\ - \frac{11}{512} \sinh \frac{x}{2} \left[(1-\vartheta)^2 \vartheta \frac{t^{\vartheta-2}}{\Gamma(\vartheta-1)} + 2(1-\vartheta) \frac{\vartheta^2 t^{2\vartheta-2}}{\Gamma(2\vartheta-1)} + \frac{\vartheta^3 t^{3\vartheta-2}}{\Gamma(3\vartheta-1)} \right].$$

The approximate solution is



$$\begin{aligned}
 u(x, t) = & \frac{1}{2} + \frac{1}{2} \cosh \frac{x}{2} - \frac{11}{32} \sinh \frac{x}{2} \left[1 - \vartheta + \vartheta \frac{t^\vartheta}{\Gamma(\vartheta + 1)} \right] + \frac{121}{512} \cosh \frac{x}{2} \left[(1 - \vartheta)^2 + 2(1 - \vartheta)\vartheta \frac{t^\vartheta}{\Gamma(\vartheta + 1)} + \frac{t^{2\vartheta}}{\Gamma(2\vartheta + 1)} \right] \\
 & - \frac{11}{128} \sinh \frac{x}{2} \left[\vartheta(1 - \vartheta) \frac{t^{\vartheta-1}}{\Gamma(\vartheta)} + \vartheta^2 \frac{t^{2\vartheta-1}}{\Gamma(2\vartheta)} \right] \\
 & - \left[(1 - \vartheta)^3 + 3(1 - \vartheta)^2\vartheta \frac{t^\vartheta}{\Gamma(\vartheta + 1)} + 3(1 - \vartheta) \frac{\vartheta^2 t^{2\vartheta}}{\Gamma(2\vartheta + 1)} + \frac{\vartheta^3 t^{3\vartheta}}{\Gamma(3\vartheta + 1)} \right] \frac{1331}{8192} \sinh \frac{x}{2} \\
 & + \frac{121}{2048} \cosh \frac{x}{2} \left[3(1 - \vartheta)^2\vartheta \frac{t^{\vartheta-1}}{\Gamma(\vartheta)} + 5(1 - \vartheta) \frac{\vartheta^2 t^{2\vartheta-1}}{\Gamma(2\vartheta)} + 2 \frac{\vartheta^3 t^{3\vartheta-1}}{\Gamma(3\vartheta)} \right] \\
 & - \frac{11}{512} \sinh \frac{x}{2} \left[(1 - \vartheta)^2\vartheta \frac{t^{\vartheta-2}}{\Gamma(\vartheta - 1)} + 2(1 - \vartheta) \frac{\vartheta^2 t^{2\vartheta-2}}{\Gamma(2\vartheta - 1)} + \frac{\vartheta^3 t^{3\vartheta-2}}{\Gamma(3\vartheta - 1)} \right] \dots
 \end{aligned}$$

When selcting $\vartheta = 1$, we get exact solution $u(x, t) = \cosh^2 \left(\frac{x}{4} - \frac{11}{24}t \right)$.





Figure 1 Plot showing the solutions for various values of ϑ , alongside the exact solution at $\vartheta = 1$, for **Example 1**.






Table 1.absolute error for Example 1

x	$\vartheta = 1$	$\vartheta = 0.9$	$\vartheta = 0.7$	$\vartheta = 0.4$
0.5	0.00049496	0.031087	0.04746	0.061608
1	0.00063554	0.039917	0.060537	0.079106
1.5	0.00081606	0.051254	0.077731	0.10157
2	0.0010478	0.065811	0.099809	0.13042
2.5	0.0013454	0.084503	0.12816	0.16747
3	0.0017276	0.1085	0.16456	0.21503
3.5	0.0022183	0.13932	0.2113	0.27611
4	0.0028483	0.17889	0.27131	0.35453
4.5	0.0036573	0.2297	0.34837	0.45523
5	0.0046961	0.29495	0.44731	0.58452

Figure 3. Graph for different values of ϑ and the exact solution at $\vartheta = 1$ for Example 2.








Figure 4. Visualization of the solutions for varies values of ϑ , including the exact solution at $\vartheta = 1$, in Example 2.

Table 2. absolute error for **Example 2**.

x	$\vartheta = 1$	$\vartheta = 0.9$	$\vartheta = 0.8$	$\vartheta = 0.7$
5	0.005757	0.0039514	0.065314	0.17893
5.5	0.0070306	0.0070339	0.079532	0.22314
6	0.008746	0.010558	0.098746	0.28137
6.5	0.011011	0.014746	0.12416	0.35728
7	0.013967	0.01986	0.15738	0.45564
7.5	0.017801	0.026222	0.20049	0.58262
8	0.022754	0.034232	0.25619	0.74621
8.5	0.029136	0.044392	0.32799	0.95667
9	0.037348	0.057341	0.42039	1.2272
9.5	0.047907	0.073892	0.53921	1.5749

Conclusion

We investigated the Fornberg-Whitham equation using the fractional Atangana-Baleanu operator and employed the Yang Adomian Decomposition Method (YADM) to obtain an approximate solution. The results demonstrated the high accuracy and efficiency of the proposed method, as they closely aligned with the exact solution. This confirms the applicability of fractional operators in modeling complex nonlinear phenomena and highlights the effectiveness of YADM in solving fractional differential equations. Future work may explore the extension of this approach to other nonlinear fractional models and compare its performance with alternative numerical and analytical techniques.

References

1. R. Hilfer, Applications Of Fractional Calculus In Physics, vol. 35. World Scientific Publishing Company, 2000. doi.org/10.1142/3779
2. J. Sabatier, O. P. Agrawal, and J. A. T. Machado, Advances in Fractional Calculus: Theoretical Developments and Applications in Physics and Engineering, Springer Netherlands, 2007. DOI: [10.1007/978-1-4020-6042-7]
3. M. S. Al-luhaibi, "An analytical treatment to fractional Fornberg-Whitham equation" , Mathematical Sciences, vol. 11, no. 1, Mar. 2017. doi.org/10.1007/s40096-017-0208-2
4. J. C. Camacho, M. Rosa, M. L. Gandarias, and M. S. Bruzón, "Classical symmetries, travelling wave solutions and conservation laws of a generalized Fornberg-Whitham equation," J Comput Appl Math, vol. 318, pp. 149-155, 2017. doi.org/10.1016/j.cam.2016.08.036
5. G. B. Whitham, "Variational methods and applications to water waves," Proc R Soc Lond A Math Phys Sci, vol. 299, no. 1456, pp. 6-25, 1967. doi.org/10.1098/rspa.1967.0119

6. B. Fornberg and G. B. Whitham, "A numerical and theoretical study of certain nonlinear wave phenomena," *Philosophical Transactions of the Royal Society of London. Series A, Mathematical and Physical Sciences*, vol. 289, no. 1361, pp. 373–404, 1978. doi.org/10.1098/rsta.1978.0064
7. D. Kumar, J. Singh, and D. Baleanu, "A new analysis of the Fornberg-Whitham equation pertaining to a fractional derivative with Mittag-Leffler-type kernel," *The European Physical Journal Plus*, vol. 133, no. 2, pp. 1-10, 2018. doi.org/10.1140/epjp/i2018-11934-y
8. M. Hashemi, A. Badali, and P. Vafadar, "Group Invariant Solutions and Conservation Laws of the Fornberg-Whitham Equation," *Zeitschrift für Naturforschung A*, vol. 69, Oct. 2014. doi.org/10.5560/zna.2014-0044
9. J. Lu, "An analytical approach to the Fornberg-Whitham type equations by using the variational iteration method," *Computers & Mathematics with Applications*, vol. 61, no. 8, pp. 2010–2013, 2011. doi.org/10.1016/j.camwa.2010.08.067
10. M. Merdan, A. Gökdoğan, A. Yıldırım, and S. T. Mohyud-Din, "Numerical Simulation of Fractional Fornberg-Whitham Equation by Differential Transformation Method," *Abstract and Applied Analysis*, vol. 2012, no. 1, p. 965367, 2012. doi.org/10.1155/2012/965367
11. M. A. Ramadan and M. S. Al-luhaibi, "New iterative method for solving the fornberg-whitham equation and comparison with homotopy perturbation transform method," *British Journal of Mathematics & Computer Science*, vol. 4, no. 9, pp. 1213–1227, 2014. doi.org/10.9734/BJMCS/2014/10024
12. F. Abidi and K. Omrani, "Numerical solutions for the nonlinear fornberg-whitham equation by He's methods," *Int J Mod Phys B*, vol. 25, no. 32, pp. 4721–4732, 2011. doi.org/10.1142/S0217979211059663
13. K. Wang and S. Liu, "Application of new iterative transform method and modified fractional homotopy analysis transform method for fractional fornberg-whitham equation," *J. Nonlinear Sci. Appl.*, vol. 9, no. 5, pp. 2419–2433, 2016. doi.org/10.22436/jnsa.009.05.34
14. J.-H. He, "Approximate solution of nonlinear differential equations with convolution product nonlinearities," *Comput Methods Appl Mech Eng*, vol. 167, no. 1–2, pp. 69–73, 1998. doi.org/10.1016/S0045-7825(98)00109-1
15. J.-H. He, "Variational iteration method for autonomous ordinary differential systems," *Appl Math Comput*, vol. 114, no. 2–3, pp. 115–123, 2000. doi.org/10.1016/S0096-3003(99)00104-6
16. G.-C. Wu and D. Baleanu, "Variational iteration method for fractional calculus-a universal approach by Laplace transform," *Adv Differ Equ*, vol. 2013, pp. 1–9, 2013. doi.org/10.1186/1687-1847-2013-18
17. N. Anjum and J.-H. He, "Laplace transform: making the variational iteration method easier," *Appl Math Lett*, vol. 92, pp. 134–138, 2019. doi.org/10.1016/j.aml.2019.01.015
18. M. Dehghan, "Finite difference procedures for solving a problem arising in modeling and design of certain optoelectronic devices," *Math Comput Simul*, vol. 71, no. 1, pp. 16–30, 2006. doi.org/10.1016/j.matcom.2005.09.001.
19. M. A. Hussein, H. K. Jassim, A. K. Jassim, "An Innovative Iterative Approach to Solving Volterra Integral Equations of Second Kind," *Acta Polytechnica*, vol. 64, no. 2, pp. 87–102, 2024. doi.org/10.14311/AP.2024.64.0087
20. Alzaki L. K. and Jassim H. K., "The approximate analytical solutions of nonlinear fractional ordinary differential equations," *Int J Nonlinear Anal Appl*, vol. 12, no. 2, pp. 527–535, 2021. doi.org/10.22075/IJNAA.2021.5340
21. Jassim H. K. and Hussein M. A., "A novel formulation of the fractional derivative with the order $\alpha \geq 0$ and without the singular kernel," *Mathematics*, vol. 10, no. 21, pp. 4123, 2022. doi.org/10.3390/math10214123
22. Caputo M. and Fabrizio M., "A new definition of fractional derivative without singular kernel," *Prog Fract Differ Appl*, vol. 1, no. 2, pp. 73–85, 2015. doi.org/10.12785/pfda/010201
23. Losada J. and Nieto J. J., "Properties of the new fractional derivative without singular kernel," *Prog Fract Differ Appl*, vol. 1, no. 2, pp. 87–92, 2015. doi.org/10.12785/pfda/010202
24. Hussein M. A., "Approximate Methods For Solving Fractional Differential Equations", *Journal of Education for Pure Science-University of Thi-Qar*, vol. 12, no. 2, pp. 32–40, 2022.

25. Saeid A. R. and Alzaki L. K., "Analytical Solutions for the Nonlinear Homogeneous Fractional Biological Equation using a Local Fractional Operator" , J Edu Pure Sci-Univ of Thi-Qar, vol. 13, no. 3, pp. 1-17, 2023.
26. Tuan N. H., Mohammadi H., and Rezapour S., "A mathematical model for COVID-19 transmission by using the Caputo fractional derivative" , Chaos Solitons Fractals, vol. 140, pp. 110-107, 2020. doi.org/10.1016/j.chaos.2020.110107
27. Elzaki TM, Razaq A. Analytical solution of fractional differential equations using the natural variation iteration method. *J Educ Pure Sci.* 2025;15(1). doi:10.32792/jeps.v15i1.624
28. Jassim H. K. and Khafif S. A., "SVIM for solving Burger's and coupled Burger's equations of fractional order" , Prog Fract Differ Appl, vol. 7, no. 1, pp. 73-78, 2021. doi.org/10.18576/pfda/070106
29. Seewn NR, Yasser MT, Tajadodi H. An Efficient Approach for Nonlinear Fractional PDEs: Elzaki Homotopy Perturbation Method. *J Edu Pure Sci-Univ Thi-Qar.* 2025;15(1):89-99.
30. Yasser MT, Jassim HK. A new integral transform for solving integral and ordinary differential equations. *Math Comput Sci.* 2025.
31. Seewn NR, Yasser MT, Ziane D. An Analytical Approach to Nonlinear Fractional Differential Equations Using Daftardar-Jafari Method. *J Edu Pure Sci-Univ Thi-Qar.* 2025;15(1):62-73.
32. K. Diethelm, The Analysis of Fractional Differential Equations: An Application-Oriented Exposition Using Differential Operators of Caputo Type. in *Lecture Notes in Mathematics*. Springer Berlin Heidelberg, 2010. doi.org/10.1007/978-3-642-14574-2
33. M. I. Syam and M. Al-Refai, "Fractional differential equations with Atangana-Baleanu fractional derivative: analysis and applications," *Chaos, Solitons & Fractals: X*, vol. 2, pp. 100013, 2019. doi.org/10.1016/j.csfx.2019.100013
34. A. Atangana and D. Baleanu, "New fractional derivatives with nonlocal and non-singular kernel: theory and application to heat transfer model," *arXiv preprint arXiv:1602.03408*, 2016.
35. X. J. Yang, "New integral transforms for solving a steady heat transfer problem," *Thermal Science*, vol. 21, pp. 79-87, 2017. doi.org/10.2298/TSCI151215045Y
36. M. Naeem, H. Yasmin, R. Shah, N. A. Shah, and J. D. Chung, "A Comparative Study of Fractional Partial Differential Equations with the Help of Yang Transform," *Symmetry (Basel)*, vol. 15, no. 1, Jan. 2023. doi.org/10.3390/sym15010199
37. H. J. Haubold, A. M. Mathai, and R. K. Saxena, "Mittag-Leffler functions and their applications," *J Appl Math*, vol. 2011, no. 1, p. 298628, 2011. doi.org/10.1155/2011/298628
38. A. K. Shukla and J. C. Prajapati, "On a generalization of Mittag-Leffler function and its properties," *J Math Anal Appl*, vol. 336, no. 2, pp. 797-811, 2007. doi.org/10.1016/j.jmaa.2007.03.018