

## The relationship between prolonged QT interval and acute stroke in Tikrit teaching hospital

Hasan I. Sultan.

Dept. of Medicine, College of Medicine, Tikrit University.

### Abstract

The QT interval is closely related to ventricular action potential and is a good noninvasive measure of ventricular repolarization and myocardial homogeneity. Prolonged QT interval has been reported to be associated with higher mortality in stroke patients. The aim of this study is to assess whether QT interval is prolonged in patients with acute stroke, and whether this prolongation if present is related to the type, location, and size of stroke. The study included 60 patients with acute stroke admitted to Tikrit teaching hospital from July 2011 to February 2012 and 12 leads ECGs were recorded at the time of admission, to measure the corrected QTc interval (QTc manual and QTc automated). A 30 age matched individuals (control group) were included for comparison. The QTc interval measurement was significantly longer in patients with acute stroke than control group. The QTc interval was significantly higher with big infarct or large hemorrhage than small lesions. There was no significant difference between right and left sided stroke regarding QTc interval values. It was concluded that QT interval was prolonged in acute stroke and this prolongation was related to the size but not the type or location of stroke.

### Introduction

Acute stroke is defined as rapidly onset of clinical signs of focal brain dysfunction lasting more than 24 hours or leading to death with no apparent cause(1,2).

There are number of electrocardiographic (ECG) changes that well known to appear during acute stroke like ST and T wave changes, which seem to be due to involvement of autonomic cardiac control centers. QT interval is closely related to ventricular action potential and is a good noninvasive measure of ventricular repolarization and myocardial homogeneity(3,4).

The rate related, corrected QT (QTc) interval

can be calculated as  $\frac{QT}{\sqrt{RR}}$  (Bazzett's formula) and normally is  $\leq 440$  milliseconds(4). Prolonged QT interval is a predictor of serious

ventricular tachyarrhythmias and sudden cardiac death in patients with ischemic heart disease(5,6), but some new studies showed that prolonged QT interval has been associated with higher mortality in stroke patients(7,8,9). The prognostic importance of QT interval in acute stroke is still underestimated in our locality. The aim of this study is to assess whether QT interval is prolonged in patients with acute stroke and whether this prolongation if present is related to the type, location and size of stroke.

### Patients and methods:

**Patients group:** 60 patients, who were admitted to Tikrit teaching hospital within 24 hours of onset of stroke during the period from July/2011 to February/2012.

**Control group:** 30 age matched individuals who were free from cardiovascular or



neurological disease, were included for comparison.

Exclusion criteria: patients presenting with the following conditions were excluded from the study (10,11):

#### **Transient ischemic attack.**

Ischemic or valvular heart disease, heart failure, arrhythmias or cardiomyopathy.

Patient taking these medications: digoxin, antiarrhythmic drugs, phenothiazine, tricyclic antidepressant drugs and macrolides antibiotics.

ECG reveals a bundle branch block or left ventricular hypertrophy.

Abnormal serum potassium or calcium concentration.

ECG recording: A 12 leads ECGs were recorded to the patients at the time of admission to hospital by using ECG machine (FX-7102 version 2, Fukuda Denshi, Japan) at a speed of 25 mm/s and amplitude of 10 mm/mV. The QT interval was measured from the onset of QRS deflection to the end of the T wave, the point of the return of T wave to the isoelectrical line, or the nadir between T and U waves(12,13). QT interval was measured manually in milliseconds (ms). Three successive QT intervals were measured in each lead and the mean was accepted as the measurement for that lead. This QT interval was corrected (QTc) for heart rate by using Bazett's formula (14)(QTc manual). Automated QT and corrected automated QT (QTc automated) intervals were obtained from ECG recorded data paper.

Neurological assessment: A complete neurological examination and brain imaging by CT scan or MRI were performed to all patients at the time of admission. In this study patients with ischemic stroke classified according to Oxford community stroke project classification as following (15):

Total anterior circulation infarct (TACI): A triad of hemiparesis (or hemisensory loss), dysphasia (or other higher cortical dysfunction) and homonymous hemianopia.

Partial anterior circulation infarct (PACI): Present with only two of the features of TACI or isolated dysphasia.

Posterior circulation infarct (PCI): Patients with brain stem or cerebellar signs and/or isolated homonymous hemianopia.

Lacunar infarction did not include in this study. Patients with hemorrhagic stroke were classified according to Afsar *et al* and Chung *et al* (11,16) into:

Large hemorrhage: > 33 mm in diameter, with or without ventricular extension or subarachnoid hemorrhage.

Small hemorrhage: ≤ 33 mm in diameter.

**Statistical analysis:** analysis was performed using SPSS version 12.0 software program. The data were reported as mean ± SD and frequencies expressed as percent. The unpaired and paired t tests were used to compare two groups of variables. ANOV test was used to compare QTc interval values among subtypes of ischemic stroke. The probability value( $P < 0.05$ ) was considered to be statistically significant.

## **Results**

There were 60 patients with acute stroke (patients group) with mean age of  $63.6 \pm 10.2$  years and 28 (46.7%) patients were males. There were 47(78.3%) patients had ischemic and 13(21.7%) patients had hemorrhagic stroke.

The control group included 30 individuals with mean age of  $60.5 \pm 9.7$  and 12 (40%) of them were males. The QTc interval measurements (QTc manual and QTc automated) were significantly longer in patients group than



control group (Table 1). However there was no significant difference between manual and automated assessment of QTc interval in both groups (Table 2).

The QTc interval measurements were found to be significantly higher in TACI and PACI than PCI in comparison subtypes of ischemic stroke (Table 3), and being significantly greater with large rather than small hemorrhagic stroke (Table 4). There was no significant difference between the location of stroke (right or left side) diagnosed by CT scan or MRI and different values of QTc interval (Table 5).

## Discussion

The present study revealed a high agreement between machine and manual QTc interval measurement, therefore Bazett's formula is still a useful method for evaluation of this interval(13,14,17). The QTc interval of patients with acute stroke was significantly higher than control group, considering exclusion of concomitant cardiac disease, that prolongation may reflect the acute disturbances of central nervous system function (11,17), which may leads to increase catecholamines secretion and sympathetic over activation (18,19). However, the QTc interval prolongation is also related to the size of lesion in the brain, being significantly longer with big infarct or large hemorrhage than small lesions, which might be due to wide involvement of cardiovascular autonomic control centers in the brain(3).

Other previous studies showed a greater QTc interval prolongation with insular involvement. Insula is the cortex of the brain that lies at the base of Sylvian fissure, it affected by proximal middle cerebral artery occlusion and associated with greater stroke severity(20,21). Moreover, the present study revealed no significant correlation between the location of stroke (right or left side) and QTc interval measurements, which seems to be due to that cardiovascular autonomic innervation

come from the two cerebral hemispheres(3,19,22).

It was concluded that QT interval was prolonged in acute stroke and this prolongation was related to the size but not to the type and location of stroke. Therefore, more studies and follow up are recommended to explore if the prolonged QT interval worsen the prognosis of stroke patients.

## References

- 1-Allen CM, Leuk CJ, Dennis M. Neurological disease: Cerebrovascular disease. In: Colledge NR, Walker BR, Ralston SH. Davidson's principles and practice of medicine. 21st ed. Edinburgh: Churchill livingstone; 2010. 1180-1190.
- 2-Familoni OB, Odusan O, Ogun SA. The pattern and prognostic features of QT intervals and dispersion in patients with acute ischemic stroke. J Nati Med Assoc 2006;98:1758-1762.
- 3-Alabd AA, Fouad A, Abdel-Nasser R, Nammias W. QT interval dispersion pattern in patients with acute ischemic stroke: Does the site of infarction matter?. Int J Angiol 2009; 18:177-181.
- 4-Goldberger AL. Electrocardiography. In: Kasper DL, Fauci AS, Longo D, Braunwald E, Hauser S, Jameson JL. Harrison's principles of internal medicine. 16th ed. New York: McGraw-Hill; 2005. 1311-1313.
- 5-Dekker JM, Crow RS, Hannan PJ, Schouten EG, Folsom AR. Heart rate-corrected QT interval prolongation predicts risk of coronary heart disease in black and white middle-aged men and women: The ARIC study. J Am CollCardiol 2004;43:565-571.
- 6-Dekker JM, Schouten EG, Klootwijk P, Pool J, Kromhout D. Association between QT interval and coronary heart disease in middle-aged and elderly men. The Zutphen study. Circulation 1994; 90:779-785.
- 7-Chao CC, Wang TL, Chong CF, Lin Y, Chen CC, Tang C, et al. Prognostic value of QT parameters in patients with acute hemorrhagic stroke: A prospective evaluation



with respect to mortality and post-hospitalization bed confinement. *J Chin Med Assoc.* 2009;72: 124–132.

8-Cardoso CR, Salles GF, Deccache W. QTc interval prolongation is a predictor of future strokes in patients with type 2 diabetes mellitus. *Stroke* 2003;34:2187-2194.

9-Maebuchi D, Arima H, Doi Y, Ninomiya T, Yonemoto K, Tanizaki Y, et al. QT interval prolongation and the risks of stroke and coronary heart disease in a general Japanese population: The Hisayama study. *Hypertens Res* 2010;33:916-921.

10-Hamdan MH. Cardiac arrhythmias: Long QT syndrome. In: Andreoli TE, Benjamin IJ, Griggs RC, Wing EJ. *Andreoli and Carpenter's Cecil essentials of medicine*. 8th ed. Philadelphia: Saunders Elsevier; 2010. 143-144.

11-Afsar N, Fak AS, Metzger JT, Melle GV, Kappenberger L, Bogousslavsky J. Acute stroke increases QT dispersion in patients without known cardiac diseases. *Arch Neurol* 2003;60:346-350.

12-Goldenberg I, Moss AJ, Zareba W. QT interval: How to measure it and what is normal. *J CardiovascElectrophysiol* 2006;17:333-336.

13-Fosser C, Duczynski G, Agin M, Wicker P, Darpo B. Comparison of manual and automated measurements of the QT interval in healthy volunteers: an analysis of five thorough QT studies. *ClinPharmacolTher* 2009;86:503-506.

14-Kasamaki Y, Ozawa Y, Ohta M, Sezai A, Yamaki T, Kaneko M, et al. Automated versus manual measurement of the QT interval and corrected QT interval. *Annals of Noninvasive Electrocardiology* 2011;16:156-164.

15-Mead GE, Lewis SC, Wardlaw JM, Dennis MS, Warlow CP. How well does the

Oxfordshire Community Stroke Project classification predict the site and size of the infarct on brain imaging?. *J Neurol Neurosurg Psychiatry* 2000;68:558–562.

16-Chung CS, Caplan LR, Han W, Pessin MS, Lee KH, Kim JM. Thalamic haemorrhage. *Brain* 1996;119:1873-1886.

17-Mulcahy J, Johnson P, James M. Electrocardiogram QT interval increases in acute stroke. *Cerebrovasc Dis* 2010;29:178-180.

18-Hachinski VC, Smith KE, Silver MD, Gibson CJ, Ciriello J. Acute myocardial and plasma catecholamine changes in experimental stroke. *Stroke* 1986;17:387-390.

19-Tokgozoglu SL, Batur MK, Top uoglu MA, Saribas O, Kes S, Oto A. Effects of stroke localization on cardiac autonomic balance and sudden death. *Stroke* 1999;30:1307-1311.

20-Fink JN, Selim MH, Kumar S, Voetsch B, Fong WC, Caplan LR. Insular cortex infarction in acute middle cerebral artery territory stroke predictor of stroke severity and vascular lesion. *Arch Neurol* 2005;62:1081-1085.

21-Christensen H, Boysen G, Christensen AF, Johannesen HH. Insular lesions, ECG abnormalities, and outcome in acute stroke. *J NeurolNeurosurg Psychiatry* 2005;76:269-271.

22-Fink JN, Frampton CM, Lyden P, Lees KR. Does hemispheric lateralization influence functional and cardiovascular outcomes after stroke? : An analysis of placebo-treated patients from prospective acute stroke trials. *Stroke* 2008;39:3335-3340.



**Table 1.**The basic characteristics of patients studied

Variables		Patients group (n:60)	Control group (n:30)	P value
Age		63.6 ± 10.2	60.5 ± 9.7	>0.05
Sex		M 28(46.7%) : F 32(53.3%)	M 12(40%) : F 18(60%)	>0.05
Ischemic stroke: n:47(78.3%)	TACI	16(34%)	-----	-----
	PACI	28(59.6%)	-----	-----
	PCI	3(6.4%)	-----	-----
Hemorrhagicstroke: n:13(21.7%)	Large	9(69.2%)	-----	-----
	Small	4(30.8%)	-----	-----
QTc manual(ms)		467.2 ± 47.8	426.03 ± 28.5	<0.05
QTc automated (ms)		462.5 ± 37.9	422.2 ± 29.7	<0.05
Fasting blood sugar (mmol/L)		8.4 ± 3.1	6.1 ± 2.3	>0.05
Blood urea (mmol/L)		7.5 ± 2.4	6.2 ± 3.1	>0.05
Serum creatinine (μmol/L)		88.5 ± 17.9	75.3 ± 3.2	>0.05
Serum sodium (mmol/L)		140.1 ± 7.6	142.3 ± 4.3	>0.05
Serum potassium (mmol/L)		4.2 ± 0.4	4.7 ± 0.6	>0.05
Serum calcium (mmol/L)		2.2 ± 0.1	2.3 ± 0.2	>0.05

**Table 2.** The difference between manual and automated QTc interval measurements in patients studied

Groups	QTc interval		P value
	QTc manual (ms)	QTc automated (ms)	
Patients group (n: 60)	467.2 ± 47.8	462.5 ± 37.9	>0.05
Control group (n: 30)	426.03 ± 28.5	422.2 ± 29.7	>0.05

**Table 3.**QT interval measurements in subtypes of ischemic stroke

Ischemic stroke(47)				P value
	TACI n: 16(34%)	PACI n: 28 (59.6%)	PCI n: 3 (6.4%)	
QTc interval				
QTc manual (ms)	474.25 ± 36.6	471.107 ± 46.1	415.67 ± 33.4	<0.05
QTc automated (ms)	469 ± 31.6	464.32 ± 32.7	405.66 ± 11.6	<0.05



**Table 4.** QTc interval measurements in subtypes of hemorrhagic stroke

Hemorrha. stroke(13)	Large	Small	P value
QTc interval	n: 9 (69.2%)	n: 4 (30.8%)	
QTc manual (ms)	498.78 ± 46.1	445.25 ± 30.7	<0.05
QTc automated (ms)	492.67 ± 48.4	431.5 ± 38.9	<0.05

**Table 5.** QT interval measurements according to the location of stroke by CT or MRI

Location		Right side	Left side	P value
QTc interval (ms)				
Ischemic stroke: 47 (78.3%)	QTc manual	n: 23 (48.9%) 468.82±51.6	n: 24 (51.1%) 465.08±41.2	>0.05
	QTc automated	457.82±36.8	460.317±33.7	>0.05
Hemorrhagic stroke: 13 (21.7%)	QTc manual	n: 6 (46.2%) 493.33±39.1	n: 7 (53.8%) 464.428±35.0	>0.05
	QTc automated	488±59.6	454.714±53.3	>0.05

### العلاقة بين استطالة فاصل QT والسكتة الدماغية الحادة في مستشفى تكريت التعليمي

حسن اسماعيل سلطان، قسم الطب الباطني، كلية الطب، جامعة تكريت

#### خلاصة

فاصل QT ذو علاقة وثيقة مع عمل الجهد الكهربائي البطيني، وهو مقياس جيد غير اجتياحي لعودة الاستقطاب البطيني وتجانس عضلة القلب. وقد سجل ترافق استطالة فاصل QT مع ارتفاع معدل الوفيات في مرضى السكتة الدماغية. ان الهدف من هذه الدراسة هو تقييم ما اذا كان هنالك استطالة في فاصل QT لدى المرضى المصابين بالسكتة الدماغية الحادة، وان كانت هذه الاستطالة موجودة هل لها ارتباط بنوع وموقع وحجم السكتة الدماغية. تضمنت هذه الدراسة ستين مريض مصابين بالسكتة الدماغية الحادة ادخلوا مستشفى تكريت التعليمي للفترة من تموز ٢٠١١ الى شباط ٢٠١٢ وسجل لهم تخطيط القلب الكهربائي في وقت الدخول وذلك لقياس فاصل QT المصحح (فاصل QT المصحح اليدوي وفاصل QT المصحح الآلي)، شملت الدراسة أيضا ثلاثين شخص (مجموعة مراقبة) وهي مطابقة للعمر من أجل المقارنة. في هذه الدراسة كانت قياسات فاصل QT المصحح أطول بشكل ملحوظ في المرضى الذين يعانون من السكتة الدماغية الحادة مقارنة بمجموعة المراقبة، وكان فاصل QT المصحح أعلى بكثير مع الاحتشاء الدماغية الضخم أو النزف الدماغية الكبير مقارنة بالآفات الصغيرة، وفي ما يتعلق بقياسات فاصل QT المصحح لم يكن هناك فرق ذو أهمية عند مقارنة السكتة الدماغية في الجانب الأيمن مع الجانب الأيسر. استنتج من هذه الدراسة انه توجد هنالك استطالة في فاصل QT لدى المرضى المصابين بالسكتة الدماغية الحادة، وهذه الاستطالة لها علاقة بحجم لا بنوع او موقع السكتة الدماغية.