



P-ISSN: 2788-9890 E-ISSN: 2788-9904

NTU Journal of Agricultural and Veterinary Sciences

Available online at: <https://journals.ntu.edu.iq/index.php/NTU-JAVS/index>



Use of Azolla the in Presence of Fiber-Degrading Enzyme Additives and its Effect onthe Productive Performance in Broiler

1st Baker Hani Taha , 2nd Anwar Mohammed Younis AL-Hamed , 3rd Abd Al-Bar Al-Farha

1. Department of Anesthesia Techniques, Medical Technical Institute/Mosul

2. Department of Animal Production, College of Agriculture and Forestry, University of Mosul

3. Infectious Diseases Department , Technical Research Center, Northern Technical University, Mosul, 41001, Iraq

Article Informations

Received: 23-07- 2024,

Accepted: 03-12-2024,

Published online: 28-09-2025

Corresponding author:

Name: Anwar Mohammed Younis
AL-Hamed

Affiliation : Department of Animal
Production, College of Agriculture
and Forestry, University of Mosul
Email:

dr.anwaralhamed@uomosul.edu.iq

Key Words:

Azolla ,
Enzyme
Fiber-Degrading
Broiler ,
Performance

ABSTRACT

In this study, we used unsexed broiler chicks of the hybrid (Ross 308), raised in animal production fields/College of Agriculture and Forestry for the period from 7/2/2024 to 19/ 3 /2024. The chicks were subjected to the treatments under study from the first day of the experiment. The experiment included five treatments: T1 : Standard ration (without Azolla and enzyme).T2: ration contained Azolla 7% and without enzyme .T3: ration contained Azolla 7% + 0.30% enzyme.T4: ration contained Azolla 7% + 0.35% enzyme.T5: ration contained Azolla 7% + 0.40% enzyme. Measurements of production characteristics were taken weekly and statistical analysis shows that there was a significant increase in live body weight, weight gain, an improvement in feed conversion ratio, a decrease in the feed consumed In enzyme addition transactions, and no significant effect of the treatments in the relative growth rate.

This study aimed to prepare a feeder containing Azolla with the addition of enzymes as a technical management method that contributes to increasing the decomposition of fibers to increase the rate of utilization as a non-traditional protein source for locally available fodder



Introduction

One of the most important elements of profitable continuous production is the availability of feed components that are included in the composition of the feed, which has become a competitor to traditional human foodstuffs [1], and their prices are considered part of the production cost. Recently, non-traditional feeds have been used to provide locally available protein sources, including Azolla, which was used in feeding large animals and animals with simple stomachs [3], [4], [5], [6], [7]. In addition to being considered a protein supplement in many studies as a source of the amino acids methionine, lysine, and histidine and containing vitamins A and B12 [8], [9], [10], Azolla is also characterized by its high content of phenols, flavonoids, carotenoids, and probiotics, and therefore it is considered an antioxidant and antibacterial. [11] Azolla contains a percentage of fibers, which may be a cause of nutritional problems because it is considered an anti-nutrient that affects the digestion of feed materials. This study came with the addition of enzymes as a nutritional supplement to the feed to contribute to increasing the rate of utilization and simplifying its content of fibers and nutrients and increasing its digestion coefficient and absorption efficiency, which is reflected in the benefit of food, as enzymes are

used to decompose the fibers that make up the cell wall [12], which are the digestive enzymes for non-starchy polysaccharides such as beta-glucanase, cellulase, mannanase, and xylanase, which are the most widely used in the field of poultry nutrition [13]. They are of great importance in treating the viscosity resulting from non-starchy polysaccharides inside

the digestive tract increasing their digestion coefficient and enhancing the work of internal enzymes, causing an increase in the availability of decomposed nutrients, which increases their absorption process. [31],[32] and thus increasing the rate of utilization of released nutrients, improving the feed conversion ratio, and reducing feeding costs [34].

Materials And Methods

The research was carried out in the poultry field of the Department of Animal Production at the College of Agriculture and Forestry / University of Mosul, the experiment was 42 days from 7/2/2024 to 19/3/2024, and the research aimed to study the effect of preparing the feed containing Azolla with enzymes. Fiber hydrolyzate is a nutritional management tool to increase the utilization rate of nutrients when used as a non-traditional locally

available source and to know its effect on productive traits. The chickens used are a hybrid type (Ross 308) and the total number of chickens is (390) with a starting weight of 42.11 grams, The chicks were treated as follows: T1 (control): feed without additives, T2: feed containing 7% Azolla, T3: adding 0.30% enzyme to the feed containing Azolla, T4: adding 0.35% enzyme to the feed containing Azolla, T5: adding 0.40% enzyme to the feed containing Azolla. The chicks were subjected to the experimental treatments from the first day of rearing and were randomly distributed to the experiment replicates (15). Feed and water were continuously available with an incubation temperature of 35°C and continuous monitoring of the chicks' movement and vitality.

Table 1. shows the percentages of feed components.

Feed ingredients %	Ratios used %			
	Starter ration		Finisher ration	
Dried Azolla	0.0	7.0	0.0	7.0
Yellow Corn	57.72	53.1	60.91	55.12
Soybean Meal	38.85	36.34	33.69	31.4
Limestone	2.38	1.83	1.85	1.83
Table Salt	0.25	0.25	0.25	0.25
Vegetable Oil	-	0.99	2.44	3.84
Dicalcium Phosphate	0.8	0.49	0.86	0.56
Total	100	100	100	100
calculated chemical analysis				
The energy represented ilo/kg.	2800	2800	3000	3000
Protein %	22	22	20	20

* Calculated according to [14]

The type of Azolla in this study is *Azolla pinnata*. The Azolla was green and dried in the shade with stirring, then added to the feed when grinding the feed components to ensure its mixing with the feed. Table (2) shows the chemical composition of the Azolla used in the study.

Table 2. shows the laboratory chemical analysis of dried Azolla .

the components	percentage
Protein	29.8
Ash	15.9
Fiber	19.7
Ether Extract	1.7
Moisture	10.15

The enzymes used in the study were a powder of enzymes from a company specialized in veterinary treatments. The composition used is shown in Table (3).

Table 3. shows the enzymes used in the study.

Enzymes	Amount in powder U/g
beta-xylanase	35000
alpha-amylase	400
beta-glucanase	2350
Bacilolysin	1700

The studied production traits were measured, namely live body weight, as the chicks were weighed weekly using a scale with a sensitivity of (5 g). The weight gain was calculated, the weekly growth rate, feed consumption, feed conversion ratio, production coefficient, and production index. The Completely Randomized Design (C.R.D) was used in data analysis using SAS (2001) test significance according to [16] New Multiple tests to find significant differences between the treatments at the probabil ity level ($p \leq 0.05$).

Results and discussion

Live body weight

Table (4) shows that average live body weight of broilers in the first and second weeks did not differ significantly ($P \geq 0.05$) between treatments, while a significant increase was observed in the third week of the fifth treatment, where it reached (785.60) g compared to T I, T2 and third treatments and its broiler feed and differed with [9] where it was found that there was a decrease in the body reached (712.43, 717.14 and 722.47) g, respectively, while it did not differ significantly from the fourth treatment, which reached (754.3) g. It was also observed that the average body weight increased significantly in the fourth week in the T5 compared to the T1 and T2, and reached (1431.83, 1304.33, 1321.31) g, respectively. Also, in the fifth week, the fifth treatment outperformed the control and second treatments, and reached (2041.23, 1894.67, 1911.11) g, respectively, while we find that there is a significant increase in the sixth week for all treatments of added the enzyme to the feed containing Azolla (the third, fourth and fifth treatments) and also the increased significantly of the second treatment over the control and reached respectively (2548.33, 2552.0, 2,630.0, 2437.0) g compared to (2429.70) g. This effect did not appear in the first and second weeks, but it appeared in the last weeks.

The addition rate of 0.40% was able to contribute to raising in live body weight and increasing the rate of feed utilization (food authorization factor, Table 8) despite the decrease in the amount consumed in the same treatments and the same weeks (Table 7). The average body weight was enhanced in the sixth week by the cumulative effect of the enzyme's work as a result of improving the internal environment of the birds' intestines, which improved the rate of Azolla digestion and increased the availability of decomposed nutrients, which increased their absorption rate. The added enzymes

also contributed to enhancing the work of internal enzymes as nutritional supplements, which led to improved nutrient utilization [17],[18], which was reflected in body weight.

These results agreed with the researcher [19], [20] as they found a significant increase in body weight when adding Azolla to the feed, while these results differed with the researcher [2], [23]

, [24] as they showed no effect of adding azolla to the feed of broilers. Regarding the use of enzymes, these results agreed with researchers [25], [26] stated that adding the enzyme mixture led to an increase in the final body weight of broilers compared to the control treatment, while they differed with researchers [27] who showed that when adding the enzyme, the body weight decreased significantly compared to the control. Researchers [28], [29], [30] did not find differences in adding the enzyme to broiler feed and differed with [9] where it was found that there was a decrease in the body weight of broilers at the age of 42 days by adding the enzyme to feed containing Azolla.

Weight Gain Rate

Table 5 shows no significant differences in weight gain during the first, second, and fifth weeks of age, while it was found that in the third week, there was a significant increase in the fourth and fifth treatments compared to the control treatment and amounted to (394.3, 422.23, 372.26) gm respectively, while the second and third treatments did not differ significantly with the previously mentioned studied treatments and amounted to (378.94, 371.17) gm respectively, and also in the fourth week we find that the fifth treatment increased significantly in the first, second, third and fourth treatments as it amounted to (646.23, 591.9, 604.17, 597.43, 594.87) gm respectively. It noted that in the sixth week, there was a significant increase in the enzyme addition treatments at different rates (0.30, 0.35, 0.40)% to the feed containing Azolla compared to the treatment (first and second) and reached (602.66, 597.4, 589.37, 535.7, 525.89) gm respectively. As for the weight gain during the total period of rearing (1-6) weeks, a significant increase was found in the treatments to which the enzyme was added (third, fourth, and fifth) compared to the treatments (first and second) and reached (2506.93, 2505.0, 2589.0, 2388.37, 2394.9) gm respectively.

These significant differences in this trait are a result of the significant differences in the body weight table (Table 4) and confirm the fact that the enzyme mechanism works in benefiting from Azolla by increasing digestion and absorption processes as

a result of reducing the viscosity of the food and enhancing the work of internal enzymes, which was reflected in the performance of the birds [31], [32]. These results were consistent with [2], [23] as he did not find a significant effect of adding Azolla on weight gain. They differed from [33], [20], [24] when they added Azolla to the feed, there was a significant increase in weight gain rate of broiler chickens at the age of 42 days. Regarding the addition of enzymes, these results agreed with [35] noted that when added enzyme mixture there was an insignificant increase in the rate of weight gain during 28 days of rearing, and differed with [27], [28], [36]. [9] stated that when adding the enzyme to the azolla feed there was a decrease in the average final weight gain.

Growth Rate

Table 6 shows no significant differences between the 1, 2, 3, 4, and fifth treatments in all weeks of rearing as well as the total rearing period from (1-6) weeks. These results agreed with the researcher [24] while these results did not agree with the researcher [37]. The reason may be due to the protein obtained from Azolla being able to give the same production performance as the feed protein in terms of growth indicators (Table 4), (Table 5), (Table 6) and economic indicators (production index and coefficient).

Feed Consumed

Table (7) shows that there are no significant differences ($P \geq 0.05$) in the first, second, and fourth weeks between the first, second, third, fourth, and fifth treatments. As for the third week, we find that the enzyme addition treatments decreased significantly compared to the control treatment and amounted to (64.519, 552.02, 570.01) g for the third, fourth, and fifth treatments, respectively, compared to the control treatment (573.28) g. In the fifth week, the second, third, fourth, and fifth treatments decreased significantly compared to the control treatment and amounted to (1013.75, 1021.19, 1047.48, 1073.44, 1166.50) g/bird. It was found that the amount of feed consumed during the sixth week. In the second treatment, it decreased compared to the control and the third treatment, and they reached (1009.71, and 1096.74) g, respectively, compared to the control treatment (1187.54) g. As a result, we find that the total amount of feed consumed for the period from (1-6) weeks decreased significantly in the second, third, fourth, and fifth treatments compared to the control treatment, as it reached (3568.4, 3974.03, 3901.24, 3968.33, 4556.36) g/bird for the total rearing period, respectively.

The reason may be that the birds in the Azolla-containing treatments consumed less feed than the control, as they aged, because the dried Azolla

absorbed water, which increased the volume of food mass in the intestines while reducing the amount of feed consumed. This gave enough time for the food to pass through the intestines. With the presence of Azolla and the enzyme, the activity of beneficial bacteria increased, which enhanced the digestion and absorption processes, and the nutrients became more abundant, which increased the rate of utilization of the feed (Food Conversion Coefficient Table 7).

These results agreed with [38], [20] where a decrease in the feed consumed was observed using Azolla, and differed with [21], [22], [39] and [23]. Regarding the addition of the enzyme, these results agreed with [27], as the amount of feed consumed decreased significantly by adding the enzyme to the feed. And these results differed from [29], [26], and [30], who showed that when adding the enzyme,

Feed conversion ratio

Table (8) showed that there were no significant differences in the first and second weeks between the first, second, third, fourth, and fifth treatments, while we find that the third, fourth, and fifth treatments decreased significantly compared to the first and second treatments during the third week (1.40, 1.41, 1.39, 1.54, 1.56, 1.54, 1.56) g feed/g weight gain, and in the fourth week the conversion factor decreased significantly in the treatments (second, third, fourth, here were no significant differences in the feed conversion factor by adding the enzyme to the feed. They differed with [9], [40] that there was no significant improvement when adding the enzyme to the Azolla feed, the best feed conversion factor when raising broilers for a total period of 42 days. fifth) compared to the control treatment, and in the sixth week, the treatments (third, fourth, fifth) decreased significantly compared to the control treatment. in the sixth week and the conversion factor (1-6) weeks, all addition treatments (second, third, fourth, and fifth) decreased compared to the first control treatment, so the values reached (1.49, 1.47, 1.51, 1.51, 1.66) g feed/g weight gain, respectively.

This may be because the enzyme contributed to the digestion and decomposition of feed materials, which led to an increase in the abundance of nutrients, which was reflected in the absorption and improved utilization of nutrients as a result of reducing the viscosity of the feed [31] and [32] and enhancing the work of internal enzymes [17], [18], which was reflected in the performance of the birds. These results agreed with the researchers [41], [26] who stated that when enzymes were added, the feed conversion factor decreased significantly. These

results did not agree with the researcher [42],[36] as the amount of feed consumed, and differed from [9], that when adding Azolla with the enzyme, there was an increase in the amount of feed consumed.

Economic Implications

Table (9) showed that the production coefficient was high in the second, third, fourth, and fifth treatments compared to the control treatment for the periods (1-5) and (1-6) weeks, reaching (32.41, 38.63, 35.34, 35.34, 29.88) g, (27.39, 33.11, 32.29, 31.18, 24.16) respectively. The production index for these treatments also increased for the period (1-5) weeks, reaching (300.02, 343.15, 328.50, 339.07) for the second, third, fourth, and fifth treatments compared to treatment (273.95) respectively. It was also noted that the enzyme addition treatments raised the production index for the period (1-6) weeks compared to the first treatment. It reached (400.05, 394.13, 397.67) respectively compared to (313.3) and this was a result of raising the performance of the weight gain trait and the average live body weight. This confirms the fact that the rate of utilization of food-degrading enzymes has contributed to raising the production performance, which raised the economic efficiency.

Conclusion

Adding enzymes to the feed containing Azolla contributed to the decomposition of fibers and increased the abundance of nutrients in the digestive tract, which increased the rate of digestion and absorption, which was reflected in the production performance and reduced the cost of production.

Conflict of interest

The researchers pledge that there is no conflict in writing the research with the interests of publishing this article.

Acknowledgments. Thanks to the Department of Animal Production / Faculty of Agriculture and Forestry for providing the field and its supplies to conduct this study.

References

[1] Thirumalaisamy, G., Muralidharan, J., Senthilkumar, S., Hema Sayee, R., & Priyadharsini, M. (2016). Cost-effective. feeding of poultry. *International Journal of Science, Environment and Technology*, 5(6),3997-4005. <https://2u.pw/Drfb4x55>

[2] Abdelatty, A. M., Mandouh, M. I., Al-Mokaddem, A. K., Mansour, H. A., Khalil, H. M. A., Elolimy, A. A., & Bionaz, M. (2020). Influence of level of inclusion

of Azolla leaf meal on growth performance, meat quality, and skeletal muscle p70S6 kinase α abundance in broiler chickens. *animal*, 14(11), 2423-2432.

<https://doi: 10.1017/S1751731120001421>

- [3] de Almeida, A. M., & Zuber, U. (2010). The effect of the Naked Neck genotype (Nana), feeding, and outdoor rearing on growth and carcass characteristics of free-range broilers in a hot climate. *Tropical Animal Health and Production*, 42,99-107. <https://doi: 10.1007/s11250-009-9391-y>
- [4] Namra, M. M. M., Hataba, N. A., & Wahed, H. A. (2010). The productive performance of growing fayoumi chicks fed restricted diets supplemented with free fresh azolla. *Egyptian Poultry Science Journal*, 30(3),747-762. <https://2u.pw/Qbk8tRdG>
- [5] Chatterjee, A., Arvind Raj, N., Ghosh, M. K., Roy, P. K., & Dutta, T. K. (2012). Azolla Meal: A potential feed supplement for calves. *NDRI News*, 17(2), 4.
- [6] Islam, M. A., & Nishibori, M. (2017). Use of multivitamins, acidifier, and Azolla in the diet of broiler chickens. *Asian-Australasian journal of animal sciences*, 30(5), 683. <https://doi.org/10.5713%2Fajas.16.0395>
- [7] Al-Hamed, A. M., & Al-Husseiny, N. A. (2023, December). Use of Azolla Plant in Broiler Diets and its Effect on Productive Performance. In *IOP Conference Series: Earth and Environmental Science* (Vol. 1252, No. 1, p. 012134). IOP Publishing. <https://doi:10.1088/1755-1315/1252/1/012134>
- [8] Katole, S. B., Lende, S. R., & Patil, S. S. (2017). A review on potential livestock feed: Azolla. *Livestock Research International*, 5(1),-1-9. <https://doi.www.jakraya.com/journa/lri>
- [9] Rana, D., Katoch, S., Mane, B. G., Rani, D., & Sankhyan, V. (2017). Carcass characteristic and physico-chemical properties of broiler chicken meat supplemented with Azolla pinnata. *Journal of Animal Research*, 7(6), 1035-1041. <https://doi.10.5958/2277-940X.2017.00155.3>
- [10] Joysowal, M., Aziz, A., Mondal, A., Singh, S. M., Boda, S. S., Chirwatkar, B., & Chhaba, B. (2018). Effect of Azolla (Azolla pinnata) feed on the growth of broiler chicken. *Journal of Entomology and Zoology Studies*, 6(3), 391-393. <https://2u.pw/L0vHW729>
- [11] Noor Nawaz, A. S., Syed, J., Dileep, N., Rakesh, K. N., & Prashith Kekuda, T. R. (2014). Antioxidant activity of Azolla pinnata and Azolla rubra–A comparative study. *Sch Acad J Biosci*, 2(10), 719-23. www.saspublisher.com
- [12] Al-Mawla, Muhammad Ibrahim Mustafa, The effect of replacing soybean meal with the addition of the xylanase enzyme and the probiotic on the productive and physiological performance of broilers. (2023) Doctoral thesis. College of Agriculture and Forestry / University of Mosul.
- [13] Roque, B. M., Jadrn, A., & Kebreab, E. (2017). Role of exogenous enzyme supplementation to improve

- nutrition and health of ruminants. Broadening Horizons, 41, 1-10. www.feedipedia.org
- [14] National Research Council, & Subcommittee on Poultry Nutrition. (1994). *Nutrient requirements of poultry: 1994*. National Academies Press. <https://2u.pw/7oySJ1g>
- [15] Anonymous. (2003). Statistical Analysis System User's Guide. (Version 9.1.3). SAS Institute Inc., Cary North Carolina, U.S.A.
- [16] Duncan, D. B. (1955). Multiple range and multiple F tests. *biometrics*, 11(1), 1-42
<http://dx.doi.org/10.2307/3001478>
- [17] Ward, N. E. (2021). Debranching enzymes in corn/soybean meal-based poultry feeds: a review. *Poultry Science*, 100(2), 765-775. <https://doi.org/10.1016/j.psj.2020.10.074>
- [18] Bedford, M. R., Partridge, G. G., Hruby, M., & Walk, C. L. (Eds.). (2022). *Enzymes in farm animal nutrition*. Cabi. <https://2u.pw/Mz8R7oeu>
- [19] Shambhvi, Katoch, S., Chauhan, P., & Mane, B. G. (2021). Effect of feeding *Azolla pinnata* in combination with direct-fed microbial on broiler performance. *Tropical Animal Health and Production*, 53, 1-9. <https://doi.org/10.1007/s11250-020-02437-w>.
- [20] Ibrahim, S., Ateya, A., & Abdo, M. (2024). Economic Evaluation of Using *Azolla* on Growth Performance of Broiler Chickens: Gene Expression Impact. *Egyptian Journal of Veterinary Sciences*, 55(1), 33-47. <https://doi.org/10.21608/ejvs.2023.215995.1519>
- [21] Abdelatty A M, Mandouh M I, Al-Mokaddem A K, Mansour H A, Khalil H M A, Elolimy A A, & Bionaz M. (2020). Influence of level of inclusion of *Azolla* leaf meal on growth performance, meat quality and skeletal muscle p70S6 kinase α abundance in broiler chickens. *animal*, 14(11), 2423-2432. <https://doi.org/10.1017/S1751731120001421>
- [22] Khan, F. U., Rafiullah, R., Lionel, K. I. N. K. P. E., Hassan, S. U., Ahmed, H. Q., Shuaib, M., & Assogba, C. M. (2023). Effect of (*Azolla Pinnata*) Meal Supplementation on Growth Performance, Carcass Traits, and Gut Histomorphology in Commercial Broilers
<http://dx.doi.org/10.21203/rs.3.rs-3712500/v1>
- [23] Samad, F. A., Idris, L. H., Abu Hassim, H., Goh, Y. M., & Loh, T. C. (2020). Effects of *Azolla* spp. as feed ingredient on the growth performance and nutrient digestibility of broiler chicken. *Journal of animal physiology and animal nutrition*, 104(6), 1704-1711. <https://doi.org/10.1111/jpn.13345>
- [24] Al-Hussaini N A.(2023). Substituting the *Azolla* Plant to Replace Part of the Protein in Broiler Diets and its Effect on Productive Performance. A Thesis. College of Agriculture and Forestry University of Mosul.
- [25] Yegani, M., & Korver, D. R. (2013). Effects of corn source and exogenous enzymes on growth performance and nutrient digestibility in broiler chickens. *Poultry Science*, 92(5), 1208-1220. <https://doi.org/10.3382/ps.2012-02390>
- [26] Van Hoeck, V., Wu, D., Somers, I., Wealleans, A., Vasanthakumari, B. L., Sanchez, A. G., & Morisset, D. (2021). Xylanase impact beyond performance: A prebiotic approach in broiler chickens. *Journal of Applied Poultry Research*, 30(4), 100193. <https://doi.org/10.1371/journal.pone.0257681>
- [27] Saleh, A. A., El-Far, A. H., Abdel-Latif, M. A., Emam, M. A., Ghanem, R., & Abd El-Hamid, H. S. (2018). Exogenous dietary enzyme formulations improve growth performance of broiler chickens fed a low-energy diet targeting the intestinal nutrient transporter genes. *PLoS One*, 13(5), e0198085. <https://doi.org/10.1371/journal.pone.0198085>
- [28] Aderibigbe, A., Cowieson, A. J., Sorbara, J. O., & Adeola, O. (2020). Growth phase and dietary α -amylase supplementation effects on nutrient digestibility and feedback enzyme secretion in broiler chickens. *Poultry Science*, 99(12), 6867-6876. <https://doi.org/10.1016/j.psj.2020.09.007>
- [29] Aderibigbe, A., Cowieson, A., Sorbara, J. O., & Adeola, O. (2020). Intestinal starch and energy digestibility in broiler chickens fed diets supplemented with α -amylase. *Poultry Science*, 99(11), 5907-5914. <https://doi.org/10.1016/j.psj.2020.08.036>
- [30] Karunaratne, N. D., Classen, H. L., Ames, N. P., Bedford, M. R., & Newkirk, R. W. (2021). Effects of hullless barley and exogenous beta-glucanase levels on ileal digesta soluble beta-glucan molecular weight, digestive tract characteristics, and performance of broiler chickens. *Poultry Science*, 100(3), 100967
<https://doi.org/10.1016/j.psj.2022.101735>
- [31] Oliveira, T. M. M. D., Nunes, R. V., Eyng, C., Berwanger, E., & Bayerle, D. F. (2016). Sunflower meal and exogenous enzymes in initial diets for broilers. *Revista Caatinga*, 29(4), 996-1005
<https://doi.org/10.1590/1983-21252016v29n425rc>
- [32] Singh, A. K., Mishra, B., Bedford, M. R., & Jha, R. (2021). Effects of supplemental xylanase and xylooligosaccharides on production performance and gut health variables of broiler chickens. *Journal of Animal Science and Biotechnology*, 12, 1-15.
<https://doi.org/10.1186/s40104-021-00617-8>
- [33] Kamel, E. R., & Hamed, E. (2021). Effect of dried *azolla* on growth performance, hematological, biochemical, antioxidant parameters, and economic efficiency of broiler chickens. *Adv. Anim. Vet. Sci*, 9(11), 1886-1894. <https://doi.org/10.17582/journal.aavs/2021/9.11.1886.1894>
- [34] Woyengo, T. A., Slominski, B. A., & Jones, R. O. (2010). Growth performance and nutrient utilization of broiler chickens fed diets supplemented with phytase alone or in combination with citric acid and multicarbohydrase. *Poultry science*, 89(10), 2221-2229. <https://doi.org/10.3382/ps.2010-00832>

- [35] Ribeiro, T., Lordelo, M. M. S., Ponte, P. I. P., Maças, B., Prates, J. A. M., Fontes, M. A., ... & Fontes, C. M. G. A. (2011). Levels of endogenous β -glucanase activity in barley affect the efficacy of exogenous enzymes used to supplement barley-based diets for poultry. *Poultry Science*, 90(6), 1245-1256. <https://doi.org/10.3382/ps.2010-01218>.
- [36] Bassi, L. S., Hejdysz, M., Pruszyńska-Oszmalek, E., Wolc, A., Cowieson, A. J., Sorbara, J. O. B., & Kaczmarek, S. A. (2023). The effect of amylase supplementation on individual variation, growth performance, and starch digestibility in broiler chickens. *Poultry Science*, 102(4), 102563. <https://doi.org/10.1016/j.psj.2023.102563>
- [37] Najim, Y. S., Mohammed, T. H., & Al-Khalani, F. M. H. (2020). The effect of the use of different levels of azolla to male broilers' diets in the production and physiological performance and economic feasibility. *Biochemical & Cellular Archives*, 20(1). <https://doi.org/10.35124/bca.2020.20.1.573>
- [38] Lavania, P., & Bairwa, K. C. (2023). Growth performance of Pratapdhan chicks supplemented with fresh Azolla in backyard system under the arid condition of Rajasthan. <https://doi.org/10.51966/jvas.2023.54.1.30-34>
- [39] Rengma, D. J., Savino, N., & Vidyarthi, V. K. (2019). Effect of dietary inclusion of Azolla powder on performance of broiler chicken. *Livestock Research International*, 7(02), 144-150. www.jakraya.com/journa/lr
- [40] Katoch, S., Rana, D., Chandel, S., & Sankhyan, V. (2017). Biological evaluation of dried Azolla in combination with carbohydrates in broiler bird <https://doi.org/10.5958/0974-8180.2017.00004.6>
- [41] Munyaka, P. M., Nandha, N. K., Kiarie, E., Nyachoti, C. M., & Khafipour, E. (2016). Impact of combined β -glucanase and xylanase enzymes on growth performance, nutrients utilization and gut microbiota in broiler chickens fed corn or wheat-based diets. *Poultry Science*, 95(3), 528-540. <https://doi.org/10.3382/ps/pev333>
- [42] Mathlouthi, N., Ballet, N., & Larbier, M. (2011). Influence of beta-glucanase supplementation on growth performances and digestive organs weights of broiler chickens fed corn, wheat and barley-based diet. *Int J Poult Sci*, 10(2), 157-159. <https://doi.org/10.3923/ijps.2011.157.159>

Table 4. Effect of added levels of enzyme % of the ratio containing Azolla on an average weekly live body weight of broilers, g/bird.

Character Treatment	Body weight (g)					
	Weeks					
	First	Second	Third	Fourth	Fifth	Sixth
T1	141.00 a	340.17 a	712.43 c	1304.33b	1894.67b	2429.70 b
Control	±2.28	± 9.61	±11.50	±40.57	±15.81	±20.72
T2	138.00 a	338.2 a	717.14 bc	1321.0 b	1911.11b	2437.0a
Azolla %7	±0.51	±9.48	± 5.93	± 2.91	± 25.145	± 35.0
T3	144.8 a	351.30a	722.47 bc	1319.0 b	1945.67b	2548.33a
Azolla7%	±0.51	±9.47	±15.93	±2.91	±25.145	±35.04
+0.30% Enzyme						
T4	149.00 a	359.73 a	754.3ab	1353.9ab	1954.6ab	2552.0a
7%Azolla +0.35%	±2.95	± 2.25	±6.43	±33.36	±51.25	±53.38
Enzyme						
T5	154.0 a	363.37 a	785.60a	1431.83a	2041.23a	0 2630.6 a
%7Azolla	±1.07	±5.03	±13.37	±26.18	±49.37	± 71.72
+ 0.40% enzyme						

Different letters vertically indicate the presence of significant differences ($P \leq 0.05$) between the means \pm S.E

Table 5. Effect of added levels of enzyme % of the ratio containing Azolla on a weight gain of broilers g/bird.

Character		Weight gain (g)					Total weight gain
Treatment	First	Second	Third	Fourth	Fifth	Sixth	1-6 weeks
T1	99.0 a	199.17a	372.26b	591.9b	590.34a	535.7b	2388.37 b
Control	±2.28	± 11.78	± 2.54	±9.11	±24.77	±14.93	±20.73
T2	95.9a	200.2a	378.94ab	604.17b	589.80a	525.89b	2394.9 b
Azolla %7	±2.5	± 9.04	±19.742	±14.25	±2.09	±11.50	±1.83
T3	102.6a	207.3a	371.17ab	597.43b	625.77a	602.66a	2506.93 a
Azolla7%+ 0.30% Enzyme	±2.50	±9.04	±19.72	±14.25	±2.06	±11.50	±8.33
T4	107 ab	210.73a	394.3a	594.87b	600.70a	597.4a	2505a
Azolla 7%+ 0.35% Enzyme	±2.95	±1.68	±7.21	± 27.14	±17.88	±13.78	±3.39
T5	112.4a	209.37a	422.23a	646.23a	609.4a	589.37a	2589.0 a
Azolla7% + 0.40% enzyme	± 3.4	± 5.69	±10.5	±16.44	±24.48	±23.42	±71.15

Different letters vertically indicate the presence of significant differences ($P \leq 0.05$) between the means \pm S.E

Table 6. Effect of added levels of enzyme to ratio containing Azolla on the growth rate of broiler

Treatment	Character	Relative growth rate%						relative growth (1-6)
		Weeks						
		First	Second	Third	Fourth	Fifth	Sixth	
T1		107.7 a	82.79 a	70.73a	58.70a	36.91a	24.74 a	63.60a
Control		±6.3	±9.2	±3.7	±4.4	±7.13	±3.12	±4.65
T2		106.50 a	84.08a	71.81a	59.28a	36.49a	24.19a	63.72a
Azolla %7		±5.4	±5.3	±5.7	±9.3	5.5±	5.33±	±7.12
T3		109.73a	83.25a	69.13a	58.5a	38.33a	26.82a	64.29a
Azolla7%+ 0.30% Enzyme		±5.4	±5.3	±5.7	± 9.35	±5.5	±5.33	±7.12
T4		112.04a	86.79a	70.84a	56.88a	36.31a	26.51a	64.90a
Azolla 7%+ 0.35% Enzyme		±8.0	±7.1	±5.9	±11.6	±6.04	±4.75	±8.13
T5		114.93a	80.93a	73.50a	58.29a	35.09a	27.59a	65.06a
Azolla7% + 0.40% enzyme		±11.1	±6.43	±9.3	±7.13	±11.3	±3.15	± 9.01

Different letters vertically indicate the presence of significant differences ($P \leq 0.05$) between the means \pm S.E

Table 7. Effect of added levels of enzyme to ratio containing Azolla on the feed intake of broiler

Treatments	Character	feed intake(g)					total feed consumed 1-6 weeks
		Weeks					
	First	Second	Third	Fourth	Fifth	Sixth	
T1	124.68a	304.75a	573.28a	1000.31a	1166.5a	1187.54a	4556.36a
Control	± 15.3	±2.45	±57.59	±35.10	±46.37	±40.52	92.238 ±
T2	119.88a	268.27a	591.15ab	924.38b	1037.38b	1009.71b	3568.40b
Azolla %7	± 3.51	± 8.67	±25.11	±41.30	±18.71	± 34.26	84.251±
T3	115.39a	284.41a	519.64b	900.00 b	1013.75b	1096.84b	3974.03b
7% Azolla ± 0.30% Enzyme	±3.38	±1.82	±22.17	±32.00	±12.08	±25.02	41.710±
T4	124.67a	292.0a	552.02b	905.99b	1021.19b	1105.19a	3901.24b
7%Azolla + 0.35% Enzyme	±2.62	±2.71	±31.15	±14.00	±15.36	±19.27	40.979±
T5	126.85a	270.96a	570.01b	990.67a	1047.48b	1113.91a	3968.33b
%7Azolla + 0.40% enzyme	± 2.76	±8.69	±5.11±	±41.30	±18.71	± 34.46	84.251±

Different letters vertically indicate the presence of significant differences ($P \leq 0.05$) between the means \pm S.E.

Table 8. Effect of added levels of an enzyme to ratio containing Azolla on the Feed conversion ratio (g feed/g weight gain) of broiler

Character	Feed conversion ratio (g feed/g weight gain)						Feed conversion ratio 1-6 w
	Treatment	First	Second	Third	Fourth	Fifth	
T1		1.21a	1.38a	1.54a	1.69a	1.97a	1.65a
Control		±0.04	±0.03	±0.02	± 0.05	± 0.08	± 0.03
T2		1.25a	1.34a	1.56a	1.53b	1.82ab	1.49b
Azolla %7		±0.01	± 0.03	±0.02	± 0.01	± 0.04	± 0.08
T3		1.22a	1.31a	1.40b	1.50b	1.62b	1.47b
7% Azolla ± 0.30% Enzyme		±0.03	±0.05	±0.03	± 0.04	± 0.02	± 0.02
T4		1.29a	1.39a	1.41b	1.52b	1.70b	1.50b
7%Azolla+ 0.35% Enzyme		±0.02	0.0±	±0.02	± 0.01	± 0.02	± 0.05
T5		1.30a	1.35a	1.39b	1.53b	1.72b	1.51b
%7Azolla + 0.40% enzyme		±0.01	0.027±	±0.01	± 0.01	± 0.04	± 0.08

Different letters vertically indicate the presence of significant differences ($P \leq 0.05$) between the means \pm S.E

Table 9. Effect of added levels of the enzyme to ratio containing Azolla on the feed consumption (g) of broiler

Indicators	1-5 weeks		1-6 weeks	
Treatment	Production index	Production factor	Production index	Production factor
T1	273.95	29.88	313.13	24.16
control				
T2	02.003	32.41	302.27	27.39
%7Azolla				
T3	343.15	38.63	400.05	33.11
+ 7%Azolla				
0.30% Enzyme				
T4	328.50	35.34	394.13	32.29
Azolla 7%+				
0.35% Enzyme				
T5	339.07	35.43	397.67	31.18
Azolla7% + 0.40% enzyme				