

Campylobacter jejuni

(2004/11/7 2004/8/21)

(*C.jejuni*) *Campylobacter jejuni*

2002

.2003

Preston *Campylobacter* blood-free medium

S

pleomorphic

% 1

° 25

° 42

Nalidixic acid

% 1.5

.Cephalothin

Isolation and Identification of *Campylobacter jejuni* from Diarrhoeal Cases of Children in Mosul City

Ghada A. Al-Tae

Amera M. Al-Rawe

Department of Biology

College of Science

Mosul University

ABSTRACT

The study include the isolation of *Campylobacter jejuni* (*C.jejuni*) from children (aging between after birth to two years) with watery or bloody diarrhea in the period

from September 2002 to February 2003. The bacterium was identified depending on succession in growth on selective medium: Preston Campylobacter blood-free medium and microscopical examination which revealed that it is gram negative pleomorphic, curved bacillus, S-shaped and gull-winged. Biochemical tests showed its positive reaction for catalase, oxidase and its ability to hydrolyse hippurate and indoxyl acetate but it was negative for urease and lipase. This bacterium can grow at 42° C while it failed to grow at 25° C and it tolerates 1% glycine while it can not grow in 1.5 % NaCl. Additionally the bacterium was sensitive to Nalidixic acid but resistant to Cephalothin.

Campylobacter

C. jejuni

.(Saenz et al., 2000)

gull-wing (S)

° 42 ° 37

H₂S

Nalidixic acid

Cephalothin

C. jejuni

C. jejuni sub *jejuni*

.(Collee et al.,1996; Blaser, 1999) sub *doylei*

C. jejuni

. *C. jejuni*

152

Cary-Blair transport media -

Preston Campylobacter Blood-free selective medium

° 42

H₂S IMViC
 Koneman et al., 1997; Prescott et al.,) Glycine
 .(1996

:

.(Koneman et al., 1997) IMViC

(13) (206)
C. jejuni (9) (% 6.3) *Campylobacter*
 Campylobacter (% 69.2) (% 4.36)
 (% 1.94) *C. coli* (4)
 .Campylobacter (% 30.8)
 (Church et al., 1995) *C. jejuni*
 294 % 17 *C. jejuni* 51
 Salih % 10 *C. jejuni*
 1994 Al-Saad
 % 6.2 (Saenz et al., 2000)
 537 8616
C. coli *C. jejuni*
 Goossens 1991 Fang .% 0.67
 1995 Giesendorf

Church et al.,) *E. coli* (1)
 .(Bryant et al., 1989) (1995)

.(Ahren et al., 1990)

:1

%				
6.32	13	Campylobacter spp.	194	152
19.48	40	<i>E. coli</i>		
16.50	34	Proteus		
8.73	18	Pseudomonas		
7.76	16	Enterobacter		
7.28	15	<i>E. faecalis</i>		
6.79	14	Klebsiella		
5.82	12	Citrobacter		
4.36	9	Salmonella		
3.39	7	Providencia		
3.39	7	Morganella		
1.94	4	Staph. Spp.		
1.45	3	<i>Y. enterocolitica</i>		
0.48	1	Serratia		
0.48	1	Shigella		
5.82	12	Candida spp.	12	
100	206		206	

:

Preston *Campylobacter* blood –free medium *C. jejuni*

Koneman et al., 1997; Baron and Fingold,)
 .(1) (1990; Merino et al., 1986

C. jejuni .(2)

curve = campylo

gull-wing

(s)

rod = bacter

.(3)

shape

C.jejuni

(2)

H₂S

<i>C.jejuni</i>		:2
	Tests	
+	Catalase test	
+	Oxidase test	
+	Hippurate hydrolysis	
+	Indoxyl acetate hydrolysis	
-	Urease test	
-	Lipase test	
-	()	
+	Motility test	
-	: ° 25	
+		
+		
-	TSI H ₂ S	
-	: %1.5	
+		
S	: Nalidixic acid	
R		
+	Cephalothin	
-	IMViC	
Positive : +		Negative : -
Sensitive : S		Resistant : R

° 25 %1.5

° 42 ° 37
thermophiles

%1

1987 ,)


.(Koneman et al., 1997; Holt et al.,1994;

C. jejuni
(4)

.1980 Harvey .(5)

:

Hippuricas



Hippuric acid		Benzoic acid	Glycocoll (glycine)
66-44			
1975	Hwang	1974	Facklam

° 37 %1 3 0.5 B

Glycine
10


.(5)
C. jejuni

(IAH)


10 (6)

IAH .


Campylobacter




2




1




4



3



6



5

<i>C.jejuni</i>	(2)	<i>C.jejuni</i>	(1)
	(4)	<i>C.jejuni</i>	(3)
	(6)		(5)

C. jejuni

Nalidixic acid

Cephalothin

Nalidixic

acid

Cephalothin

.(Melo and Pechere, 1990)

*C. jejuni**C. jejuni*. Subsp. *jejuni*

Cephalosporins

C. jejuni subsp. *dolei*

.(Koneman et al., 1997)

.1987

- Ahren, C.M., Jertborn, M., Herclik, L., Kaijser, B. and Svennerholm, A.M., 1990. Infection with bacterial enteropathogens in Swedish travellers to South-East Asia- a prospective study. *Epidemiol. Infect.*, Vol.105, pp.325-333.
- Baron, J.E., Peterson, L.R. and Finegold, S.M., 1994. *Bailey and Scott's Diagnostic Microbiology*. 9th. ed., Mosby, USA.
- Baron, E.J. and Finegold, S.M., 1990. *Bailey and Scott's Diagnostic Microbiology*. 8th ed., C.V. Mosby Company, USA.
- Blaser, M., 1999. *Campylobacter jejuni* and related species , pp. 2276-2285. In: G.L. Mandell, J.E. Bennett and R. Dolin (ed.): *Principles and Practices of Infectious Disease*, vol. 2. Churchill Livingstone, Philadelphia, Pa.
- Boton, F.J., Hutchison, D.N. and Coates, D., 1984. Blood-Free selective medium for isolation of *Campylobacter jejuni* from feces .*J. Clin. Microbiol.*, Vol.19, pp.169-171.
- Bryant, H.E., Athar, M.A. and Pai, C.H., 1989. Risk factors for *Escherichia coli* 0157:H7 infection in an urban community. *J. Infect. Dis.*, Vol.160, pp.858-864.
- Church, D.L., Gisele, C.R.T., Amin, K.M.D., T. Jadavji, M.D. and C. Trevenen, M.D., 1995. Practice Guidelines for ordering stool cultures in a pediatric population . *Am. Soc. Clin. Pathol.*, USA.
- Collee, J.G., Fraser, A.G., Marmion, B.P. and Simmons, A., 1996. *Mackie and McCartney Practical Medical Microbiology*. 14th ed, Churchill Livingston Inc., New York.

- Faklam, R.R., Padula, J.F., Thacker, L.G. and Sconyers, B.J., 1974. Presumptive identification of group A, B and D Streptococci. *Appl. Microbiol.*, Vol.27, pp.107-113.
- Fang, G., Araujo, V. and Guerrant, R.L., 1991. Enteric infections associated with exposure to animals or animal products. *Dis. Cli. NA*, Vol.5, pp.681-701.
- Goossens, H. and Giesendorf, A.J., 1995. Investigation of an outbreak of *Campylobacter* in day care. *Dis.*, Vol.172, pp.1298-1305.
- Harvey, D. and Leach, J., 1998. Analysis of coccal cell formation by *Campylobacter jejuni* using continuous culture techniques and the importance of oxidative stress. *J. Appl. Microbiol.*, Vol.85, pp.398-400.
- Holt, J.G., Krieg, N.R., Sneath, P.H.A., Staley, J.T. and Williams, S.T., 1994. *Bergey's Manual of determinative Bacteriology*. 9th. ed. Williams and Wilkins Comp. USA. Baltimore.
- Hwang, M. and Ederrer, G.M., 1975. Rapid hippurate hydrolysis method to presumptive identification of group B streptococci. *J. Clin. Microbiol.*, pp.114-115.
- Karamali, M.A., Simor, A.E., Roscoe, M., Fleming, P.C., Smith, S.S. and Lane, J., 1986. Evaltion of a Blood-Free, Charcoal-Based, Selective Medium for the isolation of *Campylobacter* organisms from feces. *J. Clin. Microbiol.*, Vol.23, pp.456-459.
- Koneman, E.W., Allen, S.D., Janda, W.M., Schreckenberger, P.C., Winn, W.C., 1997. *Color Atlas and Textbook of Diagnostic Microbiology* 5th ed. Lippincott-Raven Publishers, Philadelphia, USA.
- Lennette, E.H., Spaulding, E.H. and Truant, J.P., 1974. *Manual of Clinical Microbiology* 2nd ed. American Society for Microbiology, Washington, D.C., 263 p..
- Melo, M.A. and Pechere, J.C., 1990. Identification of *Campylobacter jejuni* surface protein that bind to eucaryotic cells in vitro. *J. Infect. Immun.*, Vol.58(6), pp.1749-1756.1986.
- Merino, F.J., Agull, A. and Villasante, P.A., 1986. Comparative efficacy of seven selective media for isolating *Campylobacter jejuni*. *J. Clin. Microbiol.*, Vol.24, pp.451-452.
- Prescott, L.M., Harley, J.P. and Klein, D.A., 1996. *Microbiology*. 3rd ed., WMC.Brown Communication, Inc., Iowa, USA.
- Saenz, Y., Zarazaga, M., Lantero, M., Gastanares, M.J., Baquero, F. and Torres, C., 2000. Antibiotic resistance in *Campylobacter* strains isolated from animals, foods and humans in Spain, pp.1997-1998.
- Salih, D.S. and Al-Saad, M.R., 1994. Isolation and Identification of thermophilic *Campylobacters* from diarrhoeal chilidren in Baghdad. *J. of IAS*, Vol.7(2).